



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:42 PM UTC

PDB ID : 1KZJ / pdb_00001kzj
Title : Crystal Structure of EcTS W80G/dUMP/CB3717 Complex
Authors : Fritz, T.A.; Liu, L.; Finer-Moore, J.S.; Stroud, R.M.
Deposited on : 2002-02-06
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	58% 37% 5%
1	B	264	59% 34% 7%
1	C	264	61% 31% 7%
1	D	264	3% 58% 38% 5%
1	E	264	2% 58% 36% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	264	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a green segment representing 61%, a yellow segment representing 34%, and a small orange segment representing 5%.

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2143	1366	370	395	12			
1	B	264	Total	C	N	O	S	0	0	0
			2143	1366	370	395	12			
1	C	264	Total	C	N	O	S	0	0	0
			2143	1366	370	395	12			
1	D	264	Total	C	N	O	S	0	0	0
			2143	1366	370	395	12			
1	E	261	Total	C	N	O	S	0	0	0
			2122	1352	367	391	12			
1	F	264	Total	C	N	O	S	0	0	0
			2143	1366	370	395	12			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	CXM	MET	modified residue	UNP P0A884
A	80	GLY	TRP	engineered mutation	UNP P0A884
B	1	CXM	MET	modified residue	UNP P0A884
B	80	GLY	TRP	engineered mutation	UNP P0A884
C	1	CXM	MET	modified residue	UNP P0A884
C	80	GLY	TRP	engineered mutation	UNP P0A884
D	1	CXM	MET	modified residue	UNP P0A884
D	80	GLY	TRP	engineered mutation	UNP P0A884
E	1	CXM	MET	modified residue	UNP P0A884
E	80	GLY	TRP	engineered mutation	UNP P0A884
F	1	CXM	MET	modified residue	UNP P0A884
F	80	GLY	TRP	engineered mutation	UNP P0A884

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (CCD ID: UMP) (formula: C₉H₁₃N₂O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	C	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	D	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	E	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	F	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is 10-PROPARGYL-5,8-DIDEAZAFOLIC ACID (CCD ID: CB3) (formula: $C_{24}H_{23}N_5O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 35	C 24	N 5	O 6	0	0
3	B	1	Total 35	C 24	N 5	O 6	0	0
3	C	1	Total 35	C 24	N 5	O 6	0	0
3	D	1	Total 35	C 24	N 5	O 6	0	0
3	E	1	Total 35	C 24	N 5	O 6	0	0
3	F	1	Total 35	C 24	N 5	O 6	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	54	Total O 54 54	0	0
4	B	54	Total O 54 54	0	0
4	C	62	Total O 62 62	0	0
4	D	34	Total O 34 34	0	0
4	E	36	Total O 36 36	0	0
4	F	52	Total O 52 52	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Thymidylate synthase




• Molecule 1: Thymidylate synthase




• Molecule 1: Thymidylate synthase






• Molecule 1: Thymidylate synthase



• Molecule 1: Thymidylate synthase



• Molecule 1: Thymidylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.55Å 84.57Å 111.52Å 90.00° 111.37° 90.00°	Depositor
Resolution (Å)	25.35 – 2.60 25.35 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.2 (25.35-2.60) 98.2 (25.35-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 2.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.215 , 0.258 0.213 , 0.256	Depositor DCC
R_{free} test set	3545 reflections (7.06%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13459	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4247e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UMP, CXM, CB3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2190	0.95	10/2972 (0.3%)
1	B	0.45	0/2190	0.95	12/2972 (0.4%)
1	C	0.44	0/2190	0.91	5/2972 (0.2%)
1	D	0.44	0/2190	0.91	11/2972 (0.4%)
1	E	0.46	0/2169	0.95	8/2944 (0.3%)
1	F	0.46	0/2190	0.95	7/2972 (0.2%)
All	All	0.45	0/13119	0.94	53/17804 (0.3%)

There are no bond length outliers.

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	170	VAL	N-CA-C	7.05	117.77	110.36
1	F	24	THR	N-CA-C	6.98	118.54	111.07
1	D	168	CYS	N-CA-C	6.84	119.51	107.61
1	A	27	LEU	N-CA-C	-6.76	97.26	108.34
1	B	170	VAL	N-CA-C	6.63	118.18	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2143	0	2072	95	0
1	B	2143	0	2072	77	0
1	C	2143	0	2072	83	0
1	D	2143	0	2073	88	0
1	E	2122	0	2048	93	0
1	F	2143	0	2072	82	0
2	A	20	0	10	0	0
2	B	20	0	10	1	0
2	C	20	0	10	1	0
2	D	20	0	11	1	0
2	E	20	0	11	1	0
2	F	20	0	10	2	0
3	A	35	0	21	2	0
3	B	35	0	21	2	0
3	C	35	0	21	3	0
3	D	35	0	21	6	0
3	E	35	0	21	5	0
3	F	35	0	21	1	0
4	A	54	0	0	2	0
4	B	54	0	0	4	0
4	C	62	0	0	6	0
4	D	34	0	0	1	0
4	E	36	0	0	1	0
4	F	52	0	0	3	0
All	All	13459	0	12597	509	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 509 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:183:LEU:HD22	1:F:187:MET:HE2	1.36	1.07
1:C:127:ARG:HH21	1:C:127:ARG:HG3	1.27	0.99
1:D:83:TRP:HH2	3:D:307:CB3:H8	1.25	0.98
1:B:233:LYS:HD2	1:B:248:GLU:HG2	1.44	0.97
1:A:59:LEU:HD23	1:A:183:LEU:HD23	1.45	0.96

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/264 (99%)	246 (94%)	15 (6%)	1 (0%)	30	51
1	B	262/264 (99%)	240 (92%)	18 (7%)	4 (2%)	8	18
1	C	262/264 (99%)	244 (93%)	15 (6%)	3 (1%)	11	25
1	D	262/264 (99%)	240 (92%)	20 (8%)	2 (1%)	16	34
1	E	259/264 (98%)	235 (91%)	20 (8%)	4 (2%)	8	18
1	F	262/264 (99%)	244 (93%)	15 (6%)	3 (1%)	11	25
All	All	1569/1584 (99%)	1449 (92%)	103 (7%)	17 (1%)	11	25

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	94	TYR
1	A	94	TYR
1	B	23	GLY
1	B	79	ILE
1	E	94	TYR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	217 (94%)	14 (6%)	17	37
1	B	231/231 (100%)	214 (93%)	17 (7%)	13	29
1	C	231/231 (100%)	212 (92%)	19 (8%)	10	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	231/231 (100%)	216 (94%)	15 (6%)	15	35
1	E	229/231 (99%)	216 (94%)	13 (6%)	18	40
1	F	231/231 (100%)	218 (94%)	13 (6%)	19	40
All	All	1384/1386 (100%)	1293 (93%)	91 (7%)	15	34

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	172	LEU
1	E	134	ASN
1	D	215	GLN
1	E	17	GLN
1	E	220	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 70 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	117	ASN
1	E	215	GLN
1	F	117	ASN
1	B	134	ASN
1	B	121	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CXM	B	1	1	9,10,11	1.34	2 (22%)	9,11,13	1.24	1 (11%)
1	CXM	D	1	1	9,10,11	1.30	1 (11%)	9,11,13	1.45	1 (11%)
1	CXM	E	1	1	9,10,11	1.07	0	9,11,13	1.29	1 (11%)
1	CXM	F	1	1	9,10,11	1.61	2 (22%)	9,11,13	1.14	1 (11%)
1	CXM	A	1	1	9,10,11	1.06	0	9,11,13	1.36	1 (11%)
1	CXM	C	1	1	9,10,11	1.07	1 (11%)	9,11,13	1.51	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	B	1	1	-	0/9/10/12	-
1	CXM	D	1	1	-	4/9/10/12	-
1	CXM	E	1	1	-	2/9/10/12	-
1	CXM	F	1	1	-	4/9/10/12	-
1	CXM	A	1	1	-	1/9/10/12	-
1	CXM	C	1	1	-	2/9/10/12	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	1	CXM	CA-N	3.05	1.51	1.46
1	B	1	CXM	CA-N	2.80	1.50	1.46
1	F	1	CXM	CN-N	2.77	1.40	1.35
1	D	1	CXM	CA-N	2.32	1.49	1.46
1	B	1	CXM	ON1-CN	2.12	1.25	1.21

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1	CXM	ON1-CN-N	-4.05	118.22	124.86
1	C	1	CXM	ON1-CN-N	-3.83	118.58	124.86
1	A	1	CXM	ON1-CN-N	-3.72	118.75	124.86
1	E	1	CXM	ON1-CN-N	-3.56	119.03	124.86
1	B	1	CXM	ON1-CN-N	-2.92	120.06	124.86

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	1	CXM	ON1-CN-N-CA
1	D	1	CXM	ON2-CN-N-CA
1	D	1	CXM	O-C-CA-CB
1	F	1	CXM	N-CA-CB-CG
1	F	1	CXM	C-CA-CB-CG

There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	1	CXM	1	0
1	D	1	CXM	3	0
1	E	1	CXM	5	0
1	F	1	CXM	1	0
1	A	1	CXM	4	0
1	C	1	CXM	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UMP	F	310	1	21,21,21	3.59	5 (23%)	30,31,31	2.86	12 (40%)
3	CB3	E	309	-	37,37,37	1.51	4 (10%)	50,51,51	2.31	7 (14%)
2	UMP	D	306	-	21,21,21	2.84	4 (19%)	30,31,31	3.03	8 (26%)
3	CB3	A	301	-	37,37,37	1.41	5 (13%)	50,51,51	2.40	10 (20%)
3	CB3	F	311	-	37,37,37	1.88	7 (18%)	50,51,51	2.91	13 (26%)
3	CB3	C	305	-	37,37,37	1.44	6 (16%)	50,51,51	2.77	15 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UMP	A	300	1	21,21,21	3.27	4 (19%)	30,31,31	3.13	9 (30%)
3	CB3	B	303	-	37,37,37	1.47	5 (13%)	50,51,51	1.77	6 (12%)
3	CB3	D	307	-	37,37,37	1.70	9 (24%)	50,51,51	1.78	10 (20%)
2	UMP	E	308	-	21,21,21	3.29	5 (23%)	30,31,31	2.78	9 (30%)
2	UMP	B	302	1	21,21,21	3.41	6 (28%)	30,31,31	3.32	12 (40%)
2	UMP	C	304	1	21,21,21	3.29	5 (23%)	30,31,31	2.95	11 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	F	310	1	-	1/10/22/22	0/2/2/2
3	CB3	E	309	-	-	6/27/28/28	0/3/3/3
2	UMP	D	306	-	-	0/10/22/22	0/2/2/2
3	CB3	A	301	-	-	3/27/28/28	0/3/3/3
3	CB3	F	311	-	-	5/27/28/28	0/3/3/3
3	CB3	C	305	-	-	8/27/28/28	0/3/3/3
2	UMP	A	300	1	-	0/10/22/22	0/2/2/2
3	CB3	B	303	-	-	5/27/28/28	0/3/3/3
3	CB3	D	307	-	-	2/27/28/28	0/3/3/3
2	UMP	E	308	-	-	0/10/22/22	0/2/2/2
2	UMP	B	302	1	-	3/10/22/22	0/2/2/2
2	UMP	C	304	1	-	0/10/22/22	0/2/2/2

The worst 5 of 65 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	UMP	O4-C4	10.21	1.44	1.24
2	C	304	UMP	C6-C5	9.79	1.57	1.35
2	E	308	UMP	O4-C4	9.46	1.43	1.24
2	B	302	UMP	C6-C5	9.10	1.56	1.35
2	F	310	UMP	O4-C4	9.06	1.42	1.24

The worst 5 of 122 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	311	CB3	CP1-N10-C9	-14.96	102.12	117.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	305	CB3	CP1-N10-C9	-12.17	104.93	117.19
3	E	309	CB3	CP2-CP1-N10	-11.30	103.02	113.45
3	A	301	CB3	CP2-CP1-N10	-10.74	103.54	113.45
2	A	300	UMP	C6-C5-C4	-9.46	107.45	119.53

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	302	UMP	C5'-O5'-P-OP2
2	B	302	UMP	C5'-O5'-P-OP3
3	F	311	CB3	C6-C9-N10-C14
3	E	309	CB3	N-CA-CB-CG
3	B	303	CB3	C6-C9-N10-C14


There are no ring outliers.



11 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	310	UMP	2	0
3	E	309	CB3	5	0
2	D	306	UMP	1	0
3	A	301	CB3	2	0
3	F	311	CB3	1	0
3	C	305	CB3	3	0
3	B	303	CB3	2	0
3	D	307	CB3	6	0
2	E	308	UMP	1	0
2	B	302	UMP	1	0
2	C	304	UMP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.









Ligand CB3 C 305




Ligand UMP A 300







Ligand CB3 D 307



Ligand UMP E 308





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/264 (99%)	-0.36	0 100 100	3, 16, 29, 39	0
1	B	263/264 (99%)	-0.25	1 (0%) 88 86	4, 17, 36, 45	0
1	C	263/264 (99%)	-0.25	0 100 100	3, 19, 33, 50	0
1	D	263/264 (99%)	-0.06	9 (3%) 48 42	5, 22, 42, 62	0
1	E	260/264 (98%)	-0.04	4 (1%) 72 68	6, 22, 43, 60	0
1	F	263/264 (99%)	-0.22	0 100 100	7, 20, 31, 44	0
All	All	1575/1584 (99%)	-0.20	14 (0%) 81 78	3, 19, 37, 62	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	22	THR	3.8
1	E	261	PRO	3.5
1	D	262	VAL	3.5
1	E	22	THR	3.2
1	D	263	ALA	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CXM	E	1	11/12	0.92	0.11	27,30,34,34	0
1	CXM	D	1	11/12	0.93	0.10	21,22,24,25	0
1	CXM	F	1	11/12	0.93	0.10	21,24,26,27	0
1	CXM	C	1	11/12	0.94	0.08	22,26,28,28	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	CXM	A	1	11/12	0.95	0.10	13,17,24,27	0
1	CXM	B	1	11/12	0.96	0.07	12,15,24,26	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CB3	E	309	35/35	0.69	0.27	67,78,92,92	0
3	CB3	D	307	35/35	0.76	0.16	43,49,63,64	0
3	CB3	B	303	35/35	0.82	0.14	24,31,50,52	0
3	CB3	F	311	35/35	0.85	0.13	18,26,47,48	0
3	CB3	C	305	35/35	0.88	0.12	10,24,51,53	0
3	CB3	A	301	35/35	0.88	0.12	9,24,36,37	0
2	UMP	E	308	20/20	0.94	0.09	12,15,21,23	0
2	UMP	D	306	20/20	0.95	0.09	18,21,23,24	0
2	UMP	B	302	20/20	0.95	0.09	16,17,21,21	0
2	UMP	F	310	20/20	0.96	0.07	12,15,20,20	0
2	UMP	C	304	20/20	0.97	0.07	12,14,17,18	0
2	UMP	A	300	20/20	0.98	0.06	4,7,11,12	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CB3 E 309:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around CB3 D 307:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CB3 B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CB3 F 311:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CB3 C 305:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around CB3 A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UMP E 308:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UMP D 306:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UMP B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UMP F 310:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UMP C 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UMP A 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.