



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 11:24 PM EDT

PDB ID : 1KZP / pdb_00001kzp
Title : PROTEIN FARNESYLTRANSFERASE COMPLEXED WITH A FARNE-
SYLATED K-RAS4B PEPTIDE PRODUCT
Authors : Long, S.B.; Casey, P.J.; Beese, L.S.
Deposited on : 2002-02-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	
2	B	437	
3	C	11	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACY	B	3003	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein Farnesyltransferase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2745	1746	481	513	5			

- Molecule 2 is a protein called Protein Farnesyltransferase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	407	Total	C	N	O	S	0	0	0
			3203	2050	550	580	23			


- Molecule 3 is a protein called Farnesylated K-Ras4B peptide product.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	5	Total	C	N	O	S	0	0	0
			35	22	5	6	2			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is ACETIC ACID (CCD ID: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is FARNESYL (CCD ID: FAR) (formula: $C_{15}H_{26}$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	C	0	0
			15	15		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	211	Total 211	O 211	0	0
7	B	201	Total 201	O 201	0	0
7	C	9	Total 9	O 9	0	0

i

- Molecule 1: Protein Farnesyltransferase alpha subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	171.24Å 171.24Å 69.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	90.5 (50.00-2.10) 96.4 (50.00-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.164 , 0.202 0.176 , 0.207	Depositor DCC
R_{free} test set	3770 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	26.9	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.028 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6428	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FAR, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2812	0.97	17/3817 (0.4%)
2	B	0.55	0/3292	0.97	9/4473 (0.2%)
3	C	0.69	0/34	0.83	0/43
All	All	0.54	0/6138	0.97	26/8333 (0.3%)

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	GLU	N-CA-C	8.73	120.48	110.97
1	A	83	GLY	CA-C-N	7.21	127.06	119.05
1	A	83	GLY	C-N-CA	7.21	127.06	119.05
2	B	121	ILE	N-CA-C	-7.16	103.80	110.53
2	B	351	LEU	N-CA-C	7.10	118.87	108.86
2	B	301	SER	N-CA-C	-6.98	103.68	111.71
1	A	348	LYS	N-CA-C	6.59	121.16	113.18
1	A	183	SER	N-CA-C	6.45	119.13	111.71
1	A	217	ASP	N-CA-C	6.39	121.92	111.37
2	B	375	HIS	N-CA-C	6.07	118.30	108.41
2	B	75	ARG	N-CA-C	6.02	118.33	111.11
1	A	85	SER	CA-C-N	6.02	126.03	119.89
1	A	85	SER	C-N-CA	6.02	126.03	119.89
1	A	150	GLU	N-CA-C	-5.75	105.09	111.36
1	A	249	GLY	N-CA-C	-5.65	104.45	112.81
1	A	371	SER	N-CA-C	5.62	118.13	111.33
2	B	192	LEU	N-CA-C	-5.62	102.97	110.55
1	A	305	SER	N-CA-C	5.59	119.84	113.19
1	A	331	GLU	N-CA-C	5.53	117.11	111.14
2	B	163	ILE	N-CA-C	-5.49	105.37	110.53
2	B	92	ALA	N-CA-C	-5.34	106.03	112.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	270	VAL	CA-C-N	5.31	125.62	119.47
1	A	270	VAL	C-N-CA	5.31	125.62	119.47
2	B	174	ARG	N-CA-C	5.23	116.98	111.28
1	A	162	GLN	CA-C-N	5.04	124.65	119.56
1	A	162	GLN	C-N-CA	5.04	124.65	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	0	2666	29	0
2	B	3203	0	3130	26	0
3	C	35	0	34	1	0
4	B	1	0	0	0	0
5	B	8	0	6	3	0
6	C	15	0	24	1	0
7	A	211	0	0	3	0
7	B	201	0	0	6	0
7	C	9	0	0	1	0
All	All	6428	0	5860	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:ILE:HG22	1:A:172:ARG:HH12	1.51	0.76
1:A:152:MET:O	1:A:156:ILE:HG23	1.88	0.72
2:B:358:ARG:HE	5:B:3003:ACY:H2	1.61	0.65
2:B:239:ILE:HB	2:B:252:THR:HA	1.79	0.64
2:B:308:LEU:HD13	2:B:330:PHE:HB3	1.80	0.64
2:B:30:HIS:HE1	7:B:1121:HOH:O	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:131:GLU:HG2	2:B:171:VAL:HG13	1.82	0.60
2:B:275:GLN:HG2	7:B:1213:HOH:O	2.02	0.59
2:B:358:ARG:CG	5:B:3003:ACY:H2	2.35	0.56
1:A:189:ILE:HD11	1:A:205:HIS:HD2	1.71	0.56
1:A:329:ASN:HB3	1:A:332:ASP:HB3	1.88	0.56
1:A:374:PRO:HG2	1:A:377:VAL:HG23	1.88	0.56
1:A:257:GLU:HG2	7:A:1342:HOH:O	2.07	0.54
2:B:358:ARG:HG3	5:B:3003:ACY:H2	1.91	0.52
1:A:303:GLN:O	1:A:307:SER:HB2	2.11	0.51
1:A:156:ILE:HG22	1:A:172:ARG:NH1	2.24	0.50
2:B:218:THR:HB	2:B:219:PRO:HD2	1.95	0.49
2:B:299:CYS:SG	6:C:12:FAR:H2	2.53	0.48
2:B:422:GLY:O	2:B:423:PHE:HB2	2.12	0.48
2:B:301:SER:O	2:B:305:ALA:HB3	2.14	0.48
2:B:333:GLN:HG3	2:B:387:MET:SD	2.53	0.47
1:A:109:ARG:HD3	7:A:1623:HOH:O	2.16	0.46
1:A:112:ARG:HA	1:A:140:LEU:CD2	2.46	0.46
1:A:112:ARG:HA	1:A:140:LEU:HD21	1.97	0.46
1:A:152:MET:O	1:A:156:ILE:CG2	2.60	0.46
1:A:75:ILE:HD11	1:A:115:ARG:NH2	2.31	0.46
1:A:205:HIS:O	1:A:209:VAL:HG23	2.16	0.45
1:A:208:TRP:NE1	1:A:212:GLU:HG3	2.32	0.45
1:A:302:LEU:HB3	1:A:306:HIS:HB2	1.98	0.45
1:A:312:ILE:HB	1:A:344:LEU:HD21	1.99	0.45
1:A:304:PRO:HG2	1:A:305:SER:H	1.82	0.45
1:A:311:LEU:C	1:A:311:LEU:HD23	2.42	0.45
1:A:77:PRO:HB2	1:A:101:ASP:HB3	1.98	0.44
1:A:96:PHE:HA	1:A:126:LEU:HD13	1.99	0.44
2:B:306:GLY:O	2:B:309:PRO:HD2	2.17	0.44
1:A:357:ARG:HH11	1:A:357:ARG:HG2	1.82	0.44
1:A:294:ASN:O	1:A:298:GLN:HG3	2.18	0.43
2:B:312:HIS:HD2	7:B:1273:HOH:O	2.00	0.43
2:B:51:VAL:O	2:B:55:ILE:HG12	2.18	0.43
2:B:253:PHE:HA	2:B:307:LEU:HD21	2.01	0.43
1:A:329:ASN:O	1:A:330:LYS:C	2.62	0.43
2:B:75:ARG:NH2	2:B:393:VAL:O	2.52	0.43
2:B:266:ARG:HG3	7:B:1450:HOH:O	2.19	0.42
2:B:259:LEU:HD12	2:B:268:LEU:HD11	2.01	0.42
2:B:83:LYS:HD3	7:B:1333:HOH:O	2.19	0.42
2:B:350:LEU:HB2	2:B:363:THR:HA	2.01	0.42
3:C:7:LYS:N	7:C:1559:HOH:O	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:ILE:CG2	1:A:172:ARG:HH12	2.27	0.42
2:B:137:ASP:HB2	7:B:1366:HOH:O	2.19	0.42
2:B:192:LEU:HD23	2:B:199:VAL:CG2	2.50	0.41
2:B:393:VAL:O	2:B:393:VAL:HG13	2.20	0.41
2:B:370:LEU:HD23	2:B:394:LEU:HD11	2.03	0.40
1:A:220:LEU:HD12	1:A:220:LEU:HA	1.83	0.40
1:A:156:ILE:O	1:A:160:GLU:HG2	2.21	0.40
1:A:308:SER:O	1:A:312:ILE:HG12	2.21	0.40
1:A:97:ARG:HD3	7:A:1472:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/377 (85%)	303 (94%)	17 (5%)	1 (0%)	36	36
2	B	405/437 (93%)	395 (98%)	10 (2%)	0	100	100
3	C	3/11 (27%)	3 (100%)	0	0	100	100
All	All	729/825 (88%)	701 (96%)	27 (4%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/338 (89%)	293 (97%)	8 (3%)	39	45
2	B	344/371 (93%)	339 (98%)	5 (2%)	57	65
3	C	4/11 (36%)	3 (75%)	1 (25%)	0	0
All	All	649/720 (90%)	635 (98%)	14 (2%)	45	53

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	LEU
1	A	156	ILE
1	A	173	ARG
1	A	257	GLU
1	A	303	GLN
1	A	330	LYS
1	A	347	GLU
1	A	364	GLN
2	B	29	GLU
2	B	68	VAL
2	B	87	ARG
2	B	121	ILE
2	B	351	LEU
3	C	9	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	GLN
1	A	108	GLN
1	A	364	GLN
2	B	56	GLN
2	B	80	HIS
2	B	410	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	FAR	C	12	3	14,14,14	0.38	0	15,16,16	0.71	0
5	ACY	B	3003	-	3,3,3	1.16	0	3,3,3	1.57	1 (33%)
5	ACY	B	3002	-	3,3,3	1.31	0	3,3,3	1.58	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FAR	C	12	3	-	0/14/14/14	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	3002	ACY	O-C-CH3	-2.16	113.65	122.53
5	B	3003	ACY	O-C-CH3	-2.15	113.72	122.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	12	FAR	1	0
5	B	3003	ACY	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/377 (85%)	0.02	6 (1%) 66 69	17, 33, 65, 76	0
2	B	407/437 (93%)	-0.21	10 (2%) 58 61	18, 29, 53, 78	0
3	C	5/11 (45%)	0.21	0 100 100	20, 23, 46, 50	0
All	All	735/825 (89%)	-0.11	16 (2%) 62 65	17, 31, 59, 78	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	423	PHE	6.1
1	A	55	PHE	3.8
2	B	19	TRP	3.6
2	B	64	PHE	3.4
2	B	18	VAL	3.0
1	A	373	ILE	2.9
2	B	17	PRO	2.8
1	A	375	ALA	2.7
2	B	380	ALA	2.5
1	A	304	PRO	2.4
2	B	325	MET	2.4
1	A	364	GLN	2.3
1	A	374	PRO	2.1
2	B	70	ARG	2.1
2	B	20	SER	2.1
2	B	266	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ACY	B	3003	4/4	0.85	0.22	53,55,56,59	0
6	FAR	C	12	15/15	0.95	0.10	12,20,39,43	0
5	ACY	B	3002	4/4	0.98	0.07	24,25,26,27	0
4	ZN	B	1001	1/1	1.00	0.02	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.