



Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 09:10 PM UTC

PDB ID : 1KZX / pdb_00001kzx
Title : Solution structure of human intestinal fatty acid binding protein with a naturally-occurring single amino acid substitution (A54T)
Authors : Zhang, F.; Luecke, C.; Baier, L.J.; Sacchettini, J.C.; Hamilton, J.A.
Deposited on : 2002-02-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	131	

2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:72, A:76-A:131 (127)	0.87	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	3, 6, 7, 8, 12, 14, 15, 17, 19, 20
2	1, 2, 10, 18
3	13, 16
4	4, 5
Single-model clusters	9; 11

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2111 atoms, of which 1046 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called INTESTINAL FATTY ACID-BINDING PROTEIN (T54).

Mol	Chain	Residues	Atoms						Trace
1	A	131	Total	C	H	N	O	S	0
			2111	674	1046	183	206	2	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	54	THR	ALA	REMARK 999	UNP P12104

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2 Scores per residue for each member of the ensemble

Colouring as in section [4.1](#) above.


4.2.1 Score per residue for model 1

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.3 Score per residue for model 3

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.4 Score per residue for model 4

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.5 Score per residue for model 5

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.6 Score per residue for model 6

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.7 Score per residue for model 7

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.8 Score per residue for model 8

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.9 Score per residue for model 9

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.10 Score per residue for model 10

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.11 Score per residue for model 11

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.12 Score per residue for model 12

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.13 Score per residue for model 13

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.14 Score per residue for model 14

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.15 Score per residue for model 15

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.16 Score per residue for model 16

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.17 Score per residue for model 17

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.18 Score per residue for model 18

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)




4.2.19 Score per residue for model 19

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)



4.2.20 Score per residue for model 20

- Molecule 1: INTESTINAL FATTY ACID-BINDING PROTEIN (T54)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing combined with torsion angle dynamics and followed by energy minimization.*

Of the 300 calculated structures, 20 were deposited, based on the following criterion: *target function.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
Discover	refinement	97

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.84±0.01	6±1/1059 (0.6± 0.1%)	1.65±0.03	15±3/1421 (1.0± 0.2%)
All	All	1.84	125/21180 (0.6%)	1.65	292/28420 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.9±1.2
All	All	0	39

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	72	LEU	CA-C	6.74	1.57	1.53	14	1
1	A	131	ASP	CG-OD1	6.44	1.37	1.25	17	15
1	A	59	GLU	CD-OE2	6.37	1.37	1.25	4	9
1	A	111	ASP	CG-OD1	6.29	1.37	1.25	12	15
1	A	59	GLU	CD-OE1	6.27	1.37	1.25	15	11
1	A	15	ASP	CG-OD1	6.25	1.37	1.25	18	12
1	A	15	ASP	CG-OD2	6.17	1.37	1.25	14	8
1	A	111	ASP	CG-OD2	6.14	1.37	1.25	5	5
1	A	131	ASP	CG-OD2	5.98	1.36	1.25	2	5
1	A	72	LEU	N-CA	5.82	1.49	1.46	14	1
1	A	86	GLY	N-CA	5.81	1.49	1.45	10	1
1	A	109	ILE	CA-C	5.44	1.57	1.52	1	3
1	A	33	HIS	ND1-CE1	5.39	1.38	1.32	9	19
1	A	33	HIS	CE1-NE2	5.25	1.37	1.32	7	19
1	A	26	VAL	CA-C	5.21	1.59	1.52	20	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	33	HIS	ND1-CE1-NE2	13.01	121.41	108.40	9	20
1	A	100	ASN	CA-CB-CG	11.10	123.70	112.60	14	7
1	A	33	HIS	CE1-NE2-CD2	-10.86	98.14	109.00	9	20
1	A	97	ASP	CA-CB-CG	10.60	123.20	112.60	14	19
1	A	13	ASN	CA-CB-CG	9.54	122.14	112.60	16	6
1	A	71	ASN	CA-CB-CG	9.45	122.05	112.60	15	8
1	A	34	ASP	CA-CB-CG	9.32	121.92	112.60	9	16
1	A	98	ASN	CA-CB-CG	9.31	121.91	112.60	2	13
1	A	2	PHE	CA-CB-CG	8.46	122.26	113.80	10	5
1	A	25	ILE	N-CA-C	8.06	119.81	110.62	5	11
1	A	87	ASN	CA-CB-CG	8.06	120.66	112.60	15	2
1	A	3	ASP	CA-CB-CG	7.58	120.18	112.60	2	7
1	A	111	ASP	CA-CB-CG	7.54	120.14	112.60	15	14
1	A	128	PHE	CA-CB-CG	7.53	121.33	113.80	20	12
1	A	55	PHE	CA-CB-CG	7.25	121.05	113.80	8	4
1	A	62	PHE	CA-CB-CG	7.06	120.86	113.80	14	11
1	A	57	ASN	CA-CB-CG	7.02	119.62	112.60	11	7
1	A	33	HIS	CA-CB-CG	6.92	120.72	113.80	5	4
1	A	47	PHE	CA-CB-CG	6.52	120.32	113.80	17	1
1	A	31	ALA	N-CA-C	6.36	119.02	111.71	16	5
1	A	33	HIS	CG-ND1-CE1	-6.36	98.50	109.30	9	20
1	A	12	GLU	CB-CA-C	6.11	122.58	110.42	12	6
1	A	98	ASN	N-CA-CB	6.10	120.80	110.49	11	2
1	A	77	GLU	CB-CG-CD	6.01	122.82	112.60	13	2
1	A	15	ASP	N-CA-C	5.95	118.25	111.11	10	1
1	A	24	ASN	CA-CB-CG	5.90	118.50	112.60	12	2
1	A	115	GLN	OE1-CD-NE2	-5.90	116.70	122.60	3	1
1	A	13	ASN	N-CA-CB	-5.88	102.37	110.42	12	5
1	A	15	ASP	CA-CB-CG	5.85	118.45	112.60	15	1
1	A	21	MET	N-CA-C	5.84	119.50	112.38	8	5
1	A	17	PHE	CA-CB-CG	5.82	119.62	113.80	15	5
1	A	97	ASP	N-CA-CB	-5.81	101.65	110.07	5	2
1	A	80	GLY	N-CA-C	5.80	119.23	110.18	7	5
1	A	131	ASP	N-CA-CB	5.74	120.26	110.50	18	1
1	A	97	ASP	N-CA-C	5.71	117.98	111.02	16	1
1	A	131	ASP	CA-CB-CG	5.65	118.25	112.60	6	1
1	A	13	ASN	N-CA-C	5.65	119.48	112.47	11	1
1	A	26	VAL	N-CA-C	5.64	115.84	110.42	7	3
1	A	100	ASN	N-CA-C	5.53	119.75	112.89	10	2
1	A	103	ASN	CA-CB-CG	5.53	118.12	112.60	13	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	18	MET	N-CA-C	5.52	116.98	111.07	2	3
1	A	13	ASN	CB-CA-C	5.52	118.52	111.40	12	1
1	A	55	PHE	N-CA-C	5.46	116.91	111.07	18	2
1	A	69	ASN	CA-CB-CG	5.45	118.05	112.60	2	3
1	A	9	ASP	CA-CB-CG	5.43	118.03	112.60	7	2
1	A	32	ALA	N-CA-C	5.39	117.15	111.28	16	2
1	A	69	ASN	OD1-CG-ND2	-5.37	117.23	122.60	20	1
1	A	129	LYS	N-CA-CB	-5.30	101.99	110.63	6	1
1	A	120	GLU	CB-CG-CD	5.29	121.60	112.60	4	2
1	A	96	THR	CA-CB-CG2	5.28	119.47	110.50	9	1
1	A	45	ASN	CA-CB-CG	5.24	117.84	112.60	16	1
1	A	100	ASN	OD1-CG-ND2	-5.17	117.43	122.60	14	1
1	A	109	ILE	CA-CB-CG1	5.12	119.11	110.40	12	1
1	A	71	ASN	N-CA-CB	5.12	119.14	110.49	8	1
1	A	28	ARG	N-CA-C	5.10	117.57	111.71	9	3
1	A	51	GLU	CB-CG-CD	5.09	121.26	112.60	8	1
1	A	14	TYR	N-CA-C	5.07	116.89	111.36	14	1
1	A	46	LYS	N-CA-CB	-5.07	103.04	110.85	6	1
1	A	109	ILE	CB-CA-C	5.07	118.46	112.68	20	1
1	A	33	HIS	CB-CG-CD2	-5.04	124.64	131.20	8	2
1	A	30	LEU	N-CA-C	5.03	116.76	111.28	15	1
1	A	14	TYR	CA-CB-CG	5.00	122.90	113.90	16	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	119	TYR	Sidechain	10
1	A	56	ARG	Sidechain	6
1	A	28	ARG	Sidechain	5
1	A	126	ARG	Sidechain	3
1	A	14	TYR	Sidechain	2
1	A	95	ARG	Sidechain	2
1	A	10	ARG	Sidechain	2
1	A	117	TYR	Sidechain	2
1	A	2	PHE	Peptide	1
1	A	88	LYS	Peptide	1
1	A	76	THR	Peptide	1
1	A	11	SER	Peptide	1
1	A	55	PHE	Sidechain	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	71	ASN	Peptide	1
1	A	98	ASN	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1043	1029	1044	1±1
All	All	20860	20580	20880	21

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:18:MET:SD	1:A:23:VAL:HG11	0.58	2.38	9	2
1:A:18:MET:SD	1:A:23:VAL:HG21	0.50	2.45	17	1
1:A:21:MET:SD	1:A:76:THR:HG21	0.44	2.52	20	1
1:A:119:TYR:OH	1:A:120:GLU:OE2	0.44	2.36	9	2
1:A:131:ASP:OD1	1:A:131:ASP:O	0.44	2.36	20	5
1:A:14:TYR:OH	1:A:34:ASP:OD2	0.43	2.36	8	1
1:A:76:THR:OG1	1:A:97:ASP:OD1	0.43	2.33	3	1
1:A:3:ASP:OD2	1:A:4:SER:OG	0.43	2.37	3	1
1:A:111:ASP:OD2	1:A:112:GLU:OE2	0.42	2.37	7	1
1:A:51:GLU:OE1	1:A:70:TYR:OH	0.42	2.37	8	1
1:A:14:TYR:CE2	1:A:18:MET:SD	0.41	3.14	20	1
1:A:43:GLU:OE1	1:A:48:THR:OG1	0.41	2.38	10	2
1:A:41:THR:OG1	1:A:43:GLU:OE1	0.40	2.37	9	1
1:A:111:ASP:OD1	1:A:112:GLU:OE2	0.40	2.39	10	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/131 (96%)	111±3 (88±2%)	13±3 (10±2%)	2±1 (1±1%)	11	57
All	All	2520/2620 (96%)	2221 (88%)	262 (10%)	37 (1%)	11	57

All 17 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	44	GLY	5
1	A	22	GLY	4
1	A	86	GLY	4
1	A	26	VAL	3
1	A	110	GLY	3
1	A	25	ILE	2
1	A	101	GLU	2
1	A	71	ASN	2
1	A	55	PHE	2
1	A	98	ASN	2
1	A	12	GLU	2
1	A	72	LEU	1
1	A	31	ALA	1
1	A	15	ASP	1
1	A	11	SER	1
1	A	4	SER	1
1	A	14	TYR	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/116 (99%)	110±2 (96±2%)	5±2 (4±2%)	28	78
All	All	2300/2320 (99%)	2203 (96%)	97 (4%)	28	78

All 50 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	10	ARG	6
1	A	98	ASN	5
1	A	12	GLU	5
1	A	97	ASP	4
1	A	120	GLU	4
1	A	95	ARG	4
1	A	7	LYS	3
1	A	59	GLU	3
1	A	122	VAL	3
1	A	71	ASN	3
1	A	16	LYS	3
1	A	77	GLU	3
1	A	107	GLU	3
1	A	84	LEU	2
1	A	101	GLU	2
1	A	20	LYS	2
1	A	19	GLU	2
1	A	63	GLU	2
1	A	15	ASP	2
1	A	43	GLU	2
1	A	90	ILE	2
1	A	45	ASN	2
1	A	109	ILE	2
1	A	42	GLN	2
1	A	131	ASP	1
1	A	18	MET	1
1	A	94	LYS	1
1	A	34	ASP	1
1	A	53	SER	1
1	A	56	ARG	1
1	A	112	GLU	1
1	A	116	THR	1
1	A	79	ARG	1
1	A	88	LYS	1
1	A	85	GLU	1
1	A	108	ILE	1
1	A	96	THR	1
1	A	28	ARG	1
1	A	46	LYS	1
1	A	76	THR	1
1	A	100	ASN	1
1	A	111	ASP	1
1	A	105	VAL	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	130	LYS	1
1	A	13	ASN	1
1	A	21	MET	1
1	A	115	GLN	1
1	A	27	LYS	1
1	A	104	THR	1
1	A	125	LYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided