



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:46 AM UTC

PDB ID : 1L0V / pdb_0000110v
Title : Quinol-Fumarate Reductase with Menaquinol Molecules
Authors : Iverson, T.M.; Luna-Chavez, C.; Croal, L.R.; Cecchini, G.; Rees, D.C.
Deposited on : 2002-02-13
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	602	63% 29% . .
1	M	602	21% 52% 20% . .
2	B	243	64% 33% . .
2	N	243	44% 47% 7% .
3	C	130	58% 35% 5% .
3	O	130	47% 45% 5% .
4	D	119	46% 48% 5% .
4	P	119	45% 45% 8% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	MQ7	D	700	-	-	X	-
6	FAD	M	803	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 17046 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate reductase flavoprotein subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	0	0
			4448	2775	802	840	31			
1	M	577	Total	C	N	O	S	0	0	0
			4448	2775	802	840	31			

- Molecule 2 is a protein called Fumarate reductase iron-sulfur protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	243	Total	C	N	O	S	0	0	0
			1888	1189	323	357	19			
2	N	243	Total	C	N	O	S	0	0	0
			1888	1189	323	357	19			


- Molecule 3 is a protein called Fumarate reductase 15 kDa hydrophobic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	130	Total	C	N	O	S	0	0	0
			1058	720	166	169	3			
3	O	130	Total	C	N	O	S	0	0	0
			1058	720	166	169	3			

- Molecule 4 is a protein called Fumarate reductase 13 kDa hydrophobic protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	119	Total	C	N	O	S	0	0	0
			926	626	151	142	7			
4	P	119	Total	C	N	O	S	0	0	0
			926	626	151	142	7			

- Molecule 5 is OXALOACETATE ION (CCD ID: OAA) (formula: $C_4H_3O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			9	4	5		
5	M	1	Total	C	O	0	0
			9	4	5		

- Molecule 6 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0
			53	27	9	15	2	
6	M	1	Total	C	N	O	P	0
			53	27	9	15	2	

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			4	2	2		
7	N	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 8 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			7	3	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	N	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 9 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			8	4	4		
9	N	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 10 is MENAQUINONE-7 (CCD ID: MQ7) (formula: $\text{C}_{46}\text{H}_{64}\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			24	22	2		
10	D	1	Total	C	O	0	0
			24	22	2		
10	N	1	Total	C	O	0	0
			24	22	2		
10	P	1	Total	C	O	0	0
			24	22	2		

- Molecule 11 is O-DODECANYL OCTAETHYLENE GLYCOL (CCD ID: CE1) (formula: $C_{28}H_{58}O_9$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	D	1	Total 37	C 28	O 9	0	0
11	D	1	Total 37	C 28	O 9	0	0
11	O	1	Total 37	C 28	O 9	0	0
11	O	1	Total 37	C 28	O 9	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Fumarate reductase flavoprotein subunit




- Molecule 1: Fumarate reductase flavoprotein subunit




Chain C:  58% 35% 5% •





- Molecule 3: Fumarate reductase 15 kDa hydrophobic protein

Chain O:  47% 45% 5% •




- Molecule 4: Fumarate reductase 13 kDa hydrophobic protein

Chain D:  46% 48% 5% •



- Molecule 4: Fumarate reductase 13 kDa hydrophobic protein

Chain P:  45% 45% 8% •



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.59Å 138.09Å 275.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-3.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.245 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	17046	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, FES, MQ7, CE1, OAA, F3S, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	2/4540 (0.0%)	1.16	33/6139 (0.5%)
1	M	0.58	1/4540 (0.0%)	1.18	54/6139 (0.9%)
2	B	1.17	10/1931 (0.5%)	1.48	24/2617 (0.9%)
2	N	1.21	10/1931 (0.5%)	1.39	26/2617 (1.0%)
3	C	0.66	0/1094	1.07	8/1496 (0.5%)
3	O	0.66	0/1094	1.10	8/1496 (0.5%)
4	D	0.70	0/956	1.08	4/1303 (0.3%)
4	P	1.49	3/956 (0.3%)	1.18	11/1303 (0.8%)
All	All	0.89	26/17042 (0.2%)	1.22	168/23110 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	N	0	3
4	P	0	1
All	All	0	5

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	9	ASP	C-N	-37.54	0.83	1.33
2	B	243	ARG	CA-C	24.28	2.04	1.52
2	N	69	VAL	C-N	-21.89	1.03	1.33
2	N	159	PRO	C-N	-21.54	1.04	1.33
2	N	64	SER	C-N	18.66	1.59	1.33

The worst 5 of 168 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	241	LYS	CA-C-N	25.27	151.42	119.84
2	B	241	LYS	C-N-CA	25.27	151.42	119.84
2	B	243	ARG	CB-CG-CD	18.72	154.36	111.30
2	N	69	VAL	CA-C-N	18.09	148.05	122.36
2	N	69	VAL	C-N-CA	18.09	148.05	122.36

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	243	ARG	Sidechain
2	N	183	SER	Mainchain
2	N	54	ARG	Mainchain
2	N	64	SER	Mainchain
4	P	9	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4448	0	4335	155	0
1	M	4448	0	4333	694	0
2	B	1888	0	1837	84	0
2	N	1888	0	1834	176	0
3	C	1058	0	1108	54	0
3	O	1058	0	1108	88	0
4	D	926	0	971	93	0
4	P	926	0	970	105	0
5	A	9	0	2	1	0
5	M	9	0	2	1	0
6	A	53	0	31	7	0
6	M	53	0	31	23	0
7	B	4	0	0	1	0
7	N	4	0	0	0	0
8	B	7	0	0	0	0
8	N	7	0	0	1	0
9	B	8	0	0	0	0
9	N	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	B	24	0	23	3	0
10	D	24	0	23	24	0
10	N	24	0	23	13	0
10	P	24	0	23	19	0
11	D	74	0	116	8	0
11	O	74	0	116	1	0
All	All	17046	0	16886	1351	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 40.

The worst 5 of 1351 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:98:LEU:CD2	2:N:132:ASN:HD21	1.28	1.42
4:P:31:MET:HE1	10:P:800:MQ7:O1	1.23	1.34
1:M:44:HIS:NE2	6:M:803:FAD:HM81	1.41	1.34
1:M:493:ASP:OD2	2:N:50:ASP:HA	1.27	1.31
1:M:98:LEU:HD23	2:N:132:ASN:ND2	1.42	1.31

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	575/602 (96%)	529 (92%)	37 (6%)	9 (2%)	7	31
1	M	575/602 (96%)	355 (62%)	153 (27%)	67 (12%)	0	1
2	B	241/243 (99%)	215 (89%)	22 (9%)	4 (2%)	7	30
2	N	241/243 (99%)	205 (85%)	34 (14%)	2 (1%)	16	45
3	C	128/130 (98%)	121 (94%)	5 (4%)	2 (2%)	7	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	O	128/130 (98%)	113 (88%)	12 (9%)	3 (2%)	5	25
4	D	117/119 (98%)	101 (86%)	13 (11%)	3 (3%)	4	23
4	P	117/119 (98%)	99 (85%)	12 (10%)	6 (5%)	1	11
All	All	2122/2188 (97%)	1738 (82%)	288 (14%)	96 (4%)	2	13

5 of 96 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	242	PRO
4	D	99	VAL
1	M	57	GLN
1	M	76	LEU
1	M	85	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/475 (97%)	447 (97%)	13 (3%)	38	62
1	M	460/475 (97%)	398 (86%)	62 (14%)	4	16
2	B	205/205 (100%)	198 (97%)	7 (3%)	32	59
2	N	205/205 (100%)	192 (94%)	13 (6%)	16	44
3	C	111/111 (100%)	103 (93%)	8 (7%)	13	40
3	O	111/111 (100%)	101 (91%)	10 (9%)	9	31
4	D	97/97 (100%)	94 (97%)	3 (3%)	35	60
4	P	97/97 (100%)	94 (97%)	3 (3%)	35	60
All	All	1746/1776 (98%)	1627 (93%)	119 (7%)	14	42

5 of 119 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	183	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	O	37	TRP
1	M	348	ILE
3	O	28	ARG
4	P	118	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	182	GLN
4	P	4	ASN
1	M	325	HIS
3	O	72	ASN
2	N	129	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FAD	M	803	1	58,58,58	1.69	14 (24%)	85,89,89	1.63	18 (21%)
10	MQ7	D	700	-	25,25,49	3.27	5 (20%)	31,34,63	2.44	11 (35%)
10	MQ7	P	800	-	25,25,49	3.22	5 (20%)	31,34,63	2.43	12 (38%)
11	CE1	O	812	-	36,36,36	1.16	0	35,35,35	1.56	7 (20%)
5	OAA	M	802	-	8,8,8	1.24	1 (12%)	8,10,10	1.24	1 (12%)
8	F3S	N	245	2	0,9,9	-	-	-	-	-
8	F3S	B	245	2	0,9,9	-	-	-	-	-
11	CE1	D	810	-	36,36,36	1.15	0	35,35,35	1.69	14 (40%)
11	CE1	D	710	-	36,36,36	1.04	0	35,35,35	1.66	14 (40%)
10	MQ7	B	701	-	25,25,49	3.27	6 (24%)	31,34,63	2.44	9 (29%)
10	MQ7	N	801	-	25,25,49	3.43	7 (28%)	31,34,63	2.55	10 (32%)
7	FES	N	244	2	0,4,4	-	-	-	-	-
9	SF4	B	246	2	0,12,12	-	-	-	-	-
11	CE1	O	811	-	36,36,36	1.24	4 (11%)	35,35,35	1.81	13 (37%)
5	OAA	A	702	-	8,8,8	1.15	0	8,10,10	1.06	1 (12%)
9	SF4	N	246	2	0,12,12	-	-	-	-	-
6	FAD	A	703	1	58,58,58	1.99	15 (25%)	85,89,89	1.83	21 (24%)
7	FES	B	244	2	0,4,4	-	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FAD	M	803	1	-	10/34/50/50	0/6/6/6
10	MQ7	D	700	-	-	2/13/33/61	0/2/2/2
10	MQ7	P	800	-	-	1/13/33/61	0/2/2/2
11	CE1	O	812	-	-	18/34/34/34	-
5	OAA	M	802	-	-	6/8/8/8	-
8	F3S	N	245	2	-	-	0/3/3/3
8	F3S	B	245	2	-	-	0/3/3/3
11	CE1	D	810	-	-	15/34/34/34	-
11	CE1	D	710	-	-	14/34/34/34	-
10	MQ7	B	701	-	-	3/13/33/61	0/2/2/2
10	MQ7	N	801	-	-	3/13/33/61	0/2/2/2
11	CE1	O	811	-	-	15/34/34/34	-
7	FES	N	244	2	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	SF4	B	246	2	-	-	0/6/5/5
5	OAA	A	702	-	-	6/8/8/8	-
9	SF4	N	246	2	-	-	0/6/5/5
6	FAD	A	703	1	-	11/34/50/50	0/6/6/6
7	FES	B	244	2	-	-	0/1/1/1

The worst 5 of 57 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	N	801	MQ7	C11-C12	-10.94	1.33	1.50
10	N	801	MQ7	C12-C13	10.15	1.56	1.33
10	D	700	MQ7	C11-C12	-10.11	1.34	1.50
10	B	701	MQ7	C12-C13	9.96	1.56	1.33
10	P	800	MQ7	C11-C12	-9.94	1.34	1.50

The worst 5 of 131 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	N	801	MQ7	C11-C3-C4	-7.83	110.33	118.58
10	D	700	MQ7	C11-C3-C4	-7.58	110.59	118.58
10	B	701	MQ7	C11-C3-C4	-7.56	110.61	118.58
10	P	800	MQ7	C11-C3-C4	-7.20	110.99	118.58
10	P	800	MQ7	C12-C11-C3	6.74	128.69	112.08

There are no chirality outliers.

5 of 104 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	702	OAA	O3-C3-C4-O4
5	A	702	OAA	O3-C3-C4-O5
5	A	702	OAA	C2-C3-C4-O4
5	A	702	OAA	C2-C3-C4-O5
5	M	802	OAA	C1-C2-C3-O3

There are no ring outliers.

13 monomers are involved in 102 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	M	803	FAD	23	0
10	D	700	MQ7	24	0





Continued on next page...



Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	P	800	MQ7	19	0
11	O	812	CE1	1	0
5	M	802	OAA	1	0
8	N	245	F3S	1	0
11	D	810	CE1	5	0
11	D	710	CE1	3	0
10	B	701	MQ7	3	0
10	N	801	MQ7	13	0
5	A	702	OAA	1	0
6	A	703	FAD	7	0
7	B	244	FES	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











Ligand MQ7 P 800	
	
Bond lengths	Bond angles
	
Torsions	Rings



Ligand CE1 O 812	
	
Bond lengths	Bond angles
	
Torsions	Rings


Ligand CE1 D 810	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand CE1 D 710	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand MQ7 B 701	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand MQ7 N 801	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand CE1 O 811	
	
Bond lengths	Bond angles
	
Torsions	Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	N	3
4	P	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	N	4:LYS	C	5:ASN	N	1.14
1	N	159:PRO	C	160:GLN	N	1.04
1	N	69:VAL	C	70:ASN	N	1.03
1	P	9:ASP	C	10:GLU	N	0.83

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.