



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:57 AM UTC

PDB ID : 5L0K / pdb_00005l0k
Title : Crystal Structure of Autotaxin and Compound PF-8380
Authors : Durbin, J.D.
Deposited on : 2016-07-27
Resolution : 2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1819 (2.76-2.72)
Clashscore	190562	1866 (2.76-2.72)
Ramachandran outliers	187476	1830 (2.76-2.72)
Sidechain outliers	187428	1831 (2.76-2.72)
RSRZ outliers	180081	1819 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	809	6% 90% 6%
1	B	809	7% 89% 7%
2	C	2	100%
3	D	3	100%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 12625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	777	Total	C	N	O	S	0	0	0
			6174	3919	1051	1155	49			
1	B	777	Total	C	N	O	S	0	0	0
			6171	3923	1044	1155	49			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ALA	ASN	engineered mutation	UNP Q64610
A	398	ALA	ASN	engineered mutation	UNP Q64610
A	410	ALA	ASN	engineered mutation	UNP Q64610
A	591	THR	ARG	engineered mutation	UNP Q64610
A	592	GLU	LYS	engineered mutation	UNP Q64610
A	806	ALA	ASN	engineered mutation	UNP Q64610
B	53	ALA	ASN	engineered mutation	UNP Q64610
B	398	ALA	ASN	engineered mutation	UNP Q64610
B	410	ALA	ASN	engineered mutation	UNP Q64610
B	591	THR	ARG	engineered mutation	UNP Q64610
B	592	GLU	LYS	engineered mutation	UNP Q64610
B	806	ALA	ASN	engineered mutation	UNP Q64610

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			39	22	2	15			


- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		
4	B	2	Total	Zn	0	0
			2	2		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		
5	B	1	Total	Ca	0	0
			1	1		

- Molecule 6 is (3,5-dichlorophenyl)methyl 4-[3-oxo-3-(2-oxo-2,3-dihydro-1,3-benzoxazol-6-yl)propyl]piperazine-1-carboxylate (CCD ID: 6ZO) (formula: C₂₂H₂₁Cl₂N₃O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	Cl	N	O	0	0
			32	22	2	3	5		
6	B	1	Total	C	Cl	N	O	0	0
			32	22	2	3	5		

- Molecule 7 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			12	6	2	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	S	0	0
			12	6	2	3	1		

- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		
8	B	1	Total	Cl	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	65	Total	O	0	0
			65	65		
9	B	52	Total	O	0	0
			52	52		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%

MAG1
MAG2

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	95.92Å 121.42Å 156.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.23 – 2.73 95.88 – 2.73	Depositor EDS
% Data completeness (in resolution range)	98.3 (96.23-2.73) 98.3 (95.88-2.73)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.226 , 0.270 0.226 , 0.267	Depositor DCC
R_{free} test set	826 reflections (1.68%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12625	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.6846e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN, EPE, CL, BMA, 6ZO, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/6352	0.77	0/8636
1	B	0.59	0/6348	0.77	0/8633
All	All	0.58	0/12700	0.77	0/17269

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6174	0	5810	31	0
1	B	6171	0	5812	38	0
2	C	28	0	25	0	0
3	D	39	0	34	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	32	0	0	0	0
6	B	32	0	0	0	0
7	A	12	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	12	0	12	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
9	A	65	0	0	0	0
9	B	52	0	0	0	0
All	All	12625	0	11705	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:372:LEU:HD22	1:B:384:LEU:HD22	1.65	0.78
1:B:78:LEU:HD12	1:B:272:THR:HG21	1.64	0.76
1:B:546:GLU:HB3	1:B:602:ALA:HB1	1.67	0.75
1:A:635:ILE:CD1	1:A:685:LEU:HD13	2.17	0.74
1:A:635:ILE:HD13	1:A:685:LEU:HD13	1.70	0.72
1:A:711:VAL:HG13	1:A:815:VAL:HG13	1.77	0.67
1:A:259:LEU:HD22	1:A:512:MET:HE3	1.79	0.64
1:A:314:GLY:N	1:A:324:MET:HE1	2.16	0.61
1:B:259:LEU:HD22	1:B:512:MET:HE3	1.83	0.60
1:A:698:VAL:HG23	1:A:701:MET:HE2	1.84	0.60
1:A:166:ILE:HD12	1:A:351:VAL:HG11	1.84	0.59
1:B:206:PRO:HB3	1:B:389:LEU:HD22	1.87	0.56
1:B:78:LEU:HD11	1:B:274:PHE:HB2	1.88	0.56
1:A:206:PRO:HB3	1:A:389:LEU:HD22	1.88	0.55
1:A:682:PRO:HG3	1:A:685:LEU:HD12	1.89	0.55
1:B:641:VAL:HG22	1:B:685:LEU:CD2	2.39	0.53
1:A:635:ILE:HD11	1:A:685:LEU:HD13	1.91	0.53
1:A:78:LEU:HD12	1:A:272:THR:HG21	1.91	0.53
1:A:704:MET:HE1	1:A:712:TRP:CD1	2.44	0.52
1:B:394:ALA:HB2	1:B:402:TYR:CZ	2.44	0.51
1:A:493:THR:HG21	1:A:515:LEU:HD23	1.92	0.50
1:A:166:ILE:CD1	1:A:351:VAL:HG11	2.41	0.50
1:A:259:LEU:HD22	1:A:512:MET:CE	2.42	0.49
1:B:724:ALA:HA	1:B:730:VAL:HG13	1.94	0.49
1:B:394:ALA:HB2	1:B:402:TYR:CE2	2.48	0.49
1:B:711:VAL:HG13	1:B:815:VAL:HG13	1.93	0.49
1:B:535:ARG:HG2	1:B:536:THR:HG23	1.94	0.48
1:A:793:PHE:CD1	1:A:795:LEU:HD13	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:493:THR:HG21	1:B:515:LEU:HD23	1.96	0.48
1:A:526:THR:CG2	1:A:849:LEU:HD22	2.43	0.47
1:A:682:PRO:CG	1:A:685:LEU:HD12	2.45	0.47
1:B:144:VAL:HG21	1:B:186:PRO:HB2	1.95	0.47
1:B:823:THR:HG23	1:B:855:LEU:HD11	1.97	0.46
1:B:328:LEU:HA	1:B:331:ILE:HD12	1.96	0.46
1:A:483:MET:HA	1:A:483:MET:HE2	1.96	0.46
1:B:259:LEU:HD22	1:B:512:MET:CE	2.44	0.46
1:B:711:VAL:CG1	1:B:795:LEU:HD21	2.45	0.46
1:B:388:THR:HG22	1:B:476:PHE:CZ	2.51	0.46
1:B:372:LEU:CD2	1:B:384:LEU:HD22	2.43	0.45
1:B:675:MET:HE2	1:B:705:TYR:CE1	2.52	0.45
1:B:568:CYS:SG	1:B:569:ASP:N	2.90	0.45
1:B:314:GLY:CA	1:B:324:MET:HE1	2.47	0.45
1:B:675:MET:HE2	1:B:705:TYR:CZ	2.52	0.44
1:A:682:PRO:HB3	1:A:716:GLN:HB3	2.00	0.44
1:A:715:PHE:O	1:A:720:VAL:HG23	2.18	0.43
1:B:641:VAL:HG22	1:B:685:LEU:HD23	2.00	0.43
1:A:210:PHE:HB3	1:A:227:ILE:HG21	2.00	0.43
1:B:830:GLU:OE2	1:B:852:LYS:NZ	2.51	0.43
1:B:648:LEU:HD21	1:B:693:TYR:CE1	2.53	0.43
1:B:794:ILE:HD13	1:B:832:LEU:CD1	2.48	0.43
1:B:166:ILE:HD12	1:B:351:VAL:HG11	2.00	0.43
1:B:641:VAL:HG22	1:B:685:LEU:HD21	2.01	0.43
1:B:178:MET:HE2	1:B:486:VAL:HG11	1.99	0.43
1:A:555:ILE:HG23	1:A:656:VAL:CG2	2.49	0.42
1:A:431:ARG:HD3	1:A:823:THR:O	2.19	0.42
1:B:526:THR:CG2	1:B:849:LEU:HD22	2.49	0.42
1:A:309:GLN:HG3	1:A:310:PRO:HA	2.01	0.42
1:B:711:VAL:HG11	1:B:795:LEU:HD21	2.02	0.42
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.55	0.42
1:A:708:PHE:CE1	1:A:795:LEU:HD23	2.55	0.42
1:B:209:THR:HG22	1:B:213:LEU:HD12	2.02	0.42
1:B:310:PRO:HB2	1:B:324:MET:HE3	2.02	0.42
1:A:268:VAL:HG21	1:A:518:LEU:HD21	2.01	0.42
1:B:408:ILE:O	1:B:412:THR:HG23	2.20	0.42
1:A:178:MET:HE1	1:A:188:ILE:HG22	2.01	0.41
1:A:268:VAL:CG2	1:A:518:LEU:HD21	2.51	0.41
1:B:394:ALA:HB2	1:B:402:TYR:CE1	2.56	0.41
1:A:388:THR:HG22	1:A:476:PHE:CZ	2.56	0.41
1:B:682:PRO:HB3	1:B:716:GLN:HB3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	771/809 (95%)	747 (97%)	23 (3%)	1 (0%)	48	69
1	B	769/809 (95%)	740 (96%)	28 (4%)	1 (0%)	48	69
All	All	1540/1618 (95%)	1487 (97%)	51 (3%)	2 (0%)	48	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	435	ALA
1	B	435	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	676/731 (92%)	671 (99%)	5 (1%)	76	85
1	B	677/731 (93%)	673 (99%)	4 (1%)	78	87
All	All	1353/1462 (92%)	1344 (99%)	9 (1%)	76	85

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	452	HIS
1	A	612	LEU
1	A	795	LEU
1	A	798	ARG
1	B	372	LEU
1	B	538	THR
1	B	667	LEU
1	B	785	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	HIS
1	A	427	HIS
1	A	481	ASN
1	A	537	ASN
1	A	731	ASN
1	B	115	ASN
1	B	134	GLN
1	B	197	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.59	0	17,19,21	1.44	2 (11%)
2	NAG	C	2	2	14,14,15	0.52	0	17,19,21	1.52	4 (23%)
3	NAG	D	1	3,1	14,14,15	0.56	0	17,19,21	1.20	1 (5%)
3	NAG	D	2	3	14,14,15	0.53	0	17,19,21	1.41	3 (17%)
3	BMA	D	3	3	11,11,12	0.52	0	15,15,17	1.15	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
3	NAG	D	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	O5-C1-C2	-4.63	104.12	111.29
3	D	1	NAG	O5-C1-C2	-3.71	105.54	111.29
2	C	2	NAG	C1-O5-C5	3.26	116.56	112.19
3	D	3	BMA	C1-O5-C5	2.75	115.87	112.19
3	D	2	NAG	C2-N2-C7	2.69	126.50	122.90
3	D	2	NAG	O5-C1-C2	-2.60	107.26	111.29
2	C	1	NAG	C1-O5-C5	2.44	115.46	112.19
2	C	2	NAG	O5-C1-C2	-2.38	107.61	111.29
2	C	2	NAG	C2-N2-C7	2.33	126.03	122.90
3	D	2	NAG	C1-O5-C5	2.24	115.19	112.19
2	C	2	NAG	C1-C2-N2	2.22	113.94	110.43

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	2	NAG	C3-C2-N2-C7
2	C	2	NAG	C3-C2-N2-C7

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	1	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EPE	B	908	-	12,12,15	0.95	1 (8%)	15,16,20	1.39	2 (13%)
7	EPE	A	907	-	12,12,15	0.96	1 (8%)	15,16,20	1.39	2 (13%)
6	6ZO	B	907	4	35,35,35	1.31	2 (5%)	49,49,49	2.96	14 (28%)
6	6ZO	A	906	4	35,35,35	1.29	2 (5%)	49,49,49	2.93	11 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EPE	B	908	-	-	0/6/14/19	0/1/1/1
7	EPE	A	907	-	-	4/6/14/19	0/1/1/1
6	6ZO	B	907	4	-	9/18/28/28	0/4/4/4
6	6ZO	A	906	4	-	8/18/28/28	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	906	6ZO	C30-N31	5.52	1.40	1.35
6	B	907	6ZO	C30-N31	5.40	1.39	1.35
7	A	907	EPE	C10-S	2.95	1.81	1.77
7	B	908	EPE	C10-S	2.82	1.81	1.77
6	B	907	6ZO	C9-N11	2.53	1.39	1.35
6	A	906	6ZO	C9-N11	2.49	1.39	1.35

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	907	6ZO	C28-N31-C30	-12.90	102.94	109.90
6	A	906	6ZO	C28-N31-C30	-12.63	103.09	109.90
6	A	906	6ZO	O29-C30-N31	8.12	113.01	108.20
6	B	907	6ZO	O29-C30-N31	8.00	112.94	108.20
6	A	906	6ZO	C26-O29-C30	-6.32	103.20	107.00
6	B	907	6ZO	C26-O29-C30	-6.26	103.24	107.00
6	A	906	6ZO	O8-C9-N11	6.02	122.28	111.77
6	B	907	6ZO	O8-C9-N11	5.91	122.09	111.77
6	A	906	6ZO	O32-C30-N31	-4.85	123.21	129.81
6	B	907	6ZO	O32-C30-N31	-4.70	123.41	129.81
6	B	907	6ZO	C26-C28-N31	4.69	110.02	105.61
6	A	906	6ZO	C26-C28-N31	4.11	109.48	105.61
6	A	906	6ZO	O29-C26-C28	3.61	111.08	108.80
6	B	907	6ZO	O8-C9-O10	-3.57	118.56	124.76
6	A	906	6ZO	O8-C9-O10	-3.42	118.82	124.76
7	A	907	EPE	C5-N4-C3	3.03	118.97	110.40
7	A	907	EPE	O1S-S-C10	2.98	111.23	106.73
6	B	907	6ZO	O29-C26-C28	2.93	110.64	108.80
6	A	906	6ZO	O10-C9-N11	-2.92	118.90	124.30
7	B	908	EPE	C5-N4-C3	2.83	118.40	110.40
6	B	907	6ZO	O10-C9-N11	-2.68	119.35	124.30
6	B	907	6ZO	C27-C28-C26	2.53	123.28	120.75
6	B	907	6ZO	O8-C7-C1	2.44	115.31	109.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	908	EPE	O2S-S-C10	2.43	110.40	106.73
6	A	906	6ZO	O8-C7-C1	2.15	114.62	109.40
6	B	907	6ZO	O29-C26-C24	2.15	129.47	124.95
6	A	906	6ZO	C4-C6-C5	2.13	119.98	117.43
6	B	907	6ZO	C4-C6-C5	2.10	119.95	117.43
6	B	907	6ZO	C27-C28-N31	-2.07	126.64	130.83

There are no chirality outliers.

All (21) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	906	6ZO	C17-C20-C21-C23
6	A	906	6ZO	C17-C20-C21-O22
6	B	907	6ZO	N11-C9-O8-C7
6	B	907	6ZO	O10-C9-O8-C7
7	A	907	EPE	C9-C10-S-O3S
6	A	906	6ZO	N11-C9-O8-C7
6	B	907	6ZO	C17-C20-C21-C23
6	A	906	6ZO	O10-C9-O8-C7
6	A	906	6ZO	O22-C21-C23-C24
6	A	906	6ZO	O22-C21-C23-C25
6	A	906	6ZO	C20-C21-C23-C24
6	B	907	6ZO	C17-C20-C21-O22
6	A	906	6ZO	C20-C21-C23-C25
7	A	907	EPE	C9-C10-S-O1S
7	A	907	EPE	C9-C10-S-O2S
6	B	907	6ZO	C20-C17-N16-C14
6	B	907	6ZO	O22-C21-C23-C24
6	B	907	6ZO	O22-C21-C23-C25
7	A	907	EPE	C10-C9-N1-C6
6	B	907	6ZO	C20-C21-C23-C24
6	B	907	6ZO	C20-C21-C23-C25

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	777/809 (96%)	0.48	46 (5%)	28 28	23, 36, 56, 69	0
1	B	777/809 (96%)	0.60	55 (7%)	22 21	24, 40, 60, 76	0
All	All	1554/1618 (96%)	0.54	101 (6%)	25 24	23, 38, 59, 76	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	460	VAL	5.0
1	A	469	PHE	4.5
1	B	243	LEU	4.4
1	B	71	PRO	4.4
1	B	459	ASP	4.0
1	B	244	ARG	4.0
1	A	242	HIS	3.8
1	A	468	CYS	3.7
1	B	556	MET	3.6
1	B	538	THR	3.4
1	B	161	VAL	3.4
1	B	108	GLY	3.4
1	B	537	ASN	3.4
1	A	591	THR	3.3
1	B	52	THR	3.3
1	A	69	GLY	3.3
1	A	53	ALA	3.2
1	A	619	SER	3.2
1	A	417	ASP	3.2
1	B	242	HIS	3.2
1	B	469	PHE	3.1
1	B	183	LYS	3.1
1	B	70	PRO	3.0
1	A	249	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	468	CYS	3.0
1	A	66	GLN	3.0
1	A	243	LEU	3.0
1	A	538	THR	3.0
1	B	858	TYR	2.9
1	A	68	VAL	2.9
1	A	193	SER	2.8
1	A	536	THR	2.8
1	A	244	ARG	2.8
1	A	236	VAL	2.8
1	A	108	GLY	2.7
1	A	251	HIS	2.7
1	A	569	ASP	2.7
1	A	810	ASP	2.7
1	B	412	THR	2.7
1	B	458	LEU	2.7
1	B	750	ASP	2.7
1	A	56	GLY	2.7
1	A	537	ASN	2.7
1	A	401	LYS	2.6
1	A	398	ALA	2.6
1	B	320	PHE	2.6
1	A	594	ARG	2.6
1	B	397	ILE	2.6
1	B	562	PHE	2.6
1	A	161	VAL	2.5
1	A	687	SER	2.5
1	B	560	SER	2.5
1	B	134	GLN	2.5
1	B	239	ALA	2.5
1	B	66	GLN	2.5
1	B	55	SER	2.4
1	A	70	PRO	2.4
1	B	314	GLY	2.4
1	B	331	ILE	2.4
1	B	110	VAL	2.4
1	A	643	SER	2.3
1	A	397	ILE	2.3
1	B	109	GLU	2.3
1	B	179	LYS	2.3
1	B	246	ARG	2.3
1	B	318	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	81	SER	2.3
1	A	107	CYS	2.3
1	B	270	ALA	2.3
1	B	536	THR	2.3
1	A	105	ASP	2.3
1	A	237	PHE	2.3
1	B	594	ARG	2.2
1	B	249	PHE	2.2
1	B	373	SER	2.2
1	B	65	LEU	2.2
1	B	558	LEU	2.2
1	B	547	VAL	2.2
1	A	158	ALA	2.2
1	B	107	CYS	2.2
1	B	638	GLN	2.2
1	B	251	HIS	2.2
1	B	542	THR	2.2
1	B	83	SER	2.2
1	B	619	SER	2.2
1	B	625	PHE	2.2
1	A	67	GLU	2.1
1	A	134	GLN	2.1
1	B	124	LEU	2.1
1	A	693	TYR	2.1
1	A	777	PHE	2.1
1	B	673	LYS	2.1
1	A	71	PRO	2.1
1	A	547	VAL	2.1
1	A	817	GLU	2.1
1	B	250	ASN	2.1
1	A	540	ARG	2.0
1	A	55	SER	2.0
1	A	59	LYS	2.0
1	B	457	PRO	2.0
1	A	124	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	D	3	11/12	0.46	0.22	78,82,84,84	0
2	NAG	C	2	14/15	0.75	0.19	53,58,60,61	0
3	NAG	D	2	14/15	0.82	0.17	56,61,65,72	0
2	NAG	C	1	14/15	0.92	0.12	31,35,39,47	0
3	NAG	D	1	14/15	0.93	0.10	41,44,46,50	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	EPE	B	908	12/15	0.85	0.19	60,67,74,74	0
6	6ZO	A	906	32/32	0.87	0.14	51,54,58,60	0
6	6ZO	B	907	32/32	0.89	0.14	47,59,62,63	0
7	EPE	A	907	12/15	0.91	0.14	49,60,69,70	0
8	CL	B	909	1/1	0.96	0.10	50,50,50,50	0
4	ZN	B	905	1/1	0.98	0.05	63,63,63,63	0
5	CA	A	905	1/1	0.98	0.06	33,33,33,33	0
5	CA	B	906	1/1	0.98	0.04	39,39,39,39	0
8	CL	A	908	1/1	0.98	0.05	34,34,34,34	0
4	ZN	A	904	1/1	0.98	0.05	54,54,54,54	0
4	ZN	B	904	1/1	0.99	0.04	37,37,37,37	0
4	ZN	A	903	1/1	0.99	0.05	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 6ZO A 906:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 6ZO B 907:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.