



wwPDB EM Validation Summary Report ⓘ

May 3, 2026 – 12:52 AM JST

PDB ID : 9LOI / pdb_00009loi
EMDB ID : EMD-62715
Title : Cryo-EM structure of human lipid phosphate phosphatase 1
Authors : Yang, M.; Qian, H.W.
Deposited on : 2024-12-12
Resolution : 2.61 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : **NOT EXECUTED**
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	229148	23984
Ramachandran outliers	224038	23583
Sidechain outliers	223484	23102

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	284	
1	B	284	
1	C	284	
1	D	284	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phospholipid phosphatase 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	252	Total	C	N	O	S	0	0
			2008	1315	331	352	10		
1	B	252	Total	C	N	O	S	0	0
			2008	1315	331	352	10		
1	C	252	Total	C	N	O	S	0	0
			2008	1315	331	352	10		
1	D	252	Total	C	N	O	S	0	0
			2008	1315	331	352	10		

- Molecule 2 is 2-(HEXADECANOYLOXY)-1-[(PHOSPHONOXY)METHYL]ETHYL HEXADECANOATE (CCD ID: LPP) (formula: C₃₅H₆₉O₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	O	P	0
			44	35	8	1	
2	B	1	Total	C	O	P	0
			44	35	8	1	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				AltConf
2	C	1	Total	C	O	P	0
			44	35	8	1	
2	D	1	Total	C	O	P	0
			44	35	8	1	

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	B	1	Total	C	N	O	0
			14	8	1	5	
3	C	1	Total	C	N	O	0
			14	8	1	5	
3	D	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 4 is 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (CCD ID: LBN) (formula: $C_{42}H_{82}NO_8P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
4	A	1	Total	C	O	P	0
			45	36	8	1	
4	B	1	Total	C	O	P	0
			45	36	8	1	
4	C	1	Total	C	O	P	0
			45	36	8	1	
4	D	1	Total	C	O	P	0
			45	36	8	1	

- Molecule 5 is [(2R)-1-octadecanoyloxy-3-[oxidanyl-[(1R,2R,3S,4R,5R,6S)-2,3,6-tris(oxidanyl)-4,5-diphosphonoxy-cyclohexyl]oxy-phosphoryl]oxy-propan-2-yl] (8Z)-icosa-5,8,11,14-tetraenoate (CCD ID: PT5) (formula: C₄₇H₈₅O₁₉P₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
5	A	1	Total	C	O	P	0
			62	40	19	3	
5	B	1	Total	C	O	P	0
			62	40	19	3	
5	C	1	Total	C	O	P	0
			62	40	19	3	
5	D	1	Total	C	O	P	0
			62	40	19	3	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Phospholipid phosphatase 1




• Molecule 1: Phospholipid phosphatase 1



• Molecule 1: Phospholipid phosphatase 1



• Molecule 1: Phospholipid phosphatase 1



L198	R199	P200		Q203	F204		L215	S216	R217	V218		S226		L229		A239		K251	E252	R253	THR	SER	SER	PHE	LYS	LYS	GLU	ARG	LYS	LYS	GLU	GLU	ASP	SER	SER	HIS	THR	THR	LEU	HIS	HIS	GLU	THR	PRO	THR	THR	GLY	ASN	HIS	TYR	PRO	PRO	SER	ASN	ASN	HIS	GLN	PRO
------	------	------	--	------	------	--	------	------	------	------	--	------	--	------	--	------	--	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	142002	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PT5, LBN, NAG, LPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	1/2060 (0.0%)	0.75	4/2797 (0.1%)
1	B	0.21	0/2060	0.65	2/2797 (0.1%)
1	C	0.21	0/2060	0.65	2/2797 (0.1%)
1	D	0.22	0/2060	0.64	2/2797 (0.1%)
All	All	0.23	1/8240 (0.0%)	0.67	10/11188 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	142	ASN	CG-OD1	-6.12	1.11	1.23

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ASN	OD1-CG-ND2	-9.78	112.82	122.60
1	A	142	ASN	CB-CA-C	-9.43	95.26	111.30
1	A	142	ASN	CB-CG-ND2	-8.47	103.69	116.40
1	A	142	ASN	N-CA-C	-5.70	97.96	107.99
1	C	88	SER	CA-C-N	5.12	127.85	120.38

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	142	ASN	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	0	2016	43	0
1	B	2008	0	2017	37	0
1	C	2008	0	2017	37	0
1	D	2008	0	2017	40	0
2	A	44	0	67	5	0
2	B	44	0	67	5	0
2	C	44	0	67	6	0
2	D	44	0	67	5	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
3	C	14	0	13	0	0
3	D	14	0	13	0	0
4	A	45	0	0	0	0
4	B	45	0	0	0	0
4	C	45	0	0	0	0
4	D	45	0	0	0	0
5	A	62	0	65	8	0
5	B	62	0	65	8	0
5	C	62	0	65	8	0
5	D	62	0	65	9	0
All	All	8692	0	8647	147	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 147 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ALA:HB1	1:A:198:LEU:HB3	1.66	0.77
1:D:195:ALA:HB1	1:D:198:LEU:HB3	1.73	0.69
1:B:195:ALA:HB1	1:B:198:LEU:HB3	1.74	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:195:ALA:HB1	1:C:198:LEU:HB3	1.73	0.68
1:A:193:ASP:OD1	1:D:6:ARG:NH2	2.29	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/284 (88%)	243 (97%)	6 (2%)	1 (0%)	30	49
1	B	250/284 (88%)	243 (97%)	6 (2%)	1 (0%)	30	49
1	C	250/284 (88%)	243 (97%)	6 (2%)	1 (0%)	30	49
1	D	250/284 (88%)	243 (97%)	6 (2%)	1 (0%)	30	49
All	All	1000/1136 (88%)	972 (97%)	24 (2%)	4 (0%)	31	49

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	B	87	ASN
1	C	87	ASN
1	D	87	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/248 (88%)	214 (99%)	3 (1%)	59	80
1	B	217/248 (88%)	215 (99%)	2 (1%)	70	86
1	C	217/248 (88%)	215 (99%)	2 (1%)	70	86
1	D	217/248 (88%)	215 (99%)	2 (1%)	70	86
All	All	868/992 (88%)	859 (99%)	9 (1%)	65	85

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	31	HIS
1	D	252	GLU
1	B	31	HIS
1	B	252	GLU
1	C	31	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	C	87	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	302	1	14,14,15	0.58	0	17,19,21	0.88	1 (5%)
2	LPP	D	301	-	43,43,43	0.90	4 (9%)	47,48,48	1.08	2 (4%)
5	PT5	C	304	-	62,62,69	0.90	2 (3%)	76,80,87	0.97	3 (3%)
3	NAG	B	302	1	14,14,15	0.59	0	17,19,21	0.89	1 (5%)
5	PT5	A	304	-	62,62,69	0.90	2 (3%)	76,80,87	0.96	3 (3%)
5	PT5	B	304	-	62,62,69	0.90	2 (3%)	76,80,87	0.97	3 (3%)
4	LBN	B	303	-	44,44,51	1.27	7 (15%)	48,49,59	1.16	3 (6%)
2	LPP	C	301	-	43,43,43	0.91	4 (9%)	47,48,48	1.09	2 (4%)
4	LBN	D	303	-	44,44,51	1.27	7 (15%)	48,49,59	1.16	3 (6%)
4	LBN	C	303	-	44,44,51	1.27	7 (15%)	48,49,59	1.16	3 (6%)
4	LBN	A	303	-	44,44,51	1.27	7 (15%)	48,49,59	1.16	3 (6%)
2	LPP	A	301	-	43,43,43	0.90	4 (9%)	47,48,48	1.09	2 (4%)
3	NAG	A	302	1	14,14,15	0.63	1 (7%)	17,19,21	0.61	0
3	NAG	D	302	1	14,14,15	0.48	0	17,19,21	0.77	1 (5%)
2	LPP	B	301	-	43,43,43	0.91	4 (9%)	47,48,48	1.09	2 (4%)
5	PT5	D	304	-	62,62,69	0.90	2 (3%)	76,80,87	0.97	3 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	302	1	-	1/6/23/26	0/1/1/1
2	LPP	D	301	-	-	15/45/45/45	-
5	PT5	C	304	-	-	21/59/83/90	0/1/1/1
3	NAG	B	302	1	-	1/6/23/26	0/1/1/1
5	PT5	A	304	-	-	21/59/83/90	0/1/1/1
5	PT5	B	304	-	-	21/59/83/90	0/1/1/1
4	LBN	B	303	-	-	22/46/46/55	-
2	LPP	C	301	-	-	15/45/45/45	-
4	LBN	D	303	-	-	23/46/46/55	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LBN	C	303	-	-	23/46/46/55	-
4	LBN	A	303	-	-	22/46/46/55	-
2	LPP	A	301	-	-	16/45/45/45	-
3	NAG	A	302	1	-	1/6/23/26	0/1/1/1
3	NAG	D	302	1	-	1/6/23/26	0/1/1/1
2	LPP	B	301	-	-	15/45/45/45	-
5	PT5	D	304	-	-	21/59/83/90	0/1/1/1

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	304	PT5	O18-C11	4.44	1.46	1.33
5	A	304	PT5	O18-C11	4.42	1.46	1.33
5	B	304	PT5	O18-C11	4.41	1.46	1.33
5	D	304	PT5	O18-C11	4.41	1.46	1.33
5	B	304	PT5	O16-C10	4.19	1.46	1.34

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	304	PT5	O16-C10-C12	4.37	120.92	111.50
5	D	304	PT5	O16-C10-C12	4.36	120.89	111.50
5	C	304	PT5	O16-C10-C12	4.33	120.84	111.50
5	A	304	PT5	O16-C10-C12	4.31	120.80	111.50
4	B	303	LBN	O7-C34-C35	3.91	119.93	111.50

There are no chirality outliers.

5 of 239 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	LPP	C6-O5-P1-O2
2	A	301	LPP	C6-O5-P1-O3
2	A	301	LPP	C6-O5-P1-O4
2	B	301	LPP	C6-O5-P1-O2
2	B	301	LPP	C6-O5-P1-O3



There are no ring outliers.



8 monomers are involved in 54 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	LPP	5	0
5	C	304	PT5	8	0
5	A	304	PT5	8	0
5	B	304	PT5	8	0
2	C	301	LPP	6	0
2	A	301	LPP	5	0
2	B	301	LPP	5	0
5	D	304	PT5	9	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.