



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:53 AM UTC

PDB ID : 7L1G / pdb_0000711g
Title : PRMT5-MEP50 Complexed with SAM
Authors : Palte, R.L.
Deposited on : 2020-12-14
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	 2% 84% 13% . .
2	B	350	 8% 74% 14% • 11%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein arginine N-methyltransferase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	625	5102	3264	877	935	26	0	4	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP O14744
A	-6	ASP	-	expression tag	UNP O14744
A	-5	TYR	-	expression tag	UNP O14744
A	-4	LYS	-	expression tag	UNP O14744
A	-3	ASP	-	expression tag	UNP O14744
A	-2	ASP	-	expression tag	UNP O14744
A	-1	ASP	-	expression tag	UNP O14744
A	0	ASP	-	expression tag	UNP O14744
A	1	LYS	-	expression tag	UNP O14744

- Molecule 2 is a protein called Methylosome protein 50.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	311	2353	1475	403	460	15	0	1	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	MET	-	initiating methionine	UNP Q9BQA1
B	-6	HIS	-	expression tag	UNP Q9BQA1
B	-5	HIS	-	expression tag	UNP Q9BQA1
B	-4	HIS	-	expression tag	UNP Q9BQA1
B	-3	HIS	-	expression tag	UNP Q9BQA1
B	-2	HIS	-	expression tag	UNP Q9BQA1
B	-1	HIS	-	expression tag	UNP Q9BQA1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	HIS	-	expression tag	UNP Q9BQA1
B	1	HIS	-	expression tag	UNP Q9BQA1

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		

- Molecule 5 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	107	Total 107	O 107	0	0
6	B	26	Total 26	O 26	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein arginine N-methyltransferase 5



- Molecule 2: Methylosome protein 50



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.55Å 138.78Å 178.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.07 – 2.47 46.07 – 2.47	Depositor EDS
% Data completeness (in resolution range)	96.6 (46.07-2.47) 97.0 (46.07-2.47)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.45Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.214 , 0.255 0.226 , 0.272	Depositor DCC
R_{free} test set	2274 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	62.7	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7677	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	0/5246	0.97	0/7135
2	B	0.91	2/2410 (0.1%)	1.01	1/3294 (0.0%)
All	All	0.88	2/7656 (0.0%)	0.98	1/10429 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	147	ILE	CA-C	5.93	1.60	1.52
2	B	27	MET	SD-CE	-5.15	1.66	1.79

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	327	VAL	N-CA-C	5.19	113.00	107.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5102	0	4989	39	0
2	B	2353	0	2266	24	0
3	A	44	0	66	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	16	0	24	0	0
4	A	2	0	0	0	0
5	A	27	0	22	2	0
6	A	107	0	0	0	0
6	B	26	0	0	0	0
All	All	7677	0	7367	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404[B]:TRP:HA	1:A:404[B]:TRP:CE3	2.26	0.70
2:B:28:GLU:HG2	2:B:48:SER:HB2	1.75	0.69
2:B:36:TYR:H	2:B:304:THR:HG21	1.58	0.69
1:A:404[B]:TRP:HA	1:A:404[B]:TRP:HE3	1.59	0.66
1:A:94:SER:O	1:A:100:ARG:HD3	2.00	0.62
2:B:37:ARG:O	2:B:307:PRO:HG3	2.01	0.60
1:A:324:TYR:HB2	1:A:368:ARG:HD2	1.83	0.59
2:B:102:ALA:HB2	2:B:122:TYR:CD1	2.38	0.58
1:A:432:ILE:HG12	1:A:457:LEU:HD13	1.85	0.58
1:A:567:ILE:HG21	1:A:579:TRP:HB2	1.86	0.56
1:A:441:ALA:HB2	1:A:555:PHE:HB2	1.87	0.56
1:A:362:MET:HG2	1:A:389:TYR:HB2	1.88	0.55
1:A:524:PRO:HG2	1:A:526:ARG:HH21	1.73	0.53
2:B:35:ARG:HB2	2:B:43:LEU:HB3	1.92	0.52
2:B:21:PRO:O	2:B:22:ASN:HB2	2.09	0.52
2:B:239:LEU:HB3	2:B:249:LEU:HB3	1.91	0.52
1:A:567:ILE:HA	1:A:572:HIS:CD2	2.44	0.51
1:A:23:VAL:HG22	1:A:29:THR:HG21	1.93	0.50
2:B:35:ARG:HA	2:B:304:THR:HG21	1.94	0.49
1:A:405:GLN:HA	1:A:409:TRP:HB2	1.93	0.49
1:A:26:ILE:HD12	1:A:43:MET:HE1	1.95	0.49
1:A:47:HIS:HB3	1:A:50:PHE:HB2	1.94	0.49
1:A:313:GLN:HE22	1:A:636:GLY:H	1.59	0.49
2:B:284:ASP:HB3	2:B:290:LEU:HD11	1.95	0.48
2:B:274:LEU:HD21	2:B:302:ASP:HA	1.95	0.48
1:A:371:LEU:HD13	1:A:433:VAL:HG12	1.95	0.48
1:A:362:MET:SD	1:A:429:ALA:HB2	2.54	0.47
1:A:324:TYR:CB	1:A:368:ARG:HD2	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:HIS:O	1:A:254:ILE:HG12	2.13	0.47
1:A:327:PHE:CE2	5:A:714:SAM:HA	2.49	0.47
1:A:485[A]:ARG:HG3	1:A:498:PHE:HZ	1.79	0.47
1:A:404[B]:TRP:HE1	3:A:708:EDO:H12	1.80	0.46
1:A:92:PRO:HG2	1:A:134:LEU:HD12	1.97	0.46
1:A:493:ASP:HB3	1:A:496:ALA:HB2	1.96	0.46
2:B:303:ALA:HB1	2:B:313:LEU:HD11	1.98	0.46
2:B:219:ALA:HB3	2:B:229:VAL:CG2	2.45	0.46
2:B:36:TYR:HB3	2:B:307:PRO:HD3	1.98	0.45
2:B:188:GLU:HG2	2:B:214:LEU:HD13	1.97	0.45
1:A:45:VAL:HG23	1:A:46:PHE:HD1	1.82	0.44
2:B:134:LEU:HD12	2:B:139:GLN:HB3	2.00	0.44
1:A:319:LEU:HD13	1:A:327:PHE:HE2	1.81	0.44
2:B:87:THR:HG22	2:B:95:LEU:HB3	1.98	0.44
1:A:592:VAL:HG21	1:A:598:ILE:HD11	2.00	0.44
2:B:44:LEU:HB2	2:B:59:TRP:HB2	2.00	0.44
1:A:348:ARG:HE	1:A:359:GLN:HE22	1.66	0.44
1:A:47:HIS:CE1	2:B:49:LEU:HD22	2.53	0.43
1:A:106:ALA:O	1:A:110:GLU:HG3	2.18	0.43
1:A:465:PRO:HB3	1:A:559:LEU:HD23	1.99	0.43
2:B:166:HIS:ND1	2:B:187:SER:HB3	2.33	0.43
1:A:301:ALA:HB1	1:A:505:ARG:HG3	2.01	0.42
1:A:319:LEU:HD22	1:A:323:THR:HG21	2.01	0.42
1:A:373:ASN:HA	1:A:376:LEU:HD12	2.01	0.42
2:B:109:ASP:HB2	2:B:114:LEU:H	1.85	0.42
1:A:260:LEU:HD23	1:A:260:LEU:HA	1.87	0.42
2:B:24:PRO:HD2	2:B:59:TRP:CH2	2.55	0.42
1:A:232:PRO:O	1:A:235:ILE:HG12	2.19	0.41
1:A:327:PHE:CZ	5:A:714:SAM:HA	2.55	0.41
2:B:184:LEU:HB3	2:B:218:LEU:HD13	2.02	0.41
2:B:274:LEU:HD12	2:B:274:LEU:N	2.35	0.41
1:A:82:ILE:O	1:A:120:PRO:HD2	2.21	0.41
1:A:92:PRO:O	1:A:100:ARG:HG3	2.21	0.41
2:B:307:PRO:HG2	2:B:308:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	627/645 (97%)	603 (96%)	23 (4%)	1 (0%)	43	61
2	B	310/350 (89%)	287 (93%)	20 (6%)	3 (1%)	12	22
All	All	937/995 (94%)	890 (95%)	43 (5%)	4 (0%)	30	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	147	ILE
2	B	22	ASN
2	B	247	CYS
1	A	441	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/570 (99%)	531 (94%)	31 (6%)	19	37
2	B	265/298 (89%)	254 (96%)	11 (4%)	26	49
All	All	827/868 (95%)	785 (95%)	42 (5%)	21	40

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	LYS
1	A	62	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	76	ARG
1	A	146	HIS
1	A	149	SER
1	A	155	VAL
1	A	169	GLU
1	A	177	GLU
1	A	220	ARG
1	A	234	SER
1	A	238	THR
1	A	240	LYS
1	A	241	LYS
1	A	276	GLU
1	A	281	LEU
1	A	289	GLN
1	A	351	GLU
1	A	361	LEU
1	A	368	ARG
1	A	385	ARG
1	A	405	GLN
1	A	445	LEU
1	A	475	ILE
1	A	483	GLU
1	A	490	LYS
1	A	504	VAL
1	A	505	ARG
1	A	526	ARG
1	A	578	SER
1	A	610	LYS
1	A	635	ILE
2	B	74	SER
2	B	107	GLU
2	B	131	VAL
2	B	141	VAL
2	B	206	ILE
2	B	208	CYS
2	B	225	SER
2	B	251	SER
2	B	283	LEU
2	B	294	GLN
2	B	304	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	263	GLN
1	A	271	HIS
1	A	290	ASN
1	A	313	GLN
1	A	359	GLN
1	A	405	GLN
1	A	532	ASN
1	A	572	HIS
2	B	22	ASN
2	B	69	ASN
2	B	111	ASN
2	B	169	GLN
2	B	178	HIS
2	B	310	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	705	-	3,3,3	0.58	0	2,2,2	0.29	0
3	EDO	A	710	-	3,3,3	0.52	0	2,2,2	0.38	0
3	EDO	A	711	-	3,3,3	0.69	0	2,2,2	0.17	0
3	EDO	B	402	-	3,3,3	0.43	0	2,2,2	0.44	0
3	EDO	A	706	-	3,3,3	0.53	0	2,2,2	0.41	0
3	EDO	B	403	-	3,3,3	0.49	0	2,2,2	0.40	0
3	EDO	A	703	-	3,3,3	0.49	0	2,2,2	0.43	0
3	EDO	B	404	-	3,3,3	0.59	0	2,2,2	0.36	0
3	EDO	A	702	-	3,3,3	0.43	0	2,2,2	0.46	0
3	EDO	A	709	-	3,3,3	0.63	0	2,2,2	0.33	0
5	SAM	A	714	-	27,29,29	0.50	0	34,42,42	0.48	0
3	EDO	A	707	-	3,3,3	0.64	0	2,2,2	0.19	0
3	EDO	A	708	-	3,3,3	0.51	0	2,2,2	0.28	0
3	EDO	A	704	-	3,3,3	0.46	0	2,2,2	0.50	0
3	EDO	A	701	-	3,3,3	0.53	0	2,2,2	0.46	0
3	EDO	B	401	-	3,3,3	0.61	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	705	-	-	0/1/1/1	-
3	EDO	A	710	-	-	1/1/1/1	-
3	EDO	A	711	-	-	0/1/1/1	-
3	EDO	B	402	-	-	0/1/1/1	-
3	EDO	A	706	-	-	1/1/1/1	-
3	EDO	B	403	-	-	0/1/1/1	-
3	EDO	A	703	-	-	0/1/1/1	-
3	EDO	B	404	-	-	0/1/1/1	-
3	EDO	A	702	-	-	1/1/1/1	-
3	EDO	A	709	-	-	1/1/1/1	-
5	SAM	A	714	-	-	11/17/33/33	0/3/3/3
3	EDO	A	707	-	-	0/1/1/1	-
3	EDO	A	708	-	-	0/1/1/1	-
3	EDO	A	704	-	-	0/1/1/1	-
3	EDO	A	701	-	-	0/1/1/1	-
3	EDO	B	401	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	714	SAM	N-CA-CB-CG
5	A	714	SAM	CA-CB-CG-SD
5	A	714	SAM	C4'-C5'-SD-CE
5	A	714	SAM	O4'-C4'-C5'-SD
5	A	714	SAM	C3'-C4'-C5'-SD
5	A	714	SAM	C-CA-CB-CG
5	A	714	SAM	CB-CG-SD-CE
3	A	706	EDO	O1-C1-C2-O2
3	A	702	EDO	O1-C1-C2-O2
3	A	710	EDO	O1-C1-C2-O2
5	A	714	SAM	O-C-CA-CB
5	A	714	SAM	OXT-C-CA-CB
3	A	709	EDO	O1-C1-C2-O2
5	A	714	SAM	OXT-C-CA-N
5	A	714	SAM	CB-CG-SD-C5'

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	714	SAM	2	0
3	A	708	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	625/645 (96%)	0.17	12 (1%) 66 63	28, 68, 107, 145	4 (0%)
2	B	311/350 (88%)	0.84	27 (8%) 16 14	40, 85, 123, 155	1 (0%)
All	All	936/995 (94%)	0.39	39 (4%) 40 36	28, 75, 114, 155	5 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	247	CYS	4.8
1	A	404[A]	TRP	4.3
2	B	248	VAL	3.4
2	B	208	CYS	3.4
2	B	23	ALA	3.4
2	B	114	LEU	3.1
2	B	19	LEU	3.1
2	B	31	LEU	3.0
2	B	249	LEU	3.0
2	B	211	PRO	2.9
1	A	327	PHE	2.8
1	A	294	PRO	2.8
1	A	577	PHE	2.8
2	B	210	ALA	2.6
2	B	290	LEU	2.6
1	A	297	TYR	2.6
2	B	327	VAL	2.5
1	A	277	PHE	2.5
1	A	139	THR	2.5
1	A	59	ALA	2.4
2	B	21	PRO	2.4
2	B	41	ALA	2.4
1	A	281	LEU	2.3
2	B	108	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	268	VAL	2.3
2	B	304	THR	2.3
2	B	316	VAL	2.3
2	B	42	LEU	2.2
2	B	115	ILE	2.2
1	A	530	ILE	2.2
2	B	244	SER	2.2
2	B	59	TRP	2.2
2	B	230	PHE	2.2
1	A	346	LEU	2.1
2	B	312	LEU	2.1
1	A	293	PRO	2.1
2	B	61	PHE	2.1
2	B	328	PRO	2.1
2	B	206	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	B	404	4/4	0.58	0.27	106,107,107,107	0
3	EDO	A	705	4/4	0.66	0.15	99,99,100,100	0
3	EDO	A	710	4/4	0.73	0.26	96,97,98,99	0
3	EDO	B	402	4/4	0.83	0.27	110,110,110,111	0
3	EDO	A	704	4/4	0.85	0.15	85,85,87,87	0
3	EDO	B	403	4/4	0.88	0.11	95,96,97,98	0
3	EDO	A	709	4/4	0.89	0.12	86,90,92,93	0
3	EDO	B	401	4/4	0.90	0.15	79,81,84,85	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	702	4/4	0.90	0.15	70,72,72,73	0
3	EDO	A	708	4/4	0.91	0.16	92,92,93,94	0
4	CL	A	713	1/1	0.91	0.13	90,90,90,90	0
4	CL	A	712	1/1	0.92	0.16	80,80,80,80	0
3	EDO	A	706	4/4	0.93	0.12	64,67,69,73	0
3	EDO	A	711	4/4	0.93	0.11	69,71,71,72	0
3	EDO	A	701	4/4	0.93	0.12	49,52,56,59	0
3	EDO	A	703	4/4	0.93	0.12	74,74,75,76	0
5	SAM	A	714	27/27	0.93	0.10	55,62,78,82	0
3	EDO	A	707	4/4	0.94	0.25	67,67,69,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.