



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 8, 2026 – 07:48 AM UTC

PDB ID : 2L29 / pdb_00002L29
BMRB ID : 17127
Title : Complex structure of E4 mutant human IGF2R domain 11 bound to IGF-II
Authors : Williams, C.; Hoppe, H.; Rezgui, D.; Strickland, M.; Frago, S.; Ellis, R.Z.;
Wattana-Amorn, P.; Prince, S.N.; Zaccheo, O.J.; Forbes, B.; Jones, E.Y.;
Crump, M.P.; Hassan, A.B.
Deposited on : 2010-08-13

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 84%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	142	
2	B	67	

2 Ensemble composition and analysis

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1515-A:1647, B:10-B:28, B:41-B:60 (172)	1.04	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 10, 12, 19, 20
2	3, 8, 13, 14, 15, 17, 18
3	9, 16
4	7, 11

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3146 atoms, of which 1548 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Insulin-like growth factor 2 receptor variant.

Mol	Chain	Residues	Atoms						Trace
1	A	142	Total	C	H	N	O	S	0
			2126	672	1049	186	208	11	

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1544	LYS	GLU	engineered mutation	UNP Q59EZ3
A	1545	SER	LYS	engineered mutation	UNP Q59EZ3
A	1547	VAL	LEU	engineered mutation	UNP Q59EZ3
A	1648	PRO	-	expression tag	UNP Q59EZ3
A	1649	GLU	-	expression tag	UNP Q59EZ3

- Molecule 2 is a protein called Insulin-like growth factor II.


Mol	Chain	Residues	Atoms						Trace
2	B	67	Total	C	H	N	O	S	0
			1020	321	499	93	101	6	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Insulin-like growth factor 2 receptor variant




- Molecule 2: Insulin-like growth factor II



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: Insulin-like growth factor 2 receptor variant



- Molecule 2: Insulin-like growth factor II





5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
TALOS	geometry optimization	
iCing	refinement	r765
ARIA	refinement	2.2
ARIA	structure solution	2.2
CNS	refinement	1.2
CNS	structure solution	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	2265
Number of shifts mapped to atoms	2265
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	84%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.20±0.04	2±1/1026 (0.2± 0.1%)	0.97±0.03	1±1/1393 (0.1± 0.1%)
2	B	1.13±0.08	1±1/309 (0.2± 0.3%)	1.04±0.05	0±0/417 (0.0± 0.0%)
All	All	1.18	46/26700 (0.2%)	0.98	17/36200 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.9±0.8
2	B	0.0±0.0	0.2±0.5
All	All	0	23

5 of 19 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	B	27	TYR	CE1-CZ	8.48	1.58	1.38	8	2
1	A	1592	TYR	CE2-CZ	-8.41	1.18	1.38	18	4
2	B	27	TYR	CE2-CZ	-8.40	1.18	1.38	8	2
1	A	1592	TYR	CE1-CZ	7.75	1.56	1.38	18	4
1	A	1542	TYR	CE1-CZ	7.49	1.56	1.38	8	6

5 of 6 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	1638	PHE	CA-CB-CG	5.74	119.54	113.80	12	4
1	A	1559	CYS	CA-C-N	5.45	125.99	120.38	9	5
1	A	1559	CYS	C-N-CA	5.45	125.99	120.38	9	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	1619	GLY	CA-C-N	5.18	124.89	119.82	12	1
1	A	1619	GLY	C-N-CA	5.18	124.89	119.82	12	1

There are no chirality outliers.

5 of 12 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	1606	TYR	Sidechain	6
1	A	1542	TYR	Sidechain	3
1	A	1592	TYR	Sidechain	3
2	B	24	ARG	Sidechain	2
2	B	59	TYR	Sidechain	2

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1002	983	981	16±4
2	B	303	276	276	3±2
All	All	26100	25180	25140	366

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

5 of 163 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:1629:LEU:HD13	1:A:1636:LEU:HG	0.87	1.44	13	10
1:A:1588:LEU:HB2	1:A:1612:PHE:HB2	0.85	1.45	1	19
1:A:1629:LEU:HG	1:A:1636:LEU:HG	0.78	1.54	7	3
1:A:1533:SER:HA	1:A:1536:ALA:HB2	0.77	1.56	19	1
1:A:1563:VAL:HG21	1:A:1573:SER:HB2	0.76	1.58	13	2

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/142 (94%)	119±3 (89±2%)	13±3 (10±2%)	1±1 (1±1%)	14	63
2	B	39/67 (58%)	34±2 (87±5%)	3±2 (9±4%)	2±1 (4±2%)	3	27
All	All	3440/4180 (82%)	3048 (89%)	329 (10%)	63 (2%)	9	52

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1560	PRO	17
2	B	42	ILE	16
2	B	60	CYS	14
1	A	1618	ALA	4
2	B	28	PHE	3

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/123 (93%)	108±1 (95±1%)	6±1 (5±1%)	24	75
2	B	33/57 (58%)	32±1 (97±2%)	1±1 (3±2%)	35	84
All	All	2940/3600 (82%)	2809 (96%)	131 (4%)	26	77

5 of 36 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1581	LEU	19
1	A	1636	LEU	17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	13	LEU	12
1	A	1626	LEU	11
1	A	1647	GLU	8

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 84% for the well-defined parts and 84% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2265
Number of shifts mapped to atoms	2265
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	198	-0.06 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	172	0.26 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}'$	178	0.21 ± 0.11	None needed (< 0.5 ppm)
^{15}N	188	0.48 ± 0.28	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 84%, i.e. 1862 atoms were assigned a chemical shift out of a possible 2214. 0 out of 29 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	801/859 (93%)	336/352 (95%)	306/344 (89%)	159/163 (98%)
Sidechain	928/1164 (80%)	637/760 (84%)	272/362 (75%)	19/42 (45%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	133/191 (70%)	82/92 (89%)	50/94 (53%)	1/5 (20%)
Overall	1862/2214 (84%)	1055/1204 (88%)	628/800 (78%)	179/210 (85%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	1572	ILE	HB	0.06	0.35 – 3.22	-6.0


7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	3071
Intra-residue ($ i-j =0$)	1109
Sequential ($ i-j =1$)	845
Medium range ($ i-j >1$ and $ i-j <5$)	338
Long range ($ i-j \geq 5$)	688
Inter-chain	42
Hydrogen bond restraints	49
Disulfide bond restraints	0
Total dihedral-angle restraints	318
Number of unmapped restraints	0
Number of restraints per residue	16.2
Number of long range restraints per residue ¹	3.5

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	88.2	0.2
0.2-0.5 (Medium)	186.9	0.5
>0.5 (Large)	191.8	4.61

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	6.2	5.72
10.0-20.0 (Medium)	0.1	14.02
>20.0 (Large)	None	None

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	1109	36.1	267	24.1	8.7	24	2.2	0.8
Backbone-Backbone	177	5.8	3	1.7	0.1	1	0.6	0.0
Backbone-Sidechain	644	21.0	211	32.8	6.9	17	2.6	0.6
Sidechain-Sidechain	288	9.4	53	18.4	1.7	6	2.1	0.2
Sequential (i-j =1)	845	27.5	305	36.1	9.9	26	3.1	0.8
Backbone-Backbone	322	10.5	35	10.9	1.1	0	0.0	0.0
Backbone-Sidechain	455	14.8	212	46.6	6.9	14	3.1	0.5
Sidechain-Sidechain	68	2.2	58	85.3	1.9	12	17.6	0.4
Medium range (i-j >1 & i-j <5)	338	11.0	168	49.7	5.5	11	3.3	0.4
Backbone-Backbone	86	2.8	9	10.5	0.3	0	0.0	0.0
Backbone-Sidechain	151	4.9	87	57.6	2.8	5	3.3	0.2
Sidechain-Sidechain	101	3.3	72	71.3	2.3	6	5.9	0.2
Long range (i-j ≥5)	688	22.4	383	55.7	12.5	29	4.2	0.9
Backbone-Backbone	108	3.5	10	9.3	0.3	1	0.9	0.0
Backbone-Sidechain	287	9.3	179	62.4	5.8	16	5.6	0.5
Sidechain-Sidechain	293	9.5	194	66.2	6.3	12	4.1	0.4
Inter-chain	42	1.4	35	83.3	1.1	3	7.1	0.1
Backbone-Backbone	2	0.1	1	50.0	0.0	0	0.0	0.0
Backbone-Sidechain	8	0.3	8	100.0	0.3	3	37.5	0.1
Sidechain-Sidechain	32	1.0	26	81.2	0.8	0	0.0	0.0
Hydrogen bond	49	1.6	3	6.1	0.1	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	3071	100.0	1161	37.8	37.8	93	3.0	3.0
Backbone-Backbone	695	22.6	58	8.3	1.9	2	0.3	0.1
Backbone-Sidechain	1594	51.9	700	43.9	22.8	55	3.5	1.8
Sidechain-Sidechain	782	25.5	403	51.5	13.1	36	4.6	1.2

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	108	126	78	153	10	475	0.53	3.09	0.43	0.42
2	105	122	68	151	12	458	0.53	4.31	0.46	0.44
3	110	128	66	134	9	447	0.5	2.98	0.35	0.42
4	111	139	72	152	8	482	0.52	3.31	0.41	0.41
5	114	122	63	151	12	462	0.6	4.56	0.54	0.44
6	106	134	74	149	11	474	0.51	2.83	0.36	0.44
7	107	125	73	174	7	486	0.54	3.27	0.47	0.41
8	104	124	72	155	10	465	0.51	3.02	0.39	0.4
9	99	124	67	167	15	472	0.51	3.18	0.38	0.41
10	107	126	74	149	13	469	0.5	2.23	0.36	0.4

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	112	116	79	153	8	468	0.54	3.49	0.44	0.43
12	114	138	63	153	13	481	0.54	3.03	0.43	0.43
13	110	136	70	159	9	484	0.52	3.29	0.4	0.42
14	114	132	72	155	16	489	0.53	3.57	0.4	0.44
15	106	136	72	146	14	474	0.54	4.03	0.48	0.42
16	109	132	70	160	16	487	0.49	3.04	0.38	0.4
17	108	114	70	149	11	452	0.52	4.61	0.44	0.42
18	103	132	70	156	13	474	0.53	2.89	0.4	0.43
19	113	127	72	163	9	484	0.48	2.24	0.33	0.4
20	113	121	63	169	11	477	0.52	3.71	0.44	0.41

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1864(IR:842, SQ:540, MR:170, LR:305, IC:7) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
49	70	24	63	7	213	1	5.0
26	33	14	49	2	124	2	10.0
27	24	12	27	5	95	3	15.0
12	12	9	24	4	61	4	20.0
18	11	12	20	3	64	5	25.0
11	12	7	17	1	48	6	30.0
17	11	4	10	1	43	7	35.0
8	15	6	14	1	44	8	40.0
9	8	11	10	1	39	9	45.0
5	8	8	18	2	41	10	50.0
4	7	5	8	0	24	11	55.0
3	5	5	8	1	22	12	60.0
5	7	3	13	0	28	13	65.0
7	8	6	10	0	31	14	70.0
7	5	4	6	3	25	15	75.0
7	10	6	8	0	31	16	80.0
7	10	7	14	0	38	17	85.0
9	13	7	18	1	48	18	90.0
12	10	7	17	0	46	19	95.0
24	26	11	29	3	93	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,924)	1:1520:A:ASN:HB3	1:1527:A:PHE:HE1	20	1.81	0.69	2.12
(1,271)	2:54:B:ALA:HB1	2:53:B:LEU:HD12	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB3	2:53:B:LEU:HD13	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB2	2:53:B:LEU:HD11	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB2	2:56:B:LEU:HD22	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB3	2:56:B:LEU:HD23	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB1	2:56:B:LEU:HD23	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB2	2:56:B:LEU:HD21	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB3	2:56:B:LEU:HD21	20	1.77	0.56	1.95
(1,271)	2:54:B:ALA:HB3	2:56:B:LEU:HD22	20	1.77	0.56	1.95
(1,2326)	2:32:B:ALA:HB2	2:31:B:PRO:HD3	20	1.73	0.3	1.85
(1,2326)	2:32:B:ALA:HB3	2:31:B:PRO:HD3	20	1.73	0.3	1.85
(1,2326)	2:32:B:ALA:HB1	2:31:B:PRO:HD3	20	1.73	0.3	1.85
(1,285)	2:17:B:LEU:HD13	2:27:B:TYR:HD2	20	1.41	1.37	0.9
(1,285)	2:17:B:LEU:HD13	2:27:B:TYR:HD1	20	1.41	1.37	0.9
(1,285)	2:17:B:LEU:HD11	2:27:B:TYR:HD1	20	1.41	1.37	0.9

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,285)	2:17:B:LEU:HD11	2:27:B:TYR:HD2	20	1.41	1.37	0.9
(1,285)	2:17:B:LEU:HD12	2:27:B:TYR:HD2	20	1.41	1.37	0.9
(1,285)	2:17:B:LEU:HD12	2:27:B:TYR:HD1	20	1.41	1.37	0.9
(1,320)	2:54:B:ALA:H	2:55:B:LEU:HD22	20	1.35	0.23	1.42
(1,320)	2:54:B:ALA:H	2:55:B:LEU:HD21	20	1.35	0.23	1.42
(1,320)	2:54:B:ALA:H	2:55:B:LEU:HD23	20	1.35	0.23	1.42
(1,320)	2:54:B:ALA:H	2:55:B:LEU:HD12	20	1.35	0.23	1.42
(1,505)	1:1618:A:ALA:HB3	1:1616:A:PRO:HD2	20	1.25	0.24	1.37
(1,505)	1:1618:A:ALA:HB2	1:1616:A:PRO:HD2	20	1.25	0.24	1.37
(1,505)	1:1618:A:ALA:HB1	1:1616:A:PRO:HD2	20	1.25	0.24	1.37
(1,364)	2:54:B:ALA:H	2:55:B:LEU:HD22	20	1.17	0.23	1.23
(1,364)	2:54:B:ALA:H	2:55:B:LEU:HD21	20	1.17	0.23	1.23
(1,364)	2:54:B:ALA:H	2:55:B:LEU:HD23	20	1.17	0.23	1.23
(1,364)	2:54:B:ALA:H	2:55:B:LEU:HD12	20	1.17	0.23	1.23
(1,3006)	2:56:B:LEU:HD11	2:42:B:ILE:HD12	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD13	2:42:B:ILE:HD11	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD13	2:42:B:ILE:HD12	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD12	2:42:B:ILE:HD12	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD13	2:42:B:ILE:HD13	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD12	2:42:B:ILE:HD11	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD11	2:42:B:ILE:HD11	20	1.12	0.37	1.0
(1,3006)	2:56:B:LEU:HD12	2:42:B:ILE:HD13	20	1.12	0.37	1.0
(1,2482)	2:13:B:LEU:HD11	2:12:B:GLU:HB3	20	1.11	0.54	1.17
(1,2482)	2:13:B:LEU:HD13	2:12:B:GLU:HB3	20	1.11	0.54	1.17
(1,2482)	2:13:B:LEU:HD12	2:12:B:GLU:HB3	20	1.11	0.54	1.17
(1,5)	1:1552:A:ILE:HD13	1:1529:A:LEU:HD13	20	1.07	0.19	1.06

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints ⓘ

9.5.1 Histogram : Distribution of distance violations ⓘ

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,285)	2:17:B:LEU:HD11	2:27:B:TYR:HD1	17	4.61
(1,285)	2:17:B:LEU:HD11	2:27:B:TYR:HD2	5	4.56
(1,2504)	2:13:B:LEU:HD22	2:9:B:CYS:HB3	2	4.31
(1,2450)	2:55:B:LEU:HD12	2:58:B:THR:HG23	15	4.03
(1,2537)	2:13:B:LEU:HD22	2:9:B:CYS:HB3	2	3.94
(1,285)	2:17:B:LEU:HD11	2:27:B:TYR:HD1	20	3.71
(1,2713)	2:51:B:CYS:H	2:55:B:LEU:HD21	15	3.65
(1,646)	1:1529:A:LEU:HD13	1:1612:A:PHE:HZ	14	3.57
(1,2508)	2:14:B:VAL:HG22	2:17:B:LEU:HD13	2	3.51
(1,2508)	2:14:B:VAL:HG22	2:17:B:LEU:HD11	11	3.49

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PHI	142	44.7	13	9.2	4.1	0	0.0	0.0
CHI1	34	10.7	2	5.9	0.6	0	0.0	0.0
PSI	142	44.7	16	11.3	5.0	1	0.7	0.3
Total	318	100.0	31	9.7	9.7	1	0.3	0.3

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations				Mean (°)	Max (°)	SD (°)	Median (°)
	PHI	CHI1	PSI	Total				
1	2	0	7	9	2.17	4.68	1.07	1.63
2	0	0	5	5	1.94	2.69	0.48	1.93
3	2	0	4	6	2.4	4.17	1.04	1.96
4	2	0	4	6	2.53	3.84	0.81	2.26
5	2	2	4	8	4.39	14.02	3.78	3.28
6	3	1	5	9	2.04	3.3	0.74	1.75
7	2	0	4	6	2.23	4.5	1.23	1.69
8	1	0	5	6	1.95	3.26	0.71	1.67
9	2	0	8	10	2.77	5.72	1.35	2.64
10	2	0	2	4	2.03	2.86	0.69	1.99
11	1	0	4	5	2.13	4.21	1.3	1.22
12	2	0	1	3	2.1	2.93	0.6	1.84
13	3	0	2	5	1.78	3.16	0.71	1.54
14	3	0	6	9	3.1	12.91	3.65	1.22
15	2	0	3	5	2.02	3.49	0.91	1.54
16	3	0	4	7	1.83	2.63	0.66	1.91
17	3	0	5	8	1.79	3.28	0.73	1.47
18	3	0	2	5	2.2	3.13	0.84	2.69
19	2	0	6	8	2.01	3.56	0.71	1.86
20	0	0	2	2	2.13	2.2	0.07	2.13

10.2.1 Bar graph : Dihedral violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints				Fraction of the ensemble	
PHI	CHI1	PSI	Total	Count ¹	%
5	1	5	11	1	5.0
1	1	1	3	2	10.0
1	0	4	5	3	15.0
3	0	2	5	4	20.0
1	0	0	1	5	25.0
1	0	0	1	6	30.0
1	0	0	1	7	35.0
0	0	0	0	8	40.0
0	0	1	1	9	45.0
0	0	1	1	10	50.0
0	0	0	0	11	55.0


Continued on next page...

Continued from previous page...

Number of violated restraints				Fraction of the ensemble	
PHI	CHI1	PSI	Total	Count ¹	%
0	0	0	0	12	60.0
0	0	0	0	13	65.0
0	0	0	0	14	70.0
0	0	0	0	15	75.0
0	0	0	0	16	80.0
0	0	1	1	17	85.0
0	0	0	0	18	90.0
0	0	0	0	19	95.0
0	0	1	1	20	100.0

¹ Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)




10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

10.4.1 Histogram : Distribution of mean dihedral-angle violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Medi
(1,137)	1:1542:A:TYR:N	1:1542:A:TYR:CA	1:1542:A:TYR:C	1:1543:A:SER:N	20	3.11	0.96	3.03
(1,246)	1:1623:A:ARG:N	1:1623:A:ARG:CA	1:1623:A:ARG:C	1:1624:A:PRO:N	17	2.34	0.84	2.47
(1,172)	1:1569:A:GLN:N	1:1569:A:GLN:CA	1:1569:A:GLN:C	1:1570:A:THR:N	10	2.05	0.74	1.73
(1,242)	1:1617:A:GLU:N	1:1617:A:GLU:CA	1:1617:A:GLU:C	1:1618:A:ALA:N	9	2.97	1.01	3.28
(1,159)	1:1558:A:ASN:C	1:1559:A:CYS:N	1:1559:A:CYS:CA	1:1559:A:CYS:C	7	1.53	0.26	1.53
(1,219)	1:1602:A:SER:C	1:1603:A:GLY:N	1:1603:A:GLY:CA	1:1603:A:GLY:C	6	1.38	0.27	1.38
(1,291)	1:1645:A:ALA:C	1:1646:A:CYS:N	1:1646:A:CYS:CA	1:1646:A:CYS:C	5	1.77	0.84	1.11
(1,45)	2:37:B:ARG:N	2:37:B:ARG:CA	2:37:B:ARG:C	2:38:B:ARG:N	4	3.11	0.33	3.08
(1,173)	1:1569:A:GLN:C	1:1570:A:THR:N	1:1570:A:THR:CA	1:1570:A:THR:C	4	2.54	0.54	2.48
(1,37)	2:30:B:ARG:N	2:30:B:ARG:CA	2:30:B:ARG:C	2:31:B:PRO:N	4	2.22	0.54	2.07

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,294)	1:1647:A:GLU:N	1:1647:A:GLU:CA	1:1647:A:GLU:C	1:1648:A:PRO:N	5	14.02
(1,294)	1:1647:A:GLU:N	1:1647:A:GLU:CA	1:1647:A:GLU:C	1:1648:A:PRO:N	14	12.91
(1,294)	1:1647:A:GLU:N	1:1647:A:GLU:CA	1:1647:A:GLU:C	1:1648:A:PRO:N	9	5.72
(1,137)	1:1542:A:TYR:N	1:1542:A:TYR:CA	1:1542:A:TYR:C	1:1543:A:SER:N	5	4.92
(1,137)	1:1542:A:TYR:N	1:1542:A:TYR:CA	1:1542:A:TYR:C	1:1543:A:SER:N	1	4.68
(1,137)	1:1542:A:TYR:N	1:1542:A:TYR:CA	1:1542:A:TYR:C	1:1543:A:SER:N	7	4.5
(1,242)	1:1617:A:GLU:N	1:1617:A:GLU:CA	1:1617:A:GLU:C	1:1618:A:ALA:N	14	4.39
(1,137)	1:1542:A:TYR:N	1:1542:A:TYR:CA	1:1542:A:TYR:C	1:1543:A:SER:N	11	4.21
(1,242)	1:1617:A:GLU:N	1:1617:A:GLU:CA	1:1617:A:GLU:C	1:1618:A:ALA:N	3	4.17
(1,137)	1:1542:A:TYR:N	1:1542:A:TYR:CA	1:1542:A:TYR:C	1:1543:A:SER:N	9	4.06