



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 26, 2026 – 06:44 AM UTC

PDB ID : 5L2E / pdb_0000512e
Title : Crystal structure of rat Glutamate receptor delta-2 extracellular domain
Authors : Cheng, S.; Ozkan, E.
Deposited on : 2016-08-01
Resolution : 4.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 4.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1077 (4.50-3.82)
Clashscore	190562	1019 (4.48-3.84)
Ramachandran outliers	187476	1035 (4.50-3.82)
Sidechain outliers	187428	1022 (4.50-3.82)
RSRZ outliers	180081	1074 (4.50-3.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	688	 74% 20% 6%
1	B	688	 77% 17% 6%
1	C	688	 75% 19% 6%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 15408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor ionotropic, delta-2, Glutamate receptor ionotropic, delta-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	648	5136	3248	889	973	26	0	0	0
1	B	648	5136	3248	889	973	26	0	0	0
1	C	648	5136	3248	889	973	26	0	0	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLY	-	expression tag	UNP Q63226
A	23	SER	-	expression tag	UNP Q63226
A	343	GLN	GLU	conflict	UNP Q63226
A	662	GLY	-	linker	UNP Q63226
A	663	THR	-	linker	UNP Q63226
A	814	HIS	-	expression tag	UNP Q63226
A	815	HIS	-	expression tag	UNP Q63226
A	816	HIS	-	expression tag	UNP Q63226
A	817	HIS	-	expression tag	UNP Q63226
A	818	HIS	-	expression tag	UNP Q63226
A	819	HIS	-	expression tag	UNP Q63226
B	22	GLY	-	expression tag	UNP Q63226
B	23	SER	-	expression tag	UNP Q63226
B	343	GLN	GLU	conflict	UNP Q63226
B	662	GLY	-	linker	UNP Q63226
B	663	THR	-	linker	UNP Q63226
B	814	HIS	-	expression tag	UNP Q63226
B	815	HIS	-	expression tag	UNP Q63226
B	816	HIS	-	expression tag	UNP Q63226
B	817	HIS	-	expression tag	UNP Q63226
B	818	HIS	-	expression tag	UNP Q63226
B	819	HIS	-	expression tag	UNP Q63226

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	22	GLY	-	expression tag	UNP Q63226
C	23	SER	-	expression tag	UNP Q63226
C	343	GLN	GLU	conflict	UNP Q63226
C	662	GLY	-	linker	UNP Q63226
C	663	THR	-	linker	UNP Q63226
C	814	HIS	-	expression tag	UNP Q63226
C	815	HIS	-	expression tag	UNP Q63226
C	816	HIS	-	expression tag	UNP Q63226
C	817	HIS	-	expression tag	UNP Q63226
C	818	HIS	-	expression tag	UNP Q63226
C	819	HIS	-	expression tag	UNP Q63226


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate receptor ionotropic, delta-2, Glutamate receptor ionotropic, delta-2




- Molecule 1: Glutamate receptor ionotropic, delta-2, Glutamate receptor ionotropic, delta-2



- Molecule 1: Glutamate receptor ionotropic, delta-2, Glutamate receptor ionotropic, delta-2

Chain C:  75% 19% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	179.17Å 179.17Å 214.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.58 – 4.15 46.58 – 4.15	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.58-4.15) 98.6 (46.58-4.15)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 4.14Å)	Xtrriage
Refinement program	PHENIX (dev_2481: ???)	Depositor
R, R_{free}	0.211 , 0.260 0.241 , 0.289	Depositor DCC
R_{free} test set	1973 reflections (6.46%)	wwPDB-VP
Wilson B-factor (Å ²)	50.0	Xtrriage
Anisotropy	0.049	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	0.378 for -h,-k,l	Xtrriage
Reported twinning fraction	0.490 for -h,-k,l	Depositor
Outliers	0 of 30109 reflections	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	15408	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.17	0/5235	0.39	0/7090
1	B	0.16	0/5235	0.38	0/7090
1	C	0.15	0/5235	0.36	0/7090
All	All	0.16	0/15705	0.38	0/21270

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5136	0	5057	85	0
1	B	5136	0	5057	73	1
1	C	5136	0	5057	77	1
All	All	15408	0	15171	233	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 233 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:458:GLU:HB2	1:C:466:LYS:HB2	1.67	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:ASP:O	1:A:395:ASN:ND2	2.21	0.74
1:C:103:GLY:HA2	1:C:133:LEU:HD21	1.69	0.74
1:B:40:ASP:OD1	1:B:324:ASN:ND2	2.22	0.72
1:C:39:LYS:NZ	1:C:319:ASN:O	2.22	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:755:ASP:O	1:C:345:ARG:NH1[6_656]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/688 (92%)	610 (96%)	26 (4%)	0	100	100
1	B	636/688 (92%)	611 (96%)	25 (4%)	0	100	100
1	C	636/688 (92%)	614 (96%)	22 (4%)	0	100	100
All	All	1908/2064 (92%)	1835 (96%)	73 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	568/604 (94%)	565 (100%)	3 (0%)	81	81
1	B	568/604 (94%)	565 (100%)	3 (0%)	81	81
1	C	568/604 (94%)	566 (100%)	2 (0%)	84	82
All	All	1704/1812 (94%)	1696 (100%)	8 (0%)	81	81

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	763	ASN
1	C	524	LEU
1	B	692	ARG
1	B	524	LEU
1	B	763	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	397	HIS
1	C	810	GLN
1	A	426	ASN
1	A	726	GLN
1	A	792	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	648/688 (94%)	-1.05	0 100 100	13, 49, 93, 119	0
1	B	648/688 (94%)	-1.02	0 100 100	21, 54, 102, 129	0
1	C	648/688 (94%)	-1.00	0 100 100	26, 60, 110, 128	0
All	All	1944/2064 (94%)	-1.02	0 100 100	13, 55, 103, 129	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.