



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 02:47 PM UTC

PDB ID : 3L49 / pdb_00003l49
Title : CRYSTAL STRUCTURE OF ABC SUGAR TRANSPORTER SUBUNIT
FROM *Rhodobacter sphaeroides* 2.4.1
Authors : Patskovsky, Y.; Ozyurt, S.; Dickey, M.; Do, J.; Wasserman, S.R.; Sauder, J.M.;
Burley, S.K.; Almo, S.C.; New York Structural GenomiX Research Consortium
(NYSGXRC); New York SGX Research Center for Structural Genomics (NYS-
GXRC)
Deposited on : 2009-12-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engl & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	
1	B	291	
1	C	291	
1	D	291	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC sugar (Ribose) transporter, periplasmic substrate-binding subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	2	0
			2152	1362	363	420	7			
1	B	282	Total	C	N	O	S	0	3	0
			2154	1364	362	421	7			
1	C	282	Total	C	N	O	S	0	3	0
			2158	1366	364	421	7			
1	D	282	Total	C	N	O	S	0	1	0
			2144	1357	360	420	7			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	MET	-	expression tag	UNP Q3J3W0
A	28	SER	-	expression tag	UNP Q3J3W0
A	310	GLU	-	expression tag	UNP Q3J3W0
A	311	GLY	-	expression tag	UNP Q3J3W0
A	312	HIS	-	expression tag	UNP Q3J3W0
A	313	HIS	-	expression tag	UNP Q3J3W0
A	314	HIS	-	expression tag	UNP Q3J3W0
A	315	HIS	-	expression tag	UNP Q3J3W0
A	316	HIS	-	expression tag	UNP Q3J3W0
A	317	HIS	-	expression tag	UNP Q3J3W0
B	27	MET	-	expression tag	UNP Q3J3W0
B	28	SER	-	expression tag	UNP Q3J3W0
B	310	GLU	-	expression tag	UNP Q3J3W0
B	311	GLY	-	expression tag	UNP Q3J3W0
B	312	HIS	-	expression tag	UNP Q3J3W0
B	313	HIS	-	expression tag	UNP Q3J3W0
B	314	HIS	-	expression tag	UNP Q3J3W0
B	315	HIS	-	expression tag	UNP Q3J3W0
B	316	HIS	-	expression tag	UNP Q3J3W0
B	317	HIS	-	expression tag	UNP Q3J3W0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	27	MET	-	expression tag	UNP Q3J3W0
C	28	SER	-	expression tag	UNP Q3J3W0
C	310	GLU	-	expression tag	UNP Q3J3W0
C	311	GLY	-	expression tag	UNP Q3J3W0
C	312	HIS	-	expression tag	UNP Q3J3W0
C	313	HIS	-	expression tag	UNP Q3J3W0
C	314	HIS	-	expression tag	UNP Q3J3W0
C	315	HIS	-	expression tag	UNP Q3J3W0
C	316	HIS	-	expression tag	UNP Q3J3W0
C	317	HIS	-	expression tag	UNP Q3J3W0
D	27	MET	-	expression tag	UNP Q3J3W0
D	28	SER	-	expression tag	UNP Q3J3W0
D	310	GLU	-	expression tag	UNP Q3J3W0
D	311	GLY	-	expression tag	UNP Q3J3W0
D	312	HIS	-	expression tag	UNP Q3J3W0
D	313	HIS	-	expression tag	UNP Q3J3W0
D	314	HIS	-	expression tag	UNP Q3J3W0
D	315	HIS	-	expression tag	UNP Q3J3W0
D	316	HIS	-	expression tag	UNP Q3J3W0
D	317	HIS	-	expression tag	UNP Q3J3W0

- Molecule 2 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			10	5	4	1		
2	B	1	Total	C	O	S	0	0
			10	5	4	1		
2	C	1	Total	C	O	S	0	0
			10	5	4	1		
2	D	1	Total	C	O	S	0	0
			10	5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	61	Total	O	0	0
			61	61		
3	B	55	Total	O	0	0
			55	55		
3	C	54	Total	O	0	0
			54	54		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	54	Total	O	0	0
			54	54		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ABC sugar (Ribose) transporter, periplasmic substrate-binding subunit




- Molecule 1: ABC sugar (Ribose) transporter, periplasmic substrate-binding subunit



- Molecule 1: ABC sugar (Ribose) transporter, periplasmic substrate-binding subunit



- Molecule 1: ABC sugar (Ribose) transporter, periplasmic substrate-binding subunit



P286	K279	Q263	V303	N309	GLU	GLY	HIS	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	67.31Å 75.71Å 77.44Å 106.41° 115.49° 96.71°	Depositor
Resolution (Å)	20.00 – 2.30 20.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.9 (20.00-2.30) 95.7 (20.00-2.30)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, R_{free}	0.193 , 0.254 0.194 , 0.257	Depositor DCC
R_{free} test set	1665 reflections (2.94%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8872	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	1/2200 (0.0%)	1.00	5/3003 (0.2%)
1	B	0.68	2/2206 (0.1%)	1.01	4/3012 (0.1%)
1	C	0.59	1/2209 (0.0%)	0.98	3/3015 (0.1%)
1	D	0.57	1/2189 (0.0%)	0.98	4/2989 (0.1%)
All	All	0.61	5/8804 (0.1%)	0.99	16/12019 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	227	VAL	CA-CB	7.49	1.57	1.54
1	B	163	VAL	CA-CB	6.38	1.60	1.53
1	A	227	VAL	CA-CB	6.34	1.57	1.54
1	B	244	ILE	C-O	-5.34	1.17	1.24
1	D	227	VAL	CA-CB	5.08	1.56	1.54

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	163	VAL	CA-C-N	7.72	127.27	119.24
1	B	163	VAL	C-N-CA	7.72	127.27	119.24
1	B	163	VAL	CB-CA-C	6.59	117.76	110.71
1	A	226	ASP	N-CA-C	6.21	117.71	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	163	VAL	CA-C-N	6.13	127.50	119.84
1	D	163	VAL	C-N-CA	6.13	127.50	119.84
1	A	43[A]	HIS	CA-C-O	5.94	128.17	121.51
1	A	43[B]	HIS	CA-C-O	5.94	128.17	121.51
1	C	163	VAL	CA-C-N	5.69	126.95	119.84
1	C	163	VAL	C-N-CA	5.69	126.95	119.84
1	C	44	ASP	N-CA-C	5.29	117.46	111.11
1	B	44	ASP	N-CA-C	5.23	117.38	111.11
1	A	225	TRP	CA-C-N	5.14	127.12	120.44
1	A	225	TRP	C-N-CA	5.14	127.12	120.44
1	D	65	ILE	CB-CA-C	5.06	117.46	111.59
1	D	30	GLU	N-CA-C	5.02	117.40	111.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	240	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2152	0	2134	12	0
1	B	2154	0	2133	8	0
1	C	2158	0	2142	7	0
1	D	2144	0	2121	10	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
3	A	61	0	0	0	0
3	B	55	0	0	0	0
3	C	54	0	0	0	0
3	D	54	0	0	0	0
All	All	8872	0	8530	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:MET:HE1	1:A:222:TRP:HD1	1.60	0.66
1:D:173:MET:HE1	1:D:185:ILE:HD11	1.77	0.66
1:A:144:MET:HE1	1:A:222:TRP:CD1	2.32	0.65
1:C:73:ASP:HB3	1:C:100:VAL:HG11	1.88	0.56
1:A:151:LYS:HG2	1:A:182:ASP:HB3	1.92	0.52
1:A:190:LEU:HD13	1:A:204:ASN:HB3	1.91	0.51
1:A:161:TYR:CZ	1:A:167:LYS:HG2	2.46	0.51
1:D:212:TYR:HB3	1:D:217:ASP:HB3	1.94	0.50
1:B:190:LEU:HD13	1:B:204:ASN:HB3	1.94	0.48
1:C:190:LEU:HD13	1:C:204:ASN:HB3	1.94	0.48
1:C:287:ARG:NH1	1:C:293:GLU:O	2.47	0.48
1:A:154:VAL:HG22	1:A:220:ALA:HB3	1.96	0.47
1:B:190:LEU:HD21	1:B:208:MET:HE3	1.96	0.47
1:A:227:VAL:HA	1:A:230:ILE:HD12	1.97	0.47
1:D:190:LEU:HD13	1:D:204:ASN:HB3	1.98	0.46
1:D:169:ARG:HD3	1:D:249:VAL:HG21	1.98	0.46
1:B:306:THR:H	1:B:309:ASN:HB2	1.79	0.46
1:C:99:ASP:OD2	1:C:99:ASP:N	2.49	0.45
1:D:151:LYS:HG3	1:D:182:ASP:HB3	1.99	0.45
1:A:83:LEU:O	1:A:88:PRO:HD3	2.16	0.45
1:D:169:ARG:NH1	1:D:224:CYS:O	2.46	0.45
1:B:145:VAL:HG23	1:B:183:VAL:HG21	1.99	0.45
1:B:212:TYR:HB3	1:B:217:ASP:HB3	1.99	0.44
1:A:105:LEU:HD22	1:A:115:LEU:HD21	2.00	0.44
1:C:264:SER:HA	1:C:265:PRO:HD3	1.91	0.43
1:A:145:VAL:HG23	1:A:183:VAL:HG21	2.00	0.43
1:D:209:LEU:HD11	1:D:236:LEU:HD23	2.01	0.43
1:B:264:SER:HA	1:B:265:PRO:HD3	1.84	0.42
1:C:144:MET:HE1	1:C:222:TRP:CD1	2.55	0.42
1:D:133:ASN:HA	1:D:136:ILE:HB	2.02	0.42
1:B:160:PHE:HB3	1:B:163:VAL:HG13	2.01	0.42
1:A:173:MET:HE3	1:A:173:MET:HB3	1.86	0.41
1:C:83:LEU:O	1:C:88:PRO:HD3	2.20	0.41
1:D:245:ARG:HA	1:D:265:PRO:O	2.20	0.41
1:B:142:LEU:HA	1:B:145:VAL:HG12	2.03	0.40
1:D:161:TYR:CZ	1:D:167:LYS:HD3	2.57	0.40
1:A:223:ALA:HB2	1:A:229:MET:HE3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/291 (97%)	273 (97%)	8 (3%)	1 (0%)	30	38
1	B	283/291 (97%)	276 (98%)	6 (2%)	1 (0%)	30	38
1	C	283/291 (97%)	278 (98%)	4 (1%)	1 (0%)	30	38
1	D	281/291 (97%)	268 (95%)	12 (4%)	1 (0%)	30	38
All	All	1129/1164 (97%)	1095 (97%)	30 (3%)	4 (0%)	30	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	250	ASP
1	B	250	ASP
1	C	250	ASP
1	D	250	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/235 (97%)	212 (93%)	17 (7%)	13	17
1	B	230/235 (98%)	216 (94%)	14 (6%)	17	24
1	C	230/235 (98%)	216 (94%)	14 (6%)	17	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	228/235 (97%)	212 (93%)	16 (7%)	14	19
All	All	917/940 (98%)	856 (93%)	61 (7%)	15	21

All (61) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	30	GLU
1	A	48	LYS
1	A	51	GLN
1	A	74	GLN
1	A	98	LEU
1	A	101	LEU
1	A	118	VAL
1	A	174	LYS
1	A	186	ILE
1	A	191	ARG
1	A	218	VAL
1	A	237	GLN
1	A	254	GLU
1	A	279	LYS
1	A	283	GLN
1	A	303	VAL
1	B	37	THR
1	B	58	GLU
1	B	63	THR
1	B	65	ILE
1	B	98	LEU
1	B	118	VAL
1	B	151	LYS
1	B	163	VAL
1	B	167	LYS
1	B	218	VAL
1	B	241	ARG
1	B	259	VAL
1	B	283	GLN
1	B	309	ASN
1	C	34	ILE
1	C	65	ILE
1	C	98	LEU
1	C	99	ASP
1	C	101	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	162	SER
1	C	173	MET
1	C	191	ARG
1	C	211	LYS
1	C	217	ASP
1	C	218	VAL
1	C	237	GLN
1	C	283	GLN
1	C	294	VAL
1	D	29	LEU
1	D	30	GLU
1	D	48	LYS
1	D	65	ILE
1	D	74	GLN
1	D	98	LEU
1	D	151	LYS
1	D	167	LYS
1	D	173	MET
1	D	184	LYS
1	D	214	ASN
1	D	221	ILE
1	D	237	GLN
1	D	279	LYS
1	D	283	GLN
1	D	303	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	ASN
1	A	128	ASN
1	A	234	GLN
1	A	292	GLN
1	B	51	GLN
1	B	53	GLN
1	B	76	GLN
1	B	79	GLN
1	B	81	GLN
1	B	128	ASN
1	B	158	ASN
1	B	234	GLN
1	B	237	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	292	GLN
1	B	309	ASN
1	C	53	GLN
1	C	81	GLN
1	C	128	ASN
1	C	196	ASN
1	C	237	GLN
1	C	273	GLN
1	D	53	GLN
1	D	127	ASN
1	D	128	ASN
1	D	172	GLN
1	D	234	GLN
1	D	237	GLN
1	D	288	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/291 (96%)	-0.32	1 (0%) 88 89	16, 33, 55, 84	2 (0%)
1	B	282/291 (96%)	-0.38	0 100 100	16, 33, 55, 76	3 (1%)
1	C	282/291 (96%)	-0.24	0 100 100	16, 34, 59, 81	3 (1%)
1	D	282/291 (96%)	-0.24	1 (0%) 88 89	15, 37, 61, 87	1 (0%)
All	All	1128/1164 (96%)	-0.30	2 (0%) 91 91	15, 34, 59, 87	9 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	216	GLY	2.3
1	A	309	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UNL	A	1	10/-	0.92	0.14	27,52,61,63	0
2	UNL	D	1	10/-	0.94	0.11	31,55,61,63	0
2	UNL	B	1	10/-	0.95	0.10	35,45,57,61	0
2	UNL	C	1	10/-	0.96	0.09	30,46,53,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.