



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 05:59 PM UTC

PDB ID : 5L44 / pdb_00005l44
Title : Structure of K-26-DCP in complex with the K-26 tripeptide
Authors : Masuyer, G.; Acharya, K.R.; Kramer, G.J.; Bachmann, B.O.
Deposited on : 2016-05-24
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	683	4% 91% 6%
1	B	683	2% 91% 6%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called K-26 dipeptidyl carboxypeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	662	Total	C	N	O	S	0	4	0
			5287	3358	922	997	10			
1	B	662	Total	C	N	O	S	0	3	0
			5274	3352	918	994	10			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is N-ACETYL-L-ILE-L-TYR-(R)-1-AMINO-2-(4-HYDROXYPHENYL)ETHYLPHOSPHONIC ACID (CCD ID: K26) (formula: $C_{25}H_{34}N_3O_8P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0
			37	25	3	8	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			37	25	3	8	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	506	Total	O	0	0
			506	506		
6	B	520	Total	O	0	0
			520	520		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: K-26 dipeptidyl carboxypeptidase



- Molecule 1: K-26 dipeptidyl carboxypeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.58Å 66.94Å 102.00Å 97.72° 90.71° 117.37°	Depositor
Resolution (Å)	33.00 – 1.75 33.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	89.2 (33.00-1.75) 89.2 (33.00-1.75)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.185 , 0.213 (Not available) , 0.238	Depositor DCC
R_{free} test set	6436 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å ²)	13.9	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.267 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11677	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K26, ZN, SO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	2/5432 (0.0%)	0.88	8/7399 (0.1%)
1	B	0.73	0/5422	0.90	4/7386 (0.1%)
All	All	0.74	2/10854 (0.0%)	0.89	12/14785 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	24	PRO	CA-C	5.78	1.55	1.51
1	A	35	VAL	CA-CB	5.45	1.57	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	254	PRO	N-CA-C	7.06	127.02	112.47
1	A	254	PRO	N-CA-C	6.70	126.27	112.47
1	B	253	VAL	CA-C-N	6.21	127.61	119.84
1	B	253	VAL	C-N-CA	6.21	127.61	119.84
1	A	253	VAL	CA-C-N	6.11	127.48	119.84
1	A	253	VAL	C-N-CA	6.11	127.48	119.84
1	A	229	GLN	CA-C-N	6.08	126.26	119.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	229	GLN	C-N-CA	6.08	126.26	119.32
1	B	128	ASP	N-CA-C	5.83	118.14	111.02
1	A	409	TYR	N-CA-C	5.50	118.70	109.95
1	A	23	LEU	CA-C-N	-5.36	115.75	119.66
1	A	23	LEU	C-N-CA	-5.36	115.75	119.66

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	253	VAL	Mainchain,Peptide
1	B	253	VAL	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5287	0	5132	25	0
1	B	5274	0	5124	25	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	5	0	0	1	0
4	B	5	0	0	1	0
5	A	37	0	32	3	0
5	B	37	0	31	0	0
6	A	506	0	0	7	0
6	B	520	0	0	4	0
All	All	11677	0	10319	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:467:HIS:CD2	6:A:1102:HOH:O	2.27	0.86
4:B:1004:SO4:O3	6:B:1101:HOH:O	1.97	0.82
5:A:1005:K26:OAJ	6:A:1102:HOH:O	2.03	0.77
4:A:1004:SO4:O2	6:A:1101:HOH:O	2.01	0.77
1:B:426:GLN:HE22	1:B:476:VAL:H	1.35	0.75
1:A:453:MET:HE1	1:A:461:MET:HE3	1.70	0.73
1:A:453:MET:HE1	1:A:461:MET:CE	2.19	0.73
1:A:426:GLN:HE22	1:A:476:VAL:H	1.38	0.69
1:A:395:ASN:HD21	1:A:635:ARG:HH22	1.41	0.68
1:B:395:ASN:HD21	1:B:635:ARG:HH22	1.43	0.66
1:B:99:LYS:HE3	1:B:176:GLN:HE22	1.62	0.65
1:A:159[A]:ARG:O	1:A:159[A]:ARG:HD2	2.02	0.59
1:B:355:LEU:CD1	1:B:461:MET:HE1	2.35	0.57
1:A:181:HIS:HD2	1:A:252:GLY:H	1.52	0.57
1:A:395:ASN:ND2	1:A:635:ARG:HH22	2.03	0.57
1:A:267:ARG:HH11	1:A:555:HIS:HE1	1.53	0.56
1:A:267:ARG:HH11	1:A:555:HIS:CE1	2.24	0.54
1:B:345:ASP:O	1:B:348:GLN:HG2	2.06	0.54
1:B:395:ASN:ND2	1:B:635:ARG:HH22	2.05	0.54
1:A:338:LEU:HD21	1:A:535:TYR:CD2	2.46	0.51
1:B:338:LEU:HD22	1:B:532:SER:HB2	1.93	0.51
1:B:355:LEU:HD12	1:B:461:MET:HE1	1.93	0.51
1:A:49:VAL:HG13	1:A:118:LEU:HD22	1.94	0.50
1:B:363:VAL:HG21	1:B:461:MET:HG2	1.93	0.50
1:B:181:HIS:HD2	1:B:252:GLY:H	1.60	0.49
1:B:267:ARG:HH11	1:B:555:HIS:HE1	1.60	0.49
1:A:593:HIS:HD2	6:A:1474:HOH:O	1.95	0.48
5:A:1005:K26:C	6:A:1102:HOH:O	2.61	0.48
1:B:338:LEU:HD21	1:B:535:TYR:CD2	2.49	0.48
1:B:46:LEU:HA	1:B:49:VAL:HG12	1.97	0.47
1:A:597:SER:OG	1:A:598:GLY:N	2.45	0.46
1:B:267:ARG:HH11	1:B:555:HIS:CE1	2.33	0.46
1:A:140:ARG:HD3	1:A:140:ARG:O	2.16	0.45
1:B:593:HIS:HD2	6:B:1483:HOH:O	1.99	0.45
1:B:245:HIS:HE1	6:B:1565:HOH:O	1.98	0.45
1:A:49:VAL:CG1	1:A:118:LEU:HD22	2.47	0.45
1:A:279:HIS:HE1	1:A:600:SER:OG	1.99	0.45
1:A:170:LEU:HD22	1:A:259:LEU:CD1	2.48	0.44
1:A:345:ASP:O	1:A:348:GLN:HG2	2.18	0.44
1:A:245:HIS:HE1	6:A:1547:HOH:O	1.99	0.43
1:A:298:MET:HE2	1:A:302:LEU:HD11	2.01	0.43
1:B:597:SER:OG	1:B:598:GLY:N	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:THR:HB	1:B:125:ILE:HD11	2.01	0.43
1:A:120:ALA:O	1:A:124:GLN:HG2	2.20	0.42
1:A:455:PHE:CD2	1:A:533:GLN:NE2	2.87	0.42
1:B:357:ARG:NH2	6:B:1106:HOH:O	2.39	0.42
1:A:593:HIS:CE1	5:A:1005:K26:HAR	2.55	0.41
1:B:279:HIS:HE1	1:B:600:SER:OG	2.02	0.41
1:B:49:VAL:HG23	1:B:118:LEU:HD22	2.03	0.41
1:A:124:GLN:HG3	6:A:1572:HOH:O	2.21	0.41
1:B:170:LEU:HD22	1:B:259:LEU:CD1	2.51	0.40
1:B:49:VAL:CG2	1:B:118:LEU:HD22	2.51	0.40
1:B:140:ARG:HD3	1:B:140:ARG:O	2.22	0.40
1:B:455:PHE:CD2	1:B:533:GLN:NE2	2.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	556/570 (98%)	554 (100%)	2 (0%)	84	80
1	B	555/570 (97%)	553 (100%)	2 (0%)	84	80
All	All	1111/1140 (98%)	1107 (100%)	4 (0%)	84	80

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	558	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	638	LEU
1	B	28	GLU
1	B	638	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (34) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	95	GLN
1	A	100	GLN
1	A	157	GLN
1	A	176	GLN
1	A	177	ASN
1	A	181	HIS
1	A	183	ASN
1	A	245	HIS
1	A	250	GLN
1	A	279	HIS
1	A	395	ASN
1	A	400	GLN
1	A	426	GLN
1	A	555	HIS
1	A	593	HIS
1	B	100	GLN
1	B	124	GLN
1	B	157	GLN
1	B	176	GLN
1	B	177	ASN
1	B	181	HIS
1	B	183	ASN
1	B	194	GLN
1	B	245	HIS
1	B	279	HIS
1	B	320	GLN
1	B	395	ASN
1	B	426	GLN
1	B	533	GLN
1	B	537	GLN
1	B	555	HIS
1	B	569	HIS
1	B	593	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	K26	A	1005	2	38,38,38	1.58	5 (13%)	49,53,53	1.42	7 (14%)
4	SO4	B	1004	-	4,4,4	0.67	0	6,6,6	0.56	0
5	K26	B	1005	2	38,38,38	1.39	5 (13%)	49,53,53	1.35	6 (12%)
4	SO4	A	1004	-	4,4,4	0.40	0	6,6,6	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	K26	A	1005	2	-	2/40/40/40	0/2/2/2
5	K26	B	1005	2	-	2/40/40/40	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1005	K26	PBK-OAK	-4.92	1.47	1.54
5	A	1005	K26	PBK-OAJ	-4.59	1.47	1.54
5	B	1005	K26	CB-CG	-3.62	1.42	1.51
5	B	1005	K26	PBK-OAJ	-3.60	1.49	1.54
5	A	1005	K26	CB-CG	-3.38	1.43	1.51
5	B	1005	K26	PBK-OAK	3.29	1.60	1.54
5	A	1005	K26	CAV-CBF	-3.23	1.43	1.51
5	A	1005	K26	PBK-CBI	3.00	1.88	1.84
5	B	1005	K26	CAV-CBF	-2.78	1.44	1.51
5	B	1005	K26	CBJ-NAX	2.04	1.50	1.45

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1005	K26	CG-CB-CA	-4.02	102.66	113.36
5	A	1005	K26	CG-CB-CA	-3.78	103.30	113.36
5	A	1005	K26	OAK-PBK-OAJ	3.58	117.31	107.58
5	A	1005	K26	CAV-CBF-CAS	3.35	127.14	120.90
5	B	1005	K26	CAV-CBI-NAY	-3.13	107.72	111.36
5	B	1005	K26	CAV-CBF-CAS	3.03	126.54	120.90
5	A	1005	K26	CAN-CAR-CBF	2.75	124.61	121.00
5	B	1005	K26	CAN-CAR-CBF	2.72	124.57	121.00
5	B	1005	K26	OAK-PBK-OAG	-2.57	107.00	113.45
5	A	1005	K26	OAK-PBK-OAG	-2.54	107.09	113.45
5	A	1005	K26	CAV-CBI-NAY	-2.11	108.90	111.36
5	B	1005	K26	C-CA-N	-2.06	105.54	111.11
5	A	1005	K26	CAV-CBF-CAR	-2.01	117.16	120.90

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1005	K26	CBI-CAV-CBF-CAR
5	A	1005	K26	CBI-CAV-CBF-CAS
5	B	1005	K26	CBI-CAV-CBF-CAR
5	B	1005	K26	CBI-CAV-CBF-CAS

There are no ring outliers.

3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1005	K26	3	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1004	SO4	1	0
4	A	1004	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	662/683 (96%)	0.27	26 (3%)	43 49	4, 15, 30, 46	4 (0%)
1	B	662/683 (96%)	0.15	14 (2%)	63 70	4, 14, 28, 43	3 (0%)
All	All	1324/1366 (96%)	0.21	40 (3%)	52 59	4, 15, 29, 46	7 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	671	LEU	4.6
1	A	124	GLN	3.6
1	A	533	GLN	3.5
1	A	116	ARG	3.5
1	B	534	LYS	3.4
1	B	535	TYR	3.3
1	A	535	TYR	3.2
1	A	58	ALA	3.1
1	B	533	GLN	2.8
1	B	671	LEU	2.8
1	B	129	GLY	2.8
1	A	59	PRO	2.7
1	A	559	THR	2.6
1	B	128	ASP	2.6
1	A	254	PRO	2.6
1	B	56	THR	2.6
1	B	558	GLN	2.6
1	A	10	GLU	2.6
1	A	114	LEU	2.5
1	A	152	LEU	2.5
1	A	534	LYS	2.5
1	B	10	GLU	2.5
1	A	273	LEU	2.4
1	A	558	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	562	GLU	2.4
1	B	127	PRO	2.2
1	A	128	ASP	2.2
1	A	560	PRO	2.2
1	A	51	ALA	2.1
1	A	154	ALA	2.1
1	B	532	SER	2.1
1	A	53	ALA	2.1
1	B	562	GLU	2.1
1	B	341	ARG	2.1
1	A	319	GLU	2.1
1	A	163	LEU	2.1
1	A	561	PRO	2.1
1	A	102	ILE	2.1
1	B	258	GLU	2.0
1	A	216	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	B	1003	1/1	0.92	0.07	25,25,25,25	0
4	SO4	A	1004	5/5	0.92	0.17	19,22,24,26	0
4	SO4	B	1004	5/5	0.94	0.14	19,22,24,25	0
5	K26	A	1005	37/37	0.95	0.09	8,14,23,27	0
5	K26	B	1005	37/37	0.95	0.09	10,15,22,28	0
3	MG	A	1003	1/1	0.96	0.06	27,27,27,27	0
3	MG	B	1002	1/1	0.99	0.04	8,8,8,8	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	1001	1/1	1.00	0.03	9,9,9,9	0
3	MG	A	1002	1/1	1.00	0.05	8,8,8,8	0
2	ZN	A	1001	1/1	1.00	0.03	9,9,9,9	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around K26 A 1005:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around K26 B 1005:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.