



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 02:54 PM UTC

PDB ID : 4L5T / pdb_00004l5t
Title : Crystal structure of the tetrameric p202 HIN2
Authors : Yin, Q.; Tian, Y.; Wu, H.
Deposited on : 2013-06-11
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	203	 62% 22% 5% 11%
1	B	203	 62% 27% • 9%
1	C	203	 53% 32% • 10%
1	D	203	 55% 29% 6% 10%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5748 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon-activable protein 202.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	0	0
			1414	905	229	271	9			
1	B	185	Total	C	N	O	S	0	0	0
			1463	935	240	277	11			
1	C	182	Total	C	N	O	S	0	0	0
			1438	919	236	272	11			
1	D	183	Total	C	N	O	S	0	0	0
			1433	913	237	272	11			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	243	SER	-	expression tag	UNP Q9R002
A	350	PRO	LEU	variant	UNP Q9R002
A	364	GLU	LYS	variant	UNP Q9R002
A	379	SER	THR	variant	UNP Q9R002
A	432	ALA	SER	variant	UNP Q9R002
B	243	SER	-	expression tag	UNP Q9R002
B	350	PRO	LEU	variant	UNP Q9R002
B	364	GLU	LYS	variant	UNP Q9R002
B	379	SER	THR	variant	UNP Q9R002
B	432	ALA	SER	variant	UNP Q9R002
C	243	SER	-	expression tag	UNP Q9R002
C	350	PRO	LEU	variant	UNP Q9R002
C	364	GLU	LYS	variant	UNP Q9R002
C	379	SER	THR	variant	UNP Q9R002
C	432	ALA	SER	variant	UNP Q9R002
D	243	SER	-	expression tag	UNP Q9R002
D	350	PRO	LEU	variant	UNP Q9R002
D	364	GLU	LYS	variant	UNP Q9R002
D	379	SER	THR	variant	UNP Q9R002
D	432	ALA	SER	variant	UNP Q9R002


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Interferon-activable protein 202



• Molecule 1: Interferon-activable protein 202



• Molecule 1: Interferon-activable protein 202



• Molecule 1: Interferon-activable protein 202





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.89Å 71.86Å 95.47Å 90.00° 102.38° 90.00°	Depositor
Resolution (Å)	46.39 – 3.40 46.39 – 3.40	Depositor EDS
% Data completeness (in resolution range)	97.9 (46.39-3.40) 93.9 (46.39-3.40)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 3.40Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.206 , 0.268 0.208 , 0.264	Depositor DCC
R_{free} test set	1118 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	80.4	Xtriage
Anisotropy	0.615	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 74.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5748	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	1/1441 (0.1%)	0.90	1/1943 (0.1%)
1	B	0.55	0/1491	0.92	2/2007 (0.1%)
1	C	0.51	0/1466	0.95	5/1975 (0.3%)
1	D	0.52	0/1460	0.99	6/1966 (0.3%)
All	All	0.53	1/5858 (0.0%)	0.94	14/7891 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	382	ASN	CA-C	-5.15	1.46	1.53

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	304	LYS	CA-C-N	10.46	137.34	120.60
1	D	304	LYS	C-N-CA	10.46	137.34	120.60
1	D	425	ALA	N-CA-C	-7.21	101.01	110.53
1	D	329	TYR	N-CA-C	-6.89	103.77	111.28
1	C	335	VAL	N-CA-C	6.24	117.27	108.17

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	329	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1414	0	1375	49	0
1	B	1463	0	1436	35	0
1	C	1438	0	1402	46	0
1	D	1433	0	1392	95	0
All	All	5748	0	5605	209	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 209 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:315:SER:N	1:D:331:SER:OG	1.58	1.35
1:D:302:LEU:HD11	1:D:328:GLU:O	1.33	1.26
1:D:282:PHE:CE1	1:D:303:GLU:CB	2.23	1.21
1:D:302:LEU:CD1	1:D:328:GLU:O	1.91	1.18
1:A:255:TYR:HB2	1:D:327:HIS:ND1	1.68	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/203 (87%)	164 (93%)	12 (7%)	0	100	100
1	B	181/203 (89%)	170 (94%)	11 (6%)	0	100	100
1	C	178/203 (88%)	165 (93%)	12 (7%)	1 (1%)	21	50
1	D	179/203 (88%)	166 (93%)	13 (7%)	0	100	100
All	All	714/812 (88%)	665 (93%)	48 (7%)	1 (0%)	48	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	321	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/182 (84%)	139 (91%)	14 (9%)	8	29
1	B	160/182 (88%)	144 (90%)	16 (10%)	7	26
1	C	156/182 (86%)	137 (88%)	19 (12%)	5	19
1	D	155/182 (85%)	141 (91%)	14 (9%)	9	30
All	All	624/728 (86%)	561 (90%)	63 (10%)	7	25

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	417	VAL
1	D	331	SER
1	C	323	SER
1	D	328	GLU
1	D	356	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	407	ASN
1	B	310	ASN
1	D	327	HIS
1	D	360	HIS
1	D	382	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	180/203 (88%)	-0.00	0	100 100	73, 114, 145, 153	0
1	B	185/203 (91%)	-0.16	0	100 100	60, 88, 116, 135	0
1	C	182/203 (89%)	-0.13	0	100 100	66, 89, 119, 133	0
1	D	183/203 (90%)	0.01	1 (0%)	87 78	84, 116, 142, 158	0
All	All	730/812 (89%)	-0.07	1 (0%)	92 91	60, 99, 141, 158	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	287	ALA	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.