



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 01:57 PM UTC

PDB ID : 5L5C / pdb_00005l5c
Title : Plexin A1 full extracellular region, domains 1 to 10, to 6 angstrom, spacegroup P4(3)2(1)2
Authors : Janssen, B.J.C.; Kong, Y.; Malinauskas, T.; Vangoor, V.R.; Coles, C.H.; Kaufmann, R.; Ni, T.; Gilbert, R.J.C.; Padilla-Parra, S.; Pasterkamp, R.J.; Jones, E.Y.
Deposited on : 2016-05-28
Resolution : 6.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 6.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1143 (8.00-4.00)
Clashscore	190562	1210 (8.00-4.00)
Ramachandran outliers	187476	1034 (8.00-4.00)
Sidechain outliers	187428	1000 (8.00-4.00)
RSRZ outliers	180081	1136 (8.00-4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1212	62% 68% 25% . .
2	B	4	25% 50% 25%
2	D	4	100%
3	C	5	20% 80%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	5	 100%
4	E	6	 67% 33%
4	I	6	 17% 83%
5	F	2	 100%
6	H	3	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	E	1	-	-	X	-
4	NAG	I	1	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 9546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Plexin-A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1171	9085	5719	1593	1715	58	0	0	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLU	-	expression tag	UNP P70206
A	35	THR	-	expression tag	UNP P70206
A	36	GLY	-	expression tag	UNP P70206
A	1237	ARG	-	expression tag	UNP P70206
A	1238	THR	-	expression tag	UNP P70206
A	1239	LYS	-	expression tag	UNP P70206
A	1240	HIS	-	expression tag	UNP P70206
A	1241	HIS	-	expression tag	UNP P70206
A	1242	HIS	-	expression tag	UNP P70206
A	1243	HIS	-	expression tag	UNP P70206
A	1244	HIS	-	expression tag	UNP P70206
A	1245	HIS	-	expression tag	UNP P70206

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	4	50	28	2	20	0	0	0
2	D	4	50	28	2	20	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	5	Total	C	N	O	0	0	0
			61	34	2	25			
3	G	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	6	Total	C	N	O	0	0	0
			72	40	2	30			
4	I	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	H	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Plexin-A1





- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 25% 50% 25%



- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 100%




- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 20% 80%



- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%




- Molecule 4: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  67% 33%




- Molecule 4: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  17% 83%




- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	198.15Å 198.15Å 228.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	119.50 – 6.00 119.50 – 6.00	Depositor EDS
% Data completeness (in resolution range)	97.0 (119.50-6.00) 97.7 (119.50-6.00)	Depositor EDS
R_{merge}	0.71	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 6.19Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.305 , 0.316 0.310 , 0.312	Depositor DCC
R_{free} test set	589 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	326.8	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 777.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	9546	wwPDB-VP
Average B, all atoms (Å ²)	283.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	5/9294 (0.1%)	1.12	34/12632 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	9

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	510	VAL	C-N	50.02	1.99	1.33
1	A	561	CYS	C-N	40.77	1.83	1.33
1	A	1043	THR	C-N	22.89	1.66	1.33
1	A	704	CYS	C-N	14.62	1.68	1.33
1	A	859	CYS	C-N	-11.57	1.18	1.33

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1043	THR	CA-C-N	-45.03	55.44	122.09
1	A	1043	THR	C-N-CA	-45.03	55.44	122.09
1	A	1043	THR	O-C-N	17.25	145.87	122.59
1	A	859	CYS	O-C-N	-17.05	100.62	123.12
1	A	561	CYS	O-C-N	14.40	140.53	122.81

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	410	ASN	Peptide
1	A	527	ASP	Peptide
1	A	807	CYS	Peptide
1	A	85	GLY	Peptide
1	A	94	TYR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9085	0	8873	266	38
2	B	50	0	43	1	0
2	D	50	0	43	0	0
3	C	61	0	51	0	0
3	G	61	0	52	0	0
4	E	72	0	61	22	0
4	I	72	0	61	0	0
5	F	28	0	25	0	0
6	H	39	0	34	0	0
7	A	28	0	26	0	0
All	All	9546	0	9269	267	38

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 267 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:704:CYS:C	1:A:705:PRO:N	1.68	1.44
1:A:661:VAL:HG13	4:E:1:NAG:C6	1.47	1.44
1:A:561:CYS:C	1:A:562:VAL:N	1.83	1.33
1:A:661:VAL:CG1	4:E:1:NAG:O6	1.88	1.20
1:A:510:VAL:C	1:A:511:GLU:N	1.99	1.20

The worst 5 of 38 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:GLN:NE2	1:A:921:GLU:N[5_454]	0.50	1.70
1:A:178:GLN:OE1	1:A:921:GLU:O[5_454]	0.98	1.22
1:A:178:GLN:NE2	1:A:921:GLU:CA[5_454]	0.98	1.22
1:A:931:HIS:CD2	1:A:982:SER:OG[8_444]	1.18	1.02
1:A:399:GLN:CB	1:A:963:ARG:NH1[4_545]	1.21	0.99

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1157/1212 (96%)	1049 (91%)	94 (8%)	14 (1%)	10 44

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	PRO
1	A	95	PRO
1	A	411	GLN
1	A	869	PRO
1	A	915	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1015/1051 (97%)	963 (95%)	52 (5%)	21 42

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	661	VAL
1	A	837	LEU
1	A	1033	LEU
1	A	683	VAL
1	A	812	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1027	ASN
1	A	983	HIS
1	A	764	GLN
1	A	931	HIS
1	A	729	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

35 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.81	0	17,19,21	1.74	3 (17%)
2	NAG	B	2	2	14,14,15	0.58	0	17,19,21	1.54	2 (11%)
2	BMA	B	3	2	11,11,12	0.63	0	15,15,17	0.72	0
2	MAN	B	4	2	11,11,12	0.64	0	15,15,17	0.94	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.62	0	17,19,21	1.21	1 (5%)
3	NAG	C	2	3	14,14,15	0.51	0	17,19,21	1.26	1 (5%)
3	BMA	C	3	3	11,11,12	0.53	0	15,15,17	0.58	0
3	MAN	C	4	3	11,11,12	0.66	0	15,15,17	0.88	1 (6%)
3	MAN	C	5	3	11,11,12	0.58	0	15,15,17	1.19	2 (13%)
2	NAG	D	1	1,2	14,14,15	0.67	0	17,19,21	1.27	1 (5%)
2	NAG	D	2	2	14,14,15	0.56	0	17,19,21	1.22	2 (11%)
2	BMA	D	3	2	11,11,12	0.81	1 (9%)	15,15,17	1.92	3 (20%)
2	MAN	D	4	2	11,11,12	0.57	0	15,15,17	0.99	1 (6%)
4	NAG	E	1	1,4	14,14,15	0.68	0	17,19,21	1.19	1 (5%)
4	NAG	E	2	4	14,14,15	0.53	0	17,19,21	1.16	2 (11%)
4	BMA	E	3	4	11,11,12	0.60	0	15,15,17	1.54	3 (20%)
4	MAN	E	4	4	11,11,12	0.62	0	15,15,17	0.87	1 (6%)
4	MAN	E	5	4	11,11,12	0.53	0	15,15,17	1.22	1 (6%)
4	MAN	E	6	4	11,11,12	0.59	0	15,15,17	0.84	1 (6%)
5	NAG	F	1	1,5	14,14,15	0.62	0	17,19,21	1.25	1 (5%)
5	NAG	F	2	5	14,14,15	0.53	0	17,19,21	1.08	1 (5%)
3	NAG	G	1	1,3	14,14,15	0.62	0	17,19,21	1.05	1 (5%)
3	NAG	G	2	3	14,14,15	0.67	0	17,19,21	1.12	1 (5%)
3	BMA	G	3	3	11,11,12	0.42	0	15,15,17	1.63	2 (13%)
3	MAN	G	4	3	11,11,12	0.55	0	15,15,17	0.87	1 (6%)
3	MAN	G	5	3	11,11,12	0.52	0	15,15,17	1.43	2 (13%)
6	NAG	H	1	1,6	14,14,15	0.59	0	17,19,21	1.10	1 (5%)
6	NAG	H	2	6	14,14,15	0.57	0	17,19,21	1.19	2 (11%)
6	BMA	H	3	6	11,11,12	0.47	0	15,15,17	1.13	2 (13%)
4	NAG	I	1	1,4	14,14,15	0.69	0	17,19,21	1.29	3 (17%)
4	NAG	I	2	4	14,14,15	0.53	0	17,19,21	1.46	3 (17%)
4	BMA	I	3	4	11,11,12	0.39	0	15,15,17	0.75	1 (6%)
4	MAN	I	4	4	11,11,12	0.61	0	15,15,17	0.89	0
4	MAN	I	5	4	11,11,12	0.62	0	15,15,17	1.13	1 (6%)
4	MAN	I	6	4	11,11,12	0.60	0	15,15,17	1.03	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
2	MAN	B	4	2	-	1/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	1/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	0/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	E	2	4	-	2/6/23/26	0/1/1/1
4	BMA	E	3	4	-	2/2/19/22	0/1/1/1
4	MAN	E	4	4	-	1/2/19/22	0/1/1/1
4	MAN	E	5	4	-	0/2/19/22	0/1/1/1
4	MAN	E	6	4	-	2/2/19/22	0/1/1/1
5	NAG	F	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	F	2	5	-	2/6/23/26	0/1/1/1
3	NAG	G	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	BMA	G	3	3	-	2/2/19/22	0/1/1/1
3	MAN	G	4	3	-	1/2/19/22	0/1/1/1
3	MAN	G	5	3	-	1/2/19/22	1/1/1/1
6	NAG	H	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	H	2	6	-	2/6/23/26	0/1/1/1
6	BMA	H	3	6	-	2/2/19/22	0/1/1/1
4	NAG	I	1	1,4	1/1/5/7	2/6/23/26	0/1/1/1
4	NAG	I	2	4	-	0/6/23/26	0/1/1/1
4	BMA	I	3	4	-	1/2/19/22	0/1/1/1
4	MAN	I	4	4	-	0/2/19/22	0/1/1/1
4	MAN	I	5	4	-	2/2/19/22	0/1/1/1
4	MAN	I	6	4	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	BMA	C2-C3	2.12	1.55	1.52

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	BMA	C1-C2-C3	6.15	118.60	109.64
2	B	1	NAG	C4-C3-C2	5.16	118.58	111.02
2	B	2	NAG	C4-C3-C2	4.90	118.20	111.02
3	G	3	BMA	C1-O5-C5	4.61	118.36	112.19
3	G	5	MAN	C1-O5-C5	4.53	118.25	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	I	1	NAG	C1

5 of 45 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	I	1	NAG	O5-C5-C6-O6
4	E	2	NAG	O5-C5-C6-O6
3	G	3	BMA	O5-C5-C6-O6
4	E	2	NAG	C4-C5-C6-O6
6	H	2	NAG	C4-C5-C6-O6



All (1) ring outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	G	5	MAN	C1-C2-C3-C4-C5-O5


3 monomers are involved in 23 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	2	NAG	1	0
2	B	1	NAG	1	0
4	E	1	NAG	22	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	A	1327	1	14,14,15	0.52	0	17,19,21	0.98	1 (5%)
7	NAG	A	1331	1	14,14,15	0.61	0	17,19,21	1.08	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	1327	1	-	1/6/23/26	0/1/1/1
7	NAG	A	1331	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1331	NAG	C1-O5-C5	3.34	116.67	112.19
7	A	1327	NAG	C4-C3-C2	2.24	114.31	111.02

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	1327	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	8

The worst 5 of 8 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	1147:ASP	C	1148:PRO	N	4.89
1	A	658:ASN	C	659:CYS	N	2.39
1	A	806:LYS	C	807:CYS	N	2.25
1	A	510:VAL	C	511:GLU	N	1.99
1	A	561:CYS	C	562:VAL	N	1.83

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1171/1212 (96%)	3.73	749 (63%) 0 2	235, 263, 432, 432	0

The worst 5 of 749 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1223	GLY	35.9
1	A	1225	GLU	26.6
1	A	1061	GLY	25.8
1	A	1218	THR	24.0
1	A	675	CYS	23.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	I	3	11/12	-0.07	0.18	431,431,431,431	0
4	BMA	E	3	11/12	-0.02	0.19	269,269,269,269	0
4	MAN	E	4	11/12	0.13	0.21	269,269,269,269	0
2	BMA	D	3	11/12	0.22	0.12	269,269,269,269	0
2	NAG	B	1	14/15	0.43	0.21	262,262,262,262	0
4	MAN	E	6	11/12	0.48	0.19	269,269,269,269	0
2	MAN	D	4	11/12	0.48	0.12	269,269,269,269	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MAN	C	4	11/12	0.51	0.17	269,269,269,269	0
2	NAG	B	2	14/15	0.55	0.13	262,262,262,262	0
2	BMA	B	3	11/12	0.58	0.11	262,262,262,262	0
4	MAN	I	6	11/12	0.64	0.22	431,431,431,431	0
3	MAN	G	4	11/12	0.68	0.13	252,252,252,252	0
5	NAG	F	1	14/15	0.68	0.15	252,252,252,252	0
3	BMA	G	3	11/12	0.71	0.13	252,252,252,252	0
5	NAG	F	2	14/15	0.71	0.14	252,252,252,252	0
6	NAG	H	1	14/15	0.73	0.18	252,252,252,252	0
6	NAG	H	2	14/15	0.75	0.20	252,252,252,252	0
3	MAN	G	5	11/12	0.78	0.21	252,252,252,252	0
2	MAN	B	4	11/12	0.81	0.25	262,262,262,262	0
3	NAG	G	2	14/15	0.81	0.32	252,252,252,252	0
3	NAG	C	2	14/15	0.82	0.13	269,269,269,269	0
6	BMA	H	3	11/12	0.83	0.13	252,252,252,252	0
3	MAN	C	5	11/12	0.86	0.11	269,269,269,269	0
4	MAN	I	4	11/12	0.86	0.31	431,431,431,431	0
4	MAN	E	5	11/12	0.87	0.21	269,269,269,269	0
3	NAG	C	1	14/15	0.87	0.16	269,269,269,269	0
3	NAG	G	1	14/15	0.88	0.26	252,252,252,252	0
3	BMA	C	3	11/12	0.89	0.17	269,269,269,269	0
2	NAG	D	2	14/15	0.90	0.15	269,269,269,269	0
4	NAG	I	2	14/15	0.90	0.27	431,431,431,431	0
2	NAG	D	1	14/15	0.91	0.24	269,269,269,269	0
4	NAG	I	1	14/15	0.92	0.52	431,431,431,431	0
4	MAN	I	5	11/12	0.92	0.49	431,431,431,431	0
4	NAG	E	2	14/15	0.98	0.16	269,269,269,269	0
4	NAG	E	1	14/15	0.99	0.16	269,269,269,269	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NAG	A	1331	14/15	0.30	0.27	268,268,268,268	0
7	NAG	A	1327	14/15	0.93	0.13	252,252,252,252	0

6.5 Other polymers [i](#)

There are no such residues in this entry.