



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:52 AM UTC

PDB ID : 9L55 / pdb_00009l55
Title : Plastid Localized Exonuclease 1 (D249A) complexed with DNA
Authors : Shi, G.; Zhang, Y.
Deposited on : 2024-12-22
Resolution : 2.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3557 (2.90-2.86)
Clashscore	190562	3801 (2.90-2.86)
Ramachandran outliers	187476	3699 (2.90-2.86)
Sidechain outliers	187428	3702 (2.90-2.86)
RSRZ outliers	180081	3558 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	19% 79% 11% 9%
1	B	331	12% 74% 18% 8%
2	E	18	11% 50% 50%
3	F	10	60% 40%
4	G	8	25% 75%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9806 atoms, of which 4555 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-3' exonuclease family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	300	Total	C	H	N	O	S	0	0	0
			4227	1400	2039	366	413	9			
1	B	303	Total	C	H	N	O	S	0	2	0
			4448	1463	2176	379	420	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	91	GLY	-	expression tag	UNP B4FJZ1
A	249	ALA	ASP	engineered mutation	UNP B4FJZ1
B	91	GLY	-	expression tag	UNP B4FJZ1
B	249	ALA	ASP	engineered mutation	UNP B4FJZ1

- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*AP*CP*CP*TP*CP*AP*CP*AP*CP*CP*AP*CP*TP*TP*AP*AP*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	E	18	Total	C	H	N	O	P	0	0	0
			542	173	182	61	108	18			

- Molecule 3 is a DNA chain called DNA (5'-D(P*AP*TP*TP*AP*AP*GP*TP*GP*GP*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	F	10	Total	C	H	N	O	P	0	0	0
			312	100	102	38	62	10			

- Molecule 4 is a DNA chain called DNA (5'-D(P*GP*TP*GP*AP*GP*GP*TP*A)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	G	8	Total	C	H	N	O	P	0	0	0
			227	80	56	34	49	8			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total 2	Mg 2	0	0
5	B	2	Total 2	Mg 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	16	Total 16	O 16	0	0
6	B	24	Total 24	O 24	0	0
6	F	3	Total 3	O 3	0	0
6	G	3	Total 3	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 5'-3' exonuclease family protein



• Molecule 1: 5'-3' exonuclease family protein




• Molecule 2: DNA (5'-D(P*TP*AP*CP*CP*TP*CP*AP*CP*AP*CP*CP*AP*CP*TP*TP*A P*AP*T)-3')



● Molecule 3: DNA (5'-D(P*AP*TP*TP*AP*AP*GP*TP*GP*GP*T)-3')



● Molecule 4: DNA (5'-D(P*GP*TP*GP*AP*GP*GP*TP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.45Å 104.40Å 117.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.38 – 2.87 30.38 – 2.87	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.38-2.87) 89.5 (30.38-2.87)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.72 (at 2.85Å)	Xtriage
Refinement program	PHENIX 1.21_5207	Depositor
R, R_{free}	0.269 , 0.315 0.269 , 0.316	Depositor DCC
R_{free} test set	1994 reflections (7.20%)	wwPDB-VP
Wilson B-factor (Å ²)	72.8	Xtriage
Anisotropy	0.690	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 89.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9806	wwPDB-VP
Average B, all atoms (Å ²)	105.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/2226	0.70	0/3025
1	B	0.30	0/2324	0.54	0/3153
2	E	0.56	0/401	0.90	0/612
3	F	0.51	0/235	0.75	0/360
4	G	0.53	0/192	0.75	0/294
All	All	0.41	0/5378	0.66	0/7444

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	2039	2079	37	0
1	B	2272	2176	2212	48	0
2	E	360	182	204	12	0
3	F	210	102	115	9	0
4	G	171	56	91	5	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	16	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	24	0	0	0	0
6	F	3	0	0	0	0
6	G	3	0	0	0	0
All	All	5251	4555	4701	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:ALA:HA	3:F:2:DT:OP1	1.53	1.08
1:A:250:LYS:HG2	3:F:2:DT:OP2	1.84	0.78
1:A:313:GLY:N	2:E:12:DA:H5'	2.01	0.76
1:A:248:PRO:O	3:F:2:DT:H5''	1.93	0.68
1:A:264:ARG:HH22	3:F:3:DT:P	2.17	0.67
1:A:143:LEU:HD12	1:A:380:PHE:CE2	2.29	0.67
2:E:2:DA:H62	4:G:7:DT:H3	1.45	0.65
1:A:248:PRO:O	3:F:2:DT:C5'	2.45	0.64
1:B:241:TYR:CE2	1:B:375:THR:HG21	2.33	0.64
1:B:107:TYR:HE1	1:B:198:THR:HG1	1.49	0.59
1:B:314:ASP:O	1:B:318:VAL:HG23	2.03	0.58
1:B:309:VAL:HG13	1:B:349:HIS:HB2	1.86	0.58
1:A:253:PHE:HA	1:A:256:LEU:HD23	1.87	0.56
1:B:324:PHE:CD2	1:B:330:LEU:HD13	2.41	0.56
1:B:312:ILE:HD12	1:B:312:ILE:N	2.22	0.55
1:A:192:ARG:HH22	3:F:1:DA:C5'	2.19	0.55
1:A:355:LEU:HD23	1:A:355:LEU:O	2.08	0.54
1:A:103:MET:HE2	1:A:202:GLY:C	2.32	0.54
1:A:140:LEU:HD22	1:A:412:ARG:HG2	1.90	0.54
2:E:1:DT:C6	2:E:1:DT:H5'	2.41	0.54
1:A:192:ARG:HH22	3:F:1:DA:H5''	1.73	0.54
1:A:132:ILE:HD11	1:A:202:GLY:HA2	1.89	0.53
1:B:253:PHE:HA	1:B:256:LEU:HD23	1.90	0.53
1:A:244:ARG:HD2	1:A:263:LEU:HD11	1.90	0.53
1:B:243:VAL:HB	1:B:260:LEU:HD22	1.91	0.53
1:B:406:VAL:HG23	1:B:410:ILE:HD11	1.92	0.52
1:B:327:LEU:CD1	1:B:331:LEU:HD21	2.39	0.51
1:B:241:TYR:HE2	1:B:375:THR:HG21	1.75	0.51
1:B:327:LEU:O	1:B:331:LEU:HD22	2.10	0.51
1:B:312:ILE:HD12	1:B:312:ILE:H	1.75	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:ARG:NH1	2:E:1:DT:OP1	2.38	0.50
1:B:287:LEU:HD11	1:B:295:VAL:HG21	1.93	0.50
1:A:139:LEU:O	1:A:143:LEU:HD23	2.11	0.50
1:B:330:LEU:HD11	1:B:346:LEU:CD1	2.42	0.49
1:A:104:TYR:CE1	1:A:199:VAL:HG21	2.47	0.49
1:A:409:ILE:HD12	1:A:409:ILE:N	2.29	0.48
2:E:8:DC:H1'	2:E:9:DA:H5'	1.95	0.48
1:B:275:PHE:CD1	1:B:279:ASP:HB2	2.50	0.47
1:B:391:ILE:HD11	1:B:410:ILE:HG23	1.95	0.47
1:B:350:SER:O	1:B:354:ILE:HD12	2.14	0.47
1:B:180:PHE:O	1:B:183[B]:MET:HG2	2.15	0.47
1:A:250:LYS:HB3	1:A:250:LYS:NZ	2.30	0.47
1:B:95:MET:HE1	1:B:142:MET:O	2.16	0.46
1:A:385:ASP:OD2	1:A:389:LYS:HG2	2.15	0.46
1:A:313:GLY:H	2:E:12:DA:H5'	1.78	0.46
1:B:316:ASN:O	1:B:317:ALA:C	2.58	0.46
1:A:218:ILE:HG23	1:A:373:PHE:CZ	2.51	0.46
1:A:139:LEU:HD13	1:A:143:LEU:HD23	1.97	0.45
1:A:313:GLY:HA3	2:E:12:DA:C5'	2.45	0.45
1:A:313:GLY:CA	2:E:12:DA:H5'	2.46	0.45
1:A:263:LEU:HD12	1:A:263:LEU:N	2.32	0.45
1:B:94:ILE:HD13	1:B:149:HIS:HB2	1.98	0.45
1:B:200:VAL:O	1:B:203:MET:HB3	2.17	0.45
1:B:112:ALA:O	1:B:115:GLN:HG3	2.17	0.45
1:B:143:LEU:HD22	1:B:380:PHE:CZ	2.52	0.45
1:B:327:LEU:HD23	1:B:357:LYS:CD	2.47	0.45
1:B:407:ASN:HA	1:B:410:ILE:HD12	1.99	0.45
4:G:7:DT:H2''	4:G:8:DA:C8	2.52	0.44
1:A:370:MET:HE2	1:A:370:MET:HB3	1.84	0.44
1:A:313:GLY:HA3	2:E:12:DA:H5''	2.00	0.43
2:E:6:DC:H2''	2:E:7:DA:H5'	2.00	0.43
1:A:250:LYS:HB3	1:A:250:LYS:HZ3	1.83	0.43
3:F:10:DT:H2''	4:G:1:DG:O4'	2.17	0.43
1:B:287:LEU:CD2	1:B:291:GLN:HB2	2.49	0.43
1:B:178:MET:HG2	1:B:183[A]:MET:HE3	2.01	0.43
1:A:274:SER:O	1:A:275:PHE:C	2.61	0.43
1:A:407:ASN:N	1:A:408:PRO:HD2	2.34	0.43
1:A:249:ALA:CA	3:F:2:DT:OP1	2.44	0.42
1:A:367:PRO:HG2	1:A:370:MET:HG3	2.00	0.42
1:B:316:ASN:O	1:B:319:LYS:N	2.52	0.42
1:B:330:LEU:HD21	1:B:346:LEU:HD11	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:3:DC:H2''	2:E:4:DC:H5'	2.00	0.42
4:G:5:DG:H2''	4:G:6:DG:H5'	2.01	0.42
1:B:99:GLY:HA3	1:B:154:PHE:CE1	2.55	0.42
1:A:185:TYR:CE2	1:A:187:ALA:HB3	2.55	0.42
1:B:275:PHE:CE1	1:B:279:ASP:HB2	2.54	0.42
1:A:178:MET:HE2	1:A:183:MET:CE	2.50	0.41
1:B:358:ASN:OD1	1:B:363:ARG:NH2	2.53	0.41
1:A:280:PHE:CZ	1:A:289:PRO:HA	2.55	0.41
1:B:183[A]:MET:HA	1:B:183[A]:MET:HE2	2.02	0.41
1:B:327:LEU:HD12	1:B:331:LEU:CD2	2.50	0.41
1:B:327:LEU:HD11	1:B:331:LEU:HD21	2.03	0.41
2:E:1:DT:H5'	2:E:1:DT:H6	1.84	0.41
1:B:250:LYS:O	1:B:253:PHE:HB2	2.20	0.41
1:B:315:ILE:CB	4:G:4:DA:OP2	2.69	0.41
1:B:324:PHE:CE2	1:B:330:LEU:HD13	2.56	0.41
1:B:327:LEU:O	1:B:330:LEU:HB3	2.20	0.41
1:A:264:ARG:HE	1:A:264:ARG:HB3	1.61	0.40
1:B:100:THR:HG21	1:B:155:ASP:HB2	2.03	0.40
1:B:241:TYR:CD2	1:B:375:THR:HG21	2.56	0.40
1:B:256:LEU:HB2	1:B:289:PRO:HB3	2.03	0.40
1:B:327:LEU:HD23	1:B:357:LYS:HD2	2.04	0.40
1:B:355:LEU:HD23	1:B:355:LEU:O	2.21	0.40
1:B:143:LEU:HB3	1:B:416:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/331 (88%)	283 (98%)	7 (2%)	0	100	100
1	B	297/331 (90%)	292 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	587/662 (89%)	575 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/281 (77%)	214 (99%)	2 (1%)	70	88
1	B	234/281 (83%)	234 (100%)	0	100	100
All	All	450/562 (80%)	448 (100%)	2 (0%)	84	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	250	LYS
1	A	398	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	305	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	300/331 (90%)	1.21	62 (20%) 2 2	49, 98, 154, 187	0
1	B	303/331 (91%)	1.00	41 (13%) 7 5	44, 102, 139, 163	1 (0%)
2	E	18/18 (100%)	1.13	2 (11%) 10 9	92, 128, 154, 155	0
3	F	10/10 (100%)	0.86	0 100 100	103, 121, 142, 143	0
4	G	8/8 (100%)	1.43	0 100 100	88, 111, 152, 160	0
All	All	639/698 (91%)	1.11	105 (16%) 4 4	44, 101, 146, 187	1 (0%)

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	183[A]	MET	4.8
1	A	407	ASN	4.7
1	A	128	TRP	4.6
1	B	407	ASN	4.2
1	A	213	MET	4.1
1	B	264	ARG	4.1
1	A	114	LEU	4.0
1	A	94	ILE	4.0
1	A	107	TYR	4.0
1	B	125	ASN	3.9
1	B	262	LEU	3.9
1	B	110	ILE	3.9
1	A	421	LYS	3.7
1	A	310	GLU	3.6
1	A	368	HIS	3.6
1	A	125	ASN	3.6
1	B	337	VAL	3.5
1	A	400	TYR	3.5
1	A	302	LYS	3.5
1	A	312	ILE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	397	LEU	3.5
1	B	362	LEU	3.5
1	A	337	VAL	3.4
1	A	346	LEU	3.3
1	B	347	ILE	3.3
1	B	421	LYS	3.3
1	A	305	ASN	3.2
1	A	239	ALA	3.2
1	A	271	GLY	3.2
1	A	308	GLY	3.2
1	B	314	ASP	3.1
1	A	410	ILE	3.1
1	A	91	GLY	3.1
1	B	302	LYS	3.0
1	B	306	ILE	3.0
1	A	272	MET	2.9
1	A	297	ALA	2.9
1	B	201	GLN	2.9
1	A	311	GLY	2.9
1	B	132	ILE	2.9
1	A	413	ALA	2.9
1	B	202	GLY	2.8
1	A	108	TYR	2.8
2	E	3	DC	2.8
1	A	106	SER	2.8
1	A	334	VAL	2.7
1	A	399	ALA	2.7
1	A	274	SER	2.7
1	A	129	VAL	2.7
1	A	394	LEU	2.7
1	A	406	VAL	2.6
1	B	105	ARG	2.6
1	A	204	GLN	2.6
1	A	417	TRP	2.6
1	B	176	LYS	2.6
1	B	311	GLY	2.6
1	B	272	MET	2.5
1	A	414	ALA	2.5
1	B	247	SER	2.5
1	B	274	SER	2.5
1	A	349	HIS	2.5
1	A	200	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	176	LYS	2.5
1	A	335	ASP	2.5
1	A	347	ILE	2.5
1	A	202	GLY	2.4
1	A	419	LYS	2.4
1	A	130	LEU	2.3
1	A	409	ILE	2.3
2	E	11	DC	2.3
1	A	327	LEU	2.3
1	B	139	LEU	2.3
1	A	391	ILE	2.3
1	B	193	THR	2.3
1	B	195	THR	2.3
1	A	242	LYS	2.3
1	B	303	ALA	2.2
1	A	343	LYS	2.2
1	A	390	PHE	2.2
1	A	404	SER	2.2
1	B	111	LEU	2.2
1	B	381	LYS	2.2
1	A	341	ARG	2.2
1	B	91	GLY	2.2
1	B	299	SER	2.2
1	A	416	LEU	2.2
1	A	187	ALA	2.2
1	B	129	VAL	2.2
1	B	383	PRO	2.2
1	B	191	ASN	2.1
1	B	198	THR	2.1
1	B	204	GLN	2.1
1	B	399	ALA	2.1
1	A	248	PRO	2.1
1	A	325	GLY	2.1
1	B	252	PHE	2.1
1	B	261	ARG	2.1
1	B	200	VAL	2.0
1	B	334	VAL	2.0
1	A	157	ASP	2.0
1	A	314	ASP	2.0
1	B	321	ILE	2.0
1	A	101	SER	2.0
1	A	387	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	303	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	A	501	1/1	0.65	0.38	98,98,98,98	0
5	MG	A	502	1/1	0.85	0.15	95,95,95,95	0
5	MG	B	502	1/1	0.91	0.12	143,143,143,143	0
5	MG	B	501	1/1	0.95	0.07	92,92,92,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.