



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:11 AM UTC

PDB ID : 1L7P / pdb_00001l7p
Title : SUBSTRATE BOUND PHOSPHOSERINE PHOSPHATASE COMPLEX
STRUCTURE
Authors : Wang, W.; Cho, H.S.; Kim, R.; Jancarik, J.; Yokota, H.; Nguyen, H.H.; Grig-
oriev, I.V.; Wemmer, D.E.; Kim, S.H.; Berkeley Structural Genomics Center
(BSGC)
Deposited on : 2002-03-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	3% 66% 29% ..
1	B	211	5% 71% 26% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHOSPHOSERINE PHOSPHATASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	S	Se	0	0	0
			1630	1039	275	312	2	2			
1	B	210	Total	C	N	O	S	Se	0	0	0
			1648	1050	278	316	2	2			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP Q58989
A	11	ASN	ASP	engineered mutation	UNP Q58989
A	43	MSE	MET	modified residue	UNP Q58989
A	174	MSE	MET	modified residue	UNP Q58989
B	501	MSE	MET	modified residue	UNP Q58989
B	511	ASN	ASP	engineered mutation	UNP Q58989
B	543	MSE	MET	modified residue	UNP Q58989
B	674	MSE	MET	modified residue	UNP Q58989

- Molecule 2 is PHOSPHOSERINE (CCD ID: SEP) (formula: C₃H₈NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			11	3	1	6	1		
2	B	1	Total	C	N	O	P	0	0
			11	3	1	6	1		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	100	Total 100	O 100	0	0
4	B	95	Total 95	O 95	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHOSPHOSERINE PHOSPHATASE



• Molecule 1: PHOSPHOSERINE PHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	70.80Å 70.43Å 90.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.90 15.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (15.00-1.90) 88.8 (15.00-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.89Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.229 , 0.266 0.234 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.470	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.033 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3505	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1640	0.94	10/2185 (0.5%)
1	B	0.36	0/1658	0.89	5/2208 (0.2%)
All	All	0.37	0/3298	0.91	15/4393 (0.3%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	511	ASN	N-CA-C	-8.75	99.24	110.53
1	A	11	ASN	N-CA-C	-8.51	99.56	110.53
1	A	94	VAL	N-CA-C	-8.21	96.75	108.58
1	A	100	GLY	N-CA-C	-6.88	104.80	115.73
1	B	600	GLY	N-CA-C	-6.83	104.53	115.66
1	A	143	ALA	N-CA-C	6.74	119.24	111.02
1	B	707	LEU	N-CA-C	5.95	118.53	111.33
1	A	195	ASP	N-CA-C	-5.92	104.41	111.69
1	B	599	SER	N-CA-C	5.33	118.15	109.24
1	A	210	ILE	N-CA-C	5.32	116.96	108.87
1	B	686	ALA	N-CA-C	5.30	118.27	110.59
1	A	99	SER	N-CA-C	5.26	118.02	109.24
1	A	186	ALA	N-CA-C	5.17	117.98	110.17
1	A	49	PHE	N-CA-C	5.14	116.57	111.07
1	A	194	ALA	N-CA-C	5.03	117.93	110.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1630	0	1738	49	0
1	B	1648	0	1757	45	0
2	A	11	0	5	0	0
2	B	11	0	5	0	0
3	B	10	0	0	0	0
4	A	100	0	0	2	0
4	B	95	0	0	2	0
All	All	3505	0	3505	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:LYS:HE3	1:A:150:LYS:HA	1.40	1.00
1:B:543:MSE:HE2	1:B:543:MSE:HA	1.56	0.86
1:A:4:LYS:O	1:A:5:LYS:HB2	1.80	0.79
1:B:672:ILE:HG23	1:B:693:LYS:HE3	1.67	0.77
1:B:570:LYS:HA	1:B:573:LYS:HE3	1.68	0.75
1:B:667:ASP:HA	1:B:686:ALA:HB2	1.70	0.73
1:A:6:LYS:NZ	1:A:211:LYS:HG3	2.07	0.70
1:B:540:LYS:O	1:B:544:GLU:HG3	1.96	0.66
1:A:149:GLU:HG3	1:A:177:LYS:HD3	1.77	0.65
1:A:167:ASP:HA	1:A:186:ALA:HB2	1.80	0.64
1:B:676:LYS:NZ	1:B:676:LYS:HB2	2.13	0.64
1:B:581:ALA:HA	1:B:703:LEU:HD12	1.80	0.63
1:A:66:GLU:HG2	1:A:70:LYS:NZ	2.14	0.62
1:B:672:ILE:HG23	1:B:693:LYS:CE	2.29	0.62
1:B:518:ASN:HB2	1:B:576:THR:HG23	1.82	0.62
1:A:198:ILE:HD12	1:A:206:ILE:HG22	1.83	0.61
1:A:158:ASN:OD1	1:A:160:GLU:HG2	2.01	0.61
1:A:200:LYS:HD3	1:A:205:GLU:OE2	2.01	0.61
1:A:203:LEU:O	1:A:206:ILE:HG12	2.01	0.61
1:A:150:LYS:HE2	4:A:932:HOH:O	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:646:GLU:HG2	1:B:650:LYS:HE3	1.84	0.60
1:A:64:PRO:HD2	1:A:67:LYS:HG3	1.85	0.57
1:B:606:VAL:O	1:B:609:ILE:HG22	2.04	0.57
1:B:667:ASP:HA	1:B:686:ALA:CB	2.35	0.56
1:B:524:GLU:OE1	1:B:574:ARG:HD3	2.06	0.56
1:B:570:LYS:HG3	1:B:573:LYS:NZ	2.21	0.55
1:A:150:LYS:HE3	1:A:150:LYS:CA	2.27	0.55
1:A:80:GLY:HA2	1:A:83:GLU:OE2	2.07	0.54
1:A:198:ILE:CD1	1:A:206:ILE:HG22	2.37	0.54
1:B:543:MSE:HA	1:B:543:MSE:CE	2.34	0.54
1:B:648:LEU:C	1:B:648:LEU:HD23	2.33	0.54
1:B:566:GLU:H	1:B:566:GLU:CD	2.16	0.54
1:A:150:LYS:HA	1:A:150:LYS:CE	2.22	0.53
1:A:69:GLU:OE2	1:A:104:ILE:HD11	2.09	0.53
1:A:150:LYS:N	1:A:150:LYS:HD2	2.25	0.52
1:A:172:ILE:HD13	1:A:190:LEU:HD12	1.93	0.51
1:B:658:ASN:OD1	1:B:660:GLU:HG2	2.10	0.51
1:A:61:LYS:O	1:A:62:ASP:HB2	2.11	0.50
1:B:574:ARG:HG3	1:B:574:ARG:O	2.10	0.50
1:B:518:ASN:HA	1:B:701:ARG:HH21	1.77	0.49
1:B:507:LEU:HG	1:B:509:LEU:HD11	1.94	0.49
1:B:653:LYS:HE3	4:B:848:HOH:O	2.11	0.49
1:B:675:PHE:HE2	1:B:694:ALA:HB2	1.78	0.49
1:A:85:ILE:HD12	1:A:95:VAL:HG11	1.94	0.49
1:A:6:LYS:HB3	1:A:93:TYR:CE1	2.48	0.49
1:B:568:VAL:O	1:B:572:ILE:HG12	2.12	0.49
1:B:537:LYS:HG2	1:B:541:GLU:OE2	2.12	0.48
1:B:672:ILE:CG2	1:B:693:LYS:HE3	2.41	0.48
1:A:113:LEU:HB2	1:A:115:LEU:HD13	1.95	0.48
1:B:599:SER:O	1:B:620:ALA:HA	2.14	0.48
1:A:83:GLU:H	1:A:83:GLU:CD	2.22	0.48
1:B:684:PHE:O	1:B:685:CYS:C	2.57	0.48
1:A:6:LYS:HZ2	1:A:211:LYS:HG3	1.78	0.48
1:B:509:LEU:HD12	1:B:509:LEU:N	2.30	0.47
1:A:11:ASN:OD1	1:A:166:GLY:HA2	2.13	0.47
1:B:574:ARG:HG3	1:B:574:ARG:HH11	1.79	0.47
1:B:570:LYS:O	1:B:573:LYS:HG2	2.15	0.47
1:A:5:LYS:HB3	1:A:157:ILE:HG12	1.97	0.46
1:A:149:GLU:OE1	1:A:177:LYS:NZ	2.48	0.46
1:A:167:ASP:HA	1:A:186:ALA:CB	2.46	0.46
1:B:633:ASP:HB2	4:B:929:HOH:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:582:GLU:O	1:B:586:LYS:HG2	2.17	0.45
1:A:9:LEU:HD23	1:A:98:VAL:HG21	1.99	0.45
1:A:148:LEU:HD23	1:A:148:LEU:C	2.42	0.45
1:B:528:GLU:OE1	1:B:570:LYS:HD3	2.16	0.45
1:B:664:ALA:HB1	1:B:674:MSE:HG2	1.98	0.45
1:B:676:LYS:HB2	1:B:676:LYS:HZ2	1.82	0.45
1:B:550:GLU:HG3	1:B:639:LEU:HB3	1.99	0.44
1:A:125:VAL:HG22	1:A:130:LEU:HD23	1.99	0.44
1:B:582:GLU:HG2	1:B:613:LEU:HD22	1.99	0.44
1:B:645:GLY:O	1:B:649:GLU:HG3	2.17	0.44
1:A:150:LYS:CA	1:A:150:LYS:CE	2.93	0.44
1:A:169:ALA:O	1:A:172:ILE:HG12	2.18	0.43
1:B:565:ILE:HD11	1:B:625:VAL:CG1	2.48	0.43
1:B:599:SER:C	1:B:601:GLY:H	2.26	0.43
1:A:99:SER:C	1:A:101:GLY:H	2.27	0.43
1:B:566:GLU:CD	1:B:566:GLU:N	2.77	0.43
1:A:66:GLU:HG2	1:A:70:LYS:HZ3	1.82	0.42
1:A:184:PHE:O	1:A:185:CYS:C	2.62	0.42
1:B:543:MSE:HE1	1:B:670:ASN:OD1	2.20	0.42
1:A:150:LYS:CE	4:A:932:HOH:O	2.63	0.41
1:A:27:ARG:HA	1:A:32:GLU:HB2	2.02	0.41
1:B:574:ARG:HG3	1:B:574:ARG:NH1	2.36	0.41
1:A:103:ASP:OD2	1:A:122:ARG:HG2	2.21	0.41
1:A:175:PHE:CD2	1:A:181:LYS:HB3	2.56	0.41
1:A:176:LYS:HE3	1:A:176:LYS:HB2	1.91	0.41
1:A:206:ILE:O	1:A:210:ILE:HG12	2.20	0.41
1:A:34:GLU:O	1:A:38:ILE:HG13	2.20	0.41
1:A:87:GLU:O	1:A:90:ASN:HB3	2.20	0.41
1:A:210:ILE:O	1:A:211:LYS:CB	2.69	0.41
1:A:96:ALA:HB2	1:A:117:TYR:HB2	2.04	0.40
1:A:4:LYS:HB3	1:A:5:LYS:H	1.69	0.40
1:A:202:ASP:HB3	1:A:205:GLU:HG2	2.02	0.40
1:B:672:ILE:HG12	1:B:693:LYS:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/211 (98%)	200 (97%)	5 (2%)	1 (0%)	24	16
1	B	208/211 (99%)	202 (97%)	6 (3%)	0	100	100
All	All	414/422 (98%)	402 (97%)	11 (3%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/176 (100%)	173 (98%)	3 (2%)	53	52
1	B	178/176 (101%)	170 (96%)	8 (4%)	24	17
All	All	354/352 (101%)	343 (97%)	11 (3%)	35	29

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ASN
1	A	150	LYS
1	A	177	LYS
1	B	519	ASN
1	B	533	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	574	ARG
1	B	576	THR
1	B	582	GLU
1	B	607	ASN
1	B	638	VAL
1	B	676	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	590	ASN
1	B	607	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SEP	A	770	-	9,10,10	1.52	1 (11%)	9,14,14	1.59	1 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	B	230	-	4,4,4	1.88	3 (75%)	6,6,6	0.46	0
3	PO4	B	240	-	4,4,4	1.72	1 (25%)	6,6,6	0.44	0
2	SEP	B	270	-	9,10,10	1.59	1 (11%)	9,14,14	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	A	770	-	-	1/10/10/10	-
2	SEP	B	270	-	-	0/10/10/10	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	270	SEP	P-OG	-3.39	1.49	1.60
2	A	770	SEP	P-OG	-3.30	1.49	1.60
3	B	240	PO4	P-O3	-2.15	1.48	1.54
3	B	230	PO4	P-O4	-2.14	1.48	1.54
3	B	230	PO4	P-O3	-2.13	1.48	1.54
3	B	230	PO4	P-O2	-2.11	1.48	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	770	SEP	OG-CB-CA	4.23	111.75	108.06

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	770	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/211 (97%)	0.26	6 (2%) 53 57	18, 27, 41, 56	0
1	B	208/211 (98%)	0.57	10 (4%) 35 38	18, 31, 46, 58	0
All	All	414/422 (98%)	0.42	16 (3%) 43 46	18, 29, 44, 58	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	535	VAL	3.7
1	A	210	ILE	3.1
1	B	527	ARG	3.0
1	A	4	LYS	2.8
1	A	211	LYS	2.7
1	B	642	ASN	2.6
1	B	574	ARG	2.5
1	B	526	ALA	2.3
1	B	530	GLY	2.3
1	B	529	ALA	2.2
1	A	160	GLU	2.2
1	B	536	LYS	2.1
1	A	201	ARG	2.1
1	A	33	GLU	2.0
1	B	641	GLU	2.0
1	B	531	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	B	240	5/5	0.88	0.14	50,50,51,52	0
2	SEP	B	270	11/11	0.93	0.10	24,30,35,37	0
3	PO4	B	230	5/5	0.96	0.09	27,28,31,31	2
2	SEP	A	770	11/11	0.98	0.06	19,22,25,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.