



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:49 PM UTC

PDB ID : 4L79 / pdb_00004l79
Title : Crystal Structure of nucleotide-free Myosin 1b residues 1-728 with bound Calmodulin
Authors : Shuman, H.; Zwolak, A.; Dominguez, R.; Ostap, E.M.
Deposited on : 2013-06-13
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	744	4% 83% 15% ..
2	B	149	12% 75% 20% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-Ib.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	734	Total	C	N	O	S	0	7	0
			6010	3820	1035	1130	25			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	729	GLY	-	expression tag	UNP Q05096
A	730	LEU	-	expression tag	UNP Q05096
A	731	ASN	-	expression tag	UNP Q05096
A	732	ASP	-	expression tag	UNP Q05096
A	733	ILE	-	expression tag	UNP Q05096
A	734	PHE	-	expression tag	UNP Q05096
A	735	GLU	-	expression tag	UNP Q05096
A	736	ALA	-	expression tag	UNP Q05096
A	737	GLN	-	expression tag	UNP Q05096
A	738	LYS	-	expression tag	UNP Q05096
A	739	ILE	-	expression tag	UNP Q05096
A	740	GLU	-	expression tag	UNP Q05096
A	741	TRP	-	expression tag	UNP Q05096
A	742	HIS	-	expression tag	UNP Q05096
A	743	GLU	-	expression tag	UNP Q05096
A	744	ASP	-	expression tag	UNP Q05096

- Molecule 2 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	147	Total	C	N	O	S	0	0	0
			1161	711	187	254	9			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	440	Total 440	O 440	0	0
4	B	19	Total 19	O 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Unconventional myosin-Ib



• Molecule 2: Calmodulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	114.18Å 45.09Å 115.55Å 90.00° 117.18° 90.00°	Depositor
Resolution (Å)	39.21 – 2.30 39.21 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.21-2.30) 99.6 (39.21-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.98 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1314)	Depositor
R, R_{free}	0.189 , 0.249 0.191 , 0.251	Depositor DCC
R_{free} test set	2381 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7631	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/6145	0.82	2/8287 (0.0%)
2	B	0.37	0/1173	0.79	4/1573 (0.3%)
All	All	0.52	0/7318	0.82	6/9860 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	137	VAL	N-CA-C	6.71	117.70	108.84
1	A	495	LYS	N-CA-C	6.13	118.09	108.96
2	B	136	GLN	N-CA-C	5.66	117.63	110.33
2	B	66	PHE	CA-C-N	-5.54	113.31	119.19
2	B	66	PHE	C-N-CA	-5.54	113.31	119.19
1	A	604	ALA	N-CA-C	5.02	117.64	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6010	0	5982	73	1
2	B	1161	0	1088	24	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	440	0	0	15	1
4	B	19	0	0	6	0
All	All	7631	0	7070	93	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:495:LYS:HE2	1:A:496:CYS:SG	2.02	0.98
1:A:462:GLU:OE2	4:A:1119:HOH:O	1.87	0.91
1:A:742:HIS:HB3	1:A:743:GLU:HB2	1.68	0.76
1:A:496:CYS:SG	1:A:498:ARG:HG3	2.26	0.74
1:A:109:GLU:OE2	4:A:1289:HOH:O	2.08	0.71
2:B:102:SER:HA	2:B:136:GLN:HG2	1.75	0.69
1:A:252:ARG:NH1	4:A:1175:HOH:O	2.22	0.68
2:B:3:ASP:OD1	4:B:213:HOH:O	2.13	0.64
1:A:433:GLU:O	4:A:1222:HOH:O	2.15	0.64
2:B:20:PHE:O	4:B:214:HOH:O	2.15	0.64
2:B:55:GLU:OE1	2:B:75:ARG:NH2	2.31	0.63
1:A:712:GLN:HG2	2:B:117:LEU:HD13	1.79	0.63
2:B:103:ALA:H	2:B:136:GLN:HG2	1.64	0.62
1:A:656:TRP:CG	1:A:657:LYS:HA	2.34	0.62
1:A:131:LYS:HG3	1:A:180:PRO:HD3	1.84	0.60
2:B:109:VAL:HG13	2:B:113:LEU:HD12	1.82	0.60
1:A:599:ASN:ND2	1:A:606:ILE:O	2.34	0.59
1:A:59:ARG:HD3	1:A:604:ALA:HB1	1.84	0.59
2:B:103:ALA:H	2:B:136:GLN:CG	2.15	0.58
1:A:492:ARG:O	4:A:1314:HOH:O	2.16	0.58
1:A:656:TRP:HA	1:A:656:TRP:CE3	2.38	0.57
1:A:496:CYS:HG	1:A:498:ARG:N	2.03	0.57
1:A:715:TYR:HA	2:B:146:MET:HE2	1.87	0.57
1:A:20:THR:OG1	1:A:48:ILE:HG12	2.05	0.57
2:B:81:ASP:OD2	4:B:215:HOH:O	2.16	0.57
1:A:639:GLU:HB2	1:A:640:PRO:HD3	1.87	0.56
1:A:145:SER:HB2	1:A:368:ILE:HD13	1.87	0.56
1:A:556:GLU:HA	1:A:561:LYS:HD3	1.87	0.56
1:A:159:ARG:NH1	1:A:393:ILE:O	2.37	0.56
1:A:110:SER:OG	1:A:163:SER:HA	2.06	0.55
1:A:741:TRP:O	1:A:742:HIS:ND1	2.39	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:742:HIS:CB	1:A:743:GLU:HB2	2.36	0.54
1:A:636:GLN:OE1	4:A:1009:HOH:O	2.18	0.54
1:A:109:GLU:HG2	1:A:597:LYS:HE3	1.90	0.54
1:A:561:LYS:NZ	4:A:1165:HOH:O	2.16	0.53
1:A:656:TRP:CB	1:A:657:LYS:HA	2.39	0.53
2:B:64:ILE:HG23	2:B:68:GLU:HG3	1.92	0.52
2:B:83:GLU:HG3	2:B:147:THR:HG21	1.89	0.52
1:A:612:VAL:O	1:A:616:ILE:HG12	2.09	0.52
1:A:734:PHE:O	4:A:1255:HOH:O	2.20	0.51
1:A:47:TYR:CZ	1:A:82:PRO:HA	2.46	0.50
1:A:73:ARG:HD3	1:A:128:VAL:HG23	1.92	0.50
2:B:122:VAL:O	2:B:126:ILE:HG12	2.10	0.50
1:A:571:GLY:HA3	4:A:1114:HOH:O	2.11	0.50
1:A:76:ASN:HB2	1:A:79:GLU:HG3	1.92	0.49
2:B:76:LYS:HB2	4:B:219:HOH:O	2.12	0.49
1:A:466:ARG:NH2	1:A:469:THR:O	2.43	0.49
2:B:73:MET:O	2:B:77:MET:HE2	2.13	0.49
1:A:444:ILE:HD11	1:A:488:HIS:ND1	2.29	0.48
1:A:333:GLN:HA	1:A:333:GLN:HE21	1.80	0.47
1:A:711:ILE:HD11	2:B:89:ALA:HB3	1.97	0.47
1:A:213:SER:O	4:A:1049:HOH:O	2.20	0.47
1:A:231:ARG:NH2	4:A:1024:HOH:O	2.47	0.47
1:A:684:LYS:NZ	4:A:1245:HOH:O	2.04	0.47
1:A:463:GLU:HG3	1:A:475:PHE:HA	1.98	0.46
2:B:76:LYS:CB	4:B:219:HOH:O	2.62	0.46
1:A:73:ARG:HG2	1:A:74:ASN:ND2	2.31	0.46
1:A:539:SER:OG	1:A:556:GLU:OE2	2.26	0.45
1:A:674:PRO:HG2	1:A:677:GLU:HG3	1.98	0.45
2:B:132:ASP:OD1	2:B:132:ASP:N	2.43	0.45
1:A:58:TYR:HB2	1:A:604:ALA:HA	1.99	0.45
2:B:8:GLU:OE1	2:B:8:GLU:N	2.48	0.45
2:B:87:ARG:HG2	2:B:91:ARG:CZ	2.47	0.44
2:B:133:GLY:HA2	2:B:134:ASP:HA	1.46	0.44
1:A:110:SER:HG	1:A:163:SER:HA	1.82	0.44
1:A:310:CYS:HB3	1:A:315:ILE:O	2.18	0.44
1:A:490:GLU:HB2	1:A:511:ARG:HB3	2.00	0.44
1:A:738:LYS:HG3	4:A:1255:HOH:O	2.18	0.44
1:A:406:TYR:OH	1:A:450:GLU:OE2	2.33	0.44
2:B:112:ASN:ND2	4:B:216:HOH:O	2.36	0.43
1:A:651:GLN:HE22	1:A:670:GLU:HG3	1.84	0.43
1:A:203:ARG:HB3	1:A:209:TYR:CE2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:477:GLU:O	1:A:481[A]:GLN:HG2	2.19	0.42
1:A:653:TRP:HA	1:A:654:PRO:HA	1.55	0.42
1:A:122:MET:HE1	1:A:171:ASP:HA	2.02	0.42
1:A:566:ARG:HE	1:A:566:ARG:HB3	1.64	0.42
1:A:689:ASN:HA	1:A:690:PRO:HD3	1.96	0.42
1:A:724:PHE:CG	2:B:39:SER:HB3	2.54	0.41
1:A:226:GLU:HB2	1:A:231:ARG:HD2	2.02	0.41
1:A:173:GLU:HG2	1:A:382:VAL:HG12	2.03	0.41
1:A:509:CYS:SG	1:A:520:LEU:HD11	2.60	0.41
1:A:645:TYR:OH	1:A:697:GLU:OE2	2.35	0.41
1:A:104:ILE:HD12	1:A:121:VAL:HG11	2.02	0.41
1:A:620:GLY:HA2	4:A:952:HOH:O	2.20	0.41
1:A:648:LEU:HD12	1:A:648:LEU:HA	1.83	0.41
1:A:626:ARG:NH2	4:A:1095:HOH:O	2.53	0.41
1:A:125:VAL:HG21	1:A:383:MET:HG2	2.02	0.41
1:A:454:ASN:HA	1:A:459:MET:HE3	2.02	0.41
1:A:650:LYS:HD3	1:A:650:LYS:HA	1.74	0.40
1:A:37:LYS:HE2	1:A:37:LYS:HB3	1.76	0.40
1:A:125:VAL:HA	1:A:128:VAL:HG12	2.03	0.40
2:B:91:ARG:HG2	2:B:139:TYR:OH	2.22	0.40
1:A:374:ALA:O	1:A:376:THR:HG22	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:GLU:OE1	1:A:495:LYS:NZ[2_746]	2.03	0.17
4:A:914:HOH:O	4:A:1018:HOH:O[2_656]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	735/744 (99%)	712 (97%)	23 (3%)	0	100	100
2	B	145/149 (97%)	144 (99%)	1 (1%)	0	100	100
All	All	880/893 (98%)	856 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	663/664 (100%)	643 (97%)	20 (3%)	36	53
2	B	126/127 (99%)	119 (94%)	7 (6%)	19	28
All	All	789/791 (100%)	762 (97%)	27 (3%)	32	49

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	LYS
1	A	289	GLU
1	A	329	VAL
1	A	377	LYS
1	A	389	TYR
1	A	444	ILE
1	A	486	HIS
1	A	494	SER
1	A	495	LYS
1	A	496	CYS
1	A	500	LEU
1	A	597	LYS
1	A	648	LEU
1	A	655	HIS
1	A	656	TRP
1	A	703	ARG
1	A	714	ILE
1	A	725	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	739	ILE
1	A	744	ASP
2	B	55	GLU
2	B	56	VAL
2	B	117	LEU
2	B	131	ILE
2	B	136	GLN
2	B	137	VAL
2	B	146	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN
1	A	102	GLN
1	A	333	GLN
1	A	424	GLN
1	A	522	GLN
1	A	586	GLN
1	A	589	ASN
1	A	731	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	734/744 (98%)	0.07	29 (3%) 42 44	14, 38, 72, 106	7 (0%)
2	B	147/149 (98%)	0.93	18 (12%) 8 9	40, 73, 98, 109	0
All	All	881/893 (98%)	0.22	47 (5%) 32 34	14, 42, 84, 109	7 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	ALA	4.3
1	A	656	TRP	3.9
2	B	131	ILE	3.6
1	A	496	CYS	3.4
1	A	6	VAL	3.1
1	A	671	LEU	3.0
1	A	604	ALA	3.0
1	A	733	ILE	3.0
1	A	739	ILE	3.0
2	B	56	VAL	2.8
2	B	135	GLY	2.8
1	A	376	THR	2.7
1	A	649	CYS	2.7
2	B	137	VAL	2.7
1	A	742	HIS	2.6
1	A	653	TRP	2.6
1	A	668	PHE	2.6
2	B	117	LEU	2.5
1	A	736	ALA	2.5
2	B	136	GLN	2.5
2	B	58	ALA	2.5
2	B	133	GLY	2.5
1	A	744	ASP	2.4
2	B	60	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	680	PHE	2.4
1	A	231	ARG	2.4
2	B	108	HIS	2.4
2	B	103	ALA	2.3
1	A	732	ASP	2.3
2	B	129	ALA	2.3
1	A	667	LEU	2.3
2	B	102	SER	2.3
1	A	335	LYS	2.2
1	A	660	ALA	2.2
2	B	134	ASP	2.2
1	A	659	PRO	2.2
2	B	97	GLY	2.2
1	A	375	GLN	2.2
1	A	704	LEU	2.2
1	A	741	TRP	2.1
1	A	289	GLU	2.1
1	A	657	LYS	2.1
1	A	605	HIS	2.1
2	B	49	LEU	2.1
1	A	498	ARG	2.1
2	B	80	THR	2.0
2	B	126	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	801	1/1	0.98	0.05	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.