



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:07 PM UTC

PDB ID : 4L7U / pdb_0000417u
Title : Human artd3 (parp3) - catalytic domain in complex with inhibitor ME0398
Authors : Karlberg, T.; Thorsell, A.G.; Lindgren, A.E.G.; Ekblad, T.; Spjut, S.; Andersson, C.D.; Weigelt, J.; Linusson, A.; Elofsson, M.; Schuler, H.
Deposited on : 2013-06-14
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	10% 80% 17% ..

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2744	1732	475	523	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	176	SER	-	expression tag	UNP Q9Y6F1
A	177	MET	-	expression tag	UNP Q9Y6F1

- Molecule 2 is methyl N-[3-(4-oxo-3,4-dihydroquinazolin-2-yl)propanoyl]-L-phenylalaninate (CCD ID: 1VC) (formula: C₂₁H₂₁N₃O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	21	3	4		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.19Å 56.75Å 56.91Å 90.00° 112.10° 90.00°	Depositor
Resolution (Å)	27.40 – 2.80 27.40 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (27.40-2.80) 99.5 (27.40-2.80)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.20	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.80Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.240 , 0.288 0.248 , 0.326	Depositor DCC
R_{free} test set	489 reflections (6.01%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	1.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.043 for l,-k,h	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2772	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1VC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/2805	1.37	14/3797 (0.4%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	318	ASP	CA-CB-CG	6.80	119.40	112.60
1	A	202	PHE	CA-CB-CG	6.46	120.27	113.80
1	A	478	ASP	CA-CB-CG	6.14	118.74	112.60
1	A	264	ASN	CA-CB-CG	5.86	118.46	112.60
1	A	387	ASN	CA-CB-CG	5.51	118.11	112.60
1	A	410	GLY	N-CA-C	5.48	120.94	112.45
1	A	265	PHE	CA-CB-CG	5.29	119.09	113.80
1	A	254	SER	CA-C-N	5.26	127.58	120.38
1	A	254	SER	C-N-CA	5.26	127.58	120.38
1	A	204	ASN	CA-CB-CG	5.25	117.85	112.60
1	A	292	ILE	CA-C-N	5.16	127.62	120.29
1	A	292	ILE	C-N-CA	5.16	127.62	120.29
1	A	390	VAL	CA-C-N	5.09	127.16	120.60
1	A	390	VAL	C-N-CA	5.09	127.16	120.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2744	0	2706	26	0
2	A	28	0	20	5	0
All	All	2772	0	2726	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:ARG:HB3	1:A:310:GLU:HG2	1.68	0.74
1:A:462:PRO:HB3	1:A:469:VAL:HG23	1.77	0.67
1:A:346:THR:HB	1:A:418:GLU:HA	1.85	0.59
1:A:322:LEU:HD22	1:A:395:LEU:HB2	1.83	0.58
1:A:387:ASN:H	2:A:601:1VC:H11	1.71	0.55
1:A:519:GLN:HG3	1:A:522:GLN:HG3	1.89	0.55
1:A:291:ASP:HB2	1:A:390:VAL:HG12	1.90	0.52
1:A:390:VAL:CG2	2:A:601:1VC:H9	2.41	0.50
1:A:514:GLU:HG2	2:A:601:1VC:H19	1.93	0.49
1:A:390:VAL:HG21	2:A:601:1VC:H9	1.94	0.48
1:A:202:PHE:HB2	1:A:218:LEU:HD11	1.94	0.47
1:A:477:PRO:HB3	1:A:497:GLY:O	2.14	0.47
1:A:485:LEU:HD12	1:A:494:VAL:HG21	1.96	0.47
1:A:394:ILE:HG22	1:A:525:LEU:HD11	1.97	0.46
1:A:355:LEU:HD21	1:A:358:ILE:HD11	1.96	0.46
1:A:409:VAL:HG22	1:A:516:LEU:HG	1.97	0.45
1:A:386:THR:HG22	2:A:601:1VC:OAB	2.17	0.45
1:A:178:LYS:HG2	1:A:311:VAL:HG13	1.99	0.44
1:A:391:VAL:O	1:A:395:LEU:HG	2.17	0.44
1:A:347:GLY:HA2	1:A:419:ASN:HB3	1.99	0.43
1:A:409:VAL:HG22	1:A:409:VAL:O	2.18	0.43
1:A:189:THR:O	1:A:193:ILE:HG12	2.19	0.43
1:A:312:PRO:HG2	1:A:317:ARG:HB2	2.01	0.42
1:A:212:ASP:HA	1:A:264:ASN:HD22	1.84	0.42
1:A:487:LEU:HD22	1:A:526:ARG:HD2	2.03	0.41
1:A:342:TYR:CZ	1:A:495:PRO:HD2	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	348/357 (98%)	334 (96%)	13 (4%)	1 (0%)	36	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	388	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/310 (97%)	281 (93%)	21 (7%)	14	40

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	LYS
1	A	226	ILE
1	A	248	GLN
1	A	279	LEU
1	A	303	GLU
1	A	316	ASP
1	A	327	GLN
1	A	335	GLU
1	A	376	LEU
1	A	409	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	431	CYS
1	A	435	HIS
1	A	436	VAL
1	A	439	MET
1	A	444	VAL
1	A	449	GLU
1	A	452	ILE
1	A	454	THR
1	A	467	ASP
1	A	485	LEU
1	A	521	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	GLN
1	A	264	ASN
1	A	296	GLN
1	A	320	GLN
1	A	327	GLN
1	A	387	ASN
1	A	522	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1VC	A	601	-	28,30,30	3.31	9 (32%)	37,40,40	3.49	17 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1VC	A	601	-	-	9/19/19/19	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	1VC	CAO-CAX	-12.11	1.28	1.49
2	A	601	1VC	CB-CG	-8.96	1.30	1.51
2	A	601	1VC	CAN-CAU	-4.47	1.42	1.51
2	A	601	1VC	CAZ-NAQ	-4.14	1.31	1.39
2	A	601	1VC	CBA-CAY	-2.84	1.42	1.47
2	A	601	1VC	O-C	2.62	1.27	1.21
2	A	601	1VC	CA-C	2.36	1.58	1.52
2	A	601	1VC	CAM-CBA	-2.27	1.36	1.39
2	A	601	1VC	CBA-CAZ	2.12	1.43	1.40

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	1VC	CB-CA-C	10.97	136.94	110.42
2	A	601	1VC	CAY-NAS-CAX	-10.75	117.69	123.90
2	A	601	1VC	CAN-CAO-CAX	-7.10	105.23	112.55
2	A	601	1VC	OAB-CAU-N	5.00	131.43	122.95
2	A	601	1VC	CAO-CAN-CAU	-4.99	102.39	112.67
2	A	601	1VC	CAA-OAT-C	-3.78	107.34	115.92
2	A	601	1VC	OAB-CAU-CAN	-3.60	115.50	122.02
2	A	601	1VC	OAT-C-CA	3.50	120.39	111.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	1VC	CB-CG-CD1	-2.80	115.68	120.90
2	A	601	1VC	CA-N-CAU	2.74	128.56	121.68
2	A	601	1VC	NAS-CAX-NAQ	2.74	126.54	123.08
2	A	601	1VC	CAL-CAZ-CBA	2.35	121.45	119.14
2	A	601	1VC	CD2-CG-CD1	2.27	121.60	118.23
2	A	601	1VC	CAL-CAZ-NAQ	-2.26	115.73	118.61
2	A	601	1VC	C-CA-N	-2.10	105.82	110.65
2	A	601	1VC	O-C-CA	-2.08	117.78	123.94
2	A	601	1VC	CG-CB-CA	-2.03	107.96	113.36

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	1VC	CA-C-OAT-CAA
2	A	601	1VC	O-C-OAT-CAA
2	A	601	1VC	N-CA-CB-CG
2	A	601	1VC	O-C-CA-N
2	A	601	1VC	OAT-C-CA-N
2	A	601	1VC	OAT-C-CA-CB
2	A	601	1VC	CAO-CAN-CAU-OAB
2	A	601	1VC	O-C-CA-CB
2	A	601	1VC	CAO-CAN-CAU-N

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	1VC	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/357 (98%)	0.96	37 (10%) 11 8	13, 38, 74, 107	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	332	GLY	4.0
1	A	365	GLY	3.6
1	A	366	GLU	3.5
1	A	245	ASP	3.4
1	A	274	ASN	3.3
1	A	505	PHE	3.2
1	A	500	VAL	3.1
1	A	375	LYS	3.0
1	A	427	ILE	2.9
1	A	353	PRO	2.9
1	A	460	LYS	2.9
1	A	434	HIS	2.8
1	A	461	SER	2.8
1	A	372	ALA	2.8
1	A	188	ALA	2.6
1	A	437	GLY	2.6
1	A	480	THR	2.6
1	A	234	GLU	2.5
1	A	241	LYS	2.5
1	A	270	PRO	2.5
1	A	497	GLY	2.4
1	A	359	TRP	2.4
1	A	345	GLN	2.4
1	A	350	HIS	2.3
1	A	246	GLY	2.3
1	A	373	HIS	2.2
1	A	532	HIS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	254	SER	2.2
1	A	294	LEU	2.2
1	A	273	ILE	2.1
1	A	489	GLY	2.1
1	A	339	ILE	2.1
1	A	499	PRO	2.1
1	A	247	GLY	2.1
1	A	436	VAL	2.1
1	A	364	GLU	2.0
1	A	336	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	1VC	A	601	28/28	0.84	0.15	28,33,43,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 1VC A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.