



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 05:24 AM UTC

PDB ID : 5L7B / pdb_00005l7b
BMRB ID : 34004
Title : Solution structure of the human SNF5/INI1 domain
Authors : Allen, M.D.; Zinzalla, G.; Bycroft, M.
Deposited on : 2016-06-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 87%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	78	46% 33% • 15% •

2 Ensemble composition and analysis

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:185-A:248 (64)	0.26	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 5, 6, 9, 10, 15, 16, 17, 18, 19
2	2, 3, 11, 12
3	4, 7, 13
4	14, 20
Single-model clusters	8

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1205 atoms, of which 597 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily B member 1.

Mol	Chain	Residues	Atoms						Trace
1	A	76	Total	C	H	N	O	S	0
			1205	383	597	95	125	5	

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	181	GLY	-	expression tag	UNP Q12824
A	182	GLY	-	expression tag	UNP Q12824
A	183	SER	-	expression tag	UNP Q12824

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1



4.2.2 Score per residue for model 2

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1






4.2.3 Score per residue for model 3

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1


Chain A: 44% 35% 15%



4.2.4 Score per residue for model 4

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1

Chain A: 41% 33% 8% 15%



4.2.5 Score per residue for model 5


- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1

Chain A: 46% 29% 6% 15%




4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2.7 Score per residue for model 7

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1



4.2.8 Score per residue for model 8

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1



4.2.9 Score per residue for model 9

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1





4.2.10 Score per residue for model 10

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1


Chain A: 42% 36% 15%



4.2.11 Score per residue for model 11

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1

Chain A: 37% 40% 5% 15%



4.2.12 Score per residue for model 12

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1


Chain A: 47% 27% 8% 15%



4.2.13 Score per residue for model 13

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1

Chain A: 41% 38% 15%




4.2.14 Score per residue for model 14

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2.15 Score per residue for model 15

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2.16 Score per residue for model 16

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2.17 Score per residue for model 17

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2.18 Score per residue for model 18

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1




4.2.19 Score per residue for model 19

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1



4.2.20 Score per residue for model 20

- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family B member 1





5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
CNS	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	918
Number of shifts mapped to atoms	918
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	87%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	516	516	516	26±4
All	All	10320	10320	10320	530

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:191:LEU:HD11	1:A:204:PHE:CE2	0.83	2.09	18	1
1:A:189:ILE:HG21	1:A:204:PHE:CE2	0.82	2.09	7	8
1:A:200:LEU:HD23	1:A:228:LEU:HD11	0.81	1.53	2	5
1:A:208:MET:HE3	1:A:245:ILE:HG22	0.80	1.52	5	9
1:A:193:MET:HG3	1:A:200:LEU:HD12	0.79	1.53	8	4
1:A:189:ILE:HG21	1:A:204:PHE:CE1	0.78	2.13	15	12
1:A:193:MET:CG	1:A:200:LEU:HD12	0.77	2.09	8	1
1:A:195:ILE:HD13	1:A:233:PHE:CE1	0.77	2.15	7	1
1:A:228:LEU:HD22	1:A:233:PHE:CZ	0.76	2.15	16	13
1:A:242:ARG:HA	1:A:245:ILE:HD12	0.75	1.57	16	5
1:A:195:ILE:HG21	1:A:232:THR:OG1	0.73	1.83	3	14
1:A:191:LEU:HD22	1:A:193:MET:HE1	0.72	1.59	14	1
1:A:241:ILE:HG22	1:A:245:ILE:HD11	0.72	1.62	14	18
1:A:228:LEU:HD22	1:A:233:PHE:CE1	0.71	2.20	15	8

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:200:LEU:HD23	1:A:228:LEU:CD1	0.70	2.15	20	3
1:A:193:MET:HG2	1:A:200:LEU:HD12	0.70	1.62	20	1
1:A:194:GLU:C	1:A:195:ILE:HD12	0.70	2.12	7	1
1:A:230:PRO:HA	1:A:234:VAL:HG23	0.69	1.64	4	20
1:A:200:LEU:C	1:A:200:LEU:HD13	0.68	2.14	15	5
1:A:228:LEU:HD22	1:A:233:PHE:CE2	0.66	2.26	17	5
1:A:200:LEU:HD12	1:A:201:ARG:N	0.65	2.06	18	15
1:A:195:ILE:HD13	1:A:233:PHE:CZ	0.65	2.26	7	1
1:A:230:PRO:CA	1:A:234:VAL:HG23	0.64	2.23	2	20
1:A:213:MET:HE3	1:A:218:PHE:CD2	0.63	2.28	16	3
1:A:208:MET:HE3	1:A:245:ILE:CG2	0.61	2.24	15	8
1:A:200:LEU:HD13	1:A:200:LEU:O	0.61	1.96	8	5
1:A:194:GLU:HA	1:A:198:GLN:O	0.60	1.97	9	19
1:A:195:ILE:HD12	1:A:233:PHE:CD2	0.60	2.31	12	1
1:A:202:ASP:CG	1:A:222:LEU:HD21	0.59	2.22	20	1
1:A:208:MET:HG2	1:A:245:ILE:HG23	0.59	1.73	20	11
1:A:200:LEU:HD22	1:A:201:ARG:N	0.59	2.11	8	2
1:A:193:MET:HE1	1:A:237:ILE:HA	0.58	1.74	6	3
1:A:208:MET:CG	1:A:245:ILE:HG23	0.58	2.29	2	2
1:A:191:LEU:CD1	1:A:204:PHE:CE2	0.56	2.87	18	2
1:A:241:ILE:O	1:A:245:ILE:HG13	0.56	2.01	18	18
1:A:195:ILE:CG1	1:A:233:PHE:CZ	0.56	2.88	2	18
1:A:195:ILE:HD12	1:A:233:PHE:CG	0.55	2.36	12	1
1:A:187:VAL:HG13	1:A:248:TYR:CD1	0.54	2.38	7	5
1:A:191:LEU:HD13	1:A:202:ASP:CB	0.54	2.32	18	1
1:A:241:ILE:HG22	1:A:245:ILE:CD1	0.54	2.31	14	13
1:A:187:VAL:HG13	1:A:248:TYR:CG	0.54	2.38	16	9
1:A:193:MET:HE1	1:A:240:ALA:CB	0.54	2.32	17	2
1:A:200:LEU:HD12	1:A:200:LEU:C	0.53	2.28	5	13
1:A:191:LEU:HD21	1:A:204:PHE:CE2	0.53	2.39	18	1
1:A:206:TRP:CD1	1:A:245:ILE:HD13	0.52	2.39	3	5
1:A:202:ASP:OD2	1:A:226:LEU:HD11	0.52	2.05	8	1
1:A:222:LEU:HD23	1:A:226:LEU:HD12	0.52	1.81	8	1
1:A:189:ILE:CG2	1:A:204:PHE:CE1	0.52	2.92	15	11
1:A:189:ILE:HG21	1:A:204:PHE:CZ	0.52	2.39	18	2
1:A:191:LEU:HD11	1:A:204:PHE:CD2	0.52	2.40	18	1
1:A:228:LEU:HB3	1:A:233:PHE:CD2	0.51	2.41	7	9
1:A:228:LEU:HB3	1:A:233:PHE:CD1	0.51	2.40	15	11
1:A:187:VAL:CG2	1:A:208:MET:CG	0.50	2.89	7	10
1:A:200:LEU:HD22	1:A:228:LEU:HD13	0.50	1.82	3	5
1:A:189:ILE:CG2	1:A:204:PHE:CE2	0.50	2.93	14	6

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:191:LEU:HD21	1:A:204:PHE:CZ	0.49	2.43	18	1
1:A:189:ILE:HG21	1:A:204:PHE:HE2	0.49	1.68	3	4
1:A:200:LEU:C	1:A:200:LEU:CD1	0.48	2.86	15	3
1:A:241:ILE:O	1:A:245:ILE:HG12	0.48	2.08	5	2
1:A:185:VAL:HG11	1:A:208:MET:SD	0.48	2.48	7	3
1:A:200:LEU:HD22	1:A:233:PHE:CD1	0.48	2.43	14	6
1:A:208:MET:HG2	1:A:245:ILE:CG2	0.47	2.39	2	3
1:A:234:VAL:HB	1:A:235:PRO:HD3	0.47	1.87	19	2
1:A:191:LEU:CD1	1:A:204:PHE:CE1	0.47	2.98	3	3
1:A:195:ILE:HG12	1:A:233:PHE:CZ	0.47	2.45	20	7
1:A:198:GLN:CD	1:A:228:LEU:HD23	0.47	2.35	18	3
1:A:198:GLN:NE2	1:A:228:LEU:HD23	0.47	2.25	7	3
1:A:202:ASP:OD1	1:A:226:LEU:HD11	0.47	2.10	20	1
1:A:198:GLN:OE1	1:A:228:LEU:HD23	0.47	2.09	18	2
1:A:206:TRP:CD1	1:A:245:ILE:CD1	0.46	2.98	18	5
1:A:200:LEU:HD22	1:A:233:PHE:CD2	0.46	2.45	13	5
1:A:193:MET:HE1	1:A:240:ALA:HB2	0.46	1.86	17	3
1:A:189:ILE:HG21	1:A:204:PHE:HE1	0.46	1.68	1	3
1:A:200:LEU:CD2	1:A:233:PHE:CD1	0.45	2.99	4	3
1:A:200:LEU:CD2	1:A:233:PHE:CD2	0.45	3.00	16	4
1:A:200:LEU:CD2	1:A:233:PHE:CE2	0.44	3.01	1	3
1:A:210:GLU:N	1:A:210:GLU:OE1	0.44	2.51	5	2
1:A:233:PHE:O	1:A:237:ILE:HG13	0.44	2.13	19	12
1:A:195:ILE:HG21	1:A:232:THR:CB	0.44	2.42	15	1
1:A:190:ARG:C	1:A:191:LEU:HD23	0.44	2.38	12	1
1:A:195:ILE:CG1	1:A:233:PHE:CE2	0.44	3.01	8	2
1:A:230:PRO:O	1:A:234:VAL:HG23	0.44	2.13	18	11
1:A:191:LEU:HD12	1:A:204:PHE:CE1	0.43	2.48	3	2
1:A:187:VAL:HG13	1:A:248:TYR:CD2	0.43	2.49	3	1
1:A:234:VAL:HB	1:A:235:PRO:CD	0.43	2.43	10	18
1:A:198:GLN:NE2	1:A:228:LEU:CD2	0.43	2.81	8	4
1:A:210:GLU:N	1:A:210:GLU:OE2	0.43	2.52	9	4
1:A:217:MET:HG3	1:A:218:PHE:N	0.43	2.28	2	1
1:A:191:LEU:HD13	1:A:237:ILE:HG23	0.43	1.91	15	1
1:A:191:LEU:HD13	1:A:202:ASP:HB3	0.42	1.92	18	1
1:A:230:PRO:O	1:A:234:VAL:HB	0.42	2.14	8	3
1:A:232:THR:C	1:A:235:PRO:HD2	0.41	2.40	4	1
1:A:195:ILE:HG12	1:A:233:PHE:CE2	0.41	2.50	8	2
1:A:234:VAL:N	1:A:235:PRO:HD2	0.41	2.31	20	1
1:A:187:VAL:CG1	1:A:248:TYR:CG	0.41	3.04	12	1
1:A:195:ILE:HD12	1:A:195:ILE:N	0.41	2.30	7	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:191:LEU:CD1	1:A:237:ILE:HG23	0.41	2.44	15	1
1:A:202:ASP:OD2	1:A:222:LEU:HD21	0.41	2.15	20	1
1:A:220:GLU:OE1	1:A:234:VAL:HG21	0.41	2.15	2	1
1:A:248:TYR:CD2	1:A:248:TYR:C	0.41	2.99	10	1
1:A:219:SER:OG	1:A:238:ALA:HB2	0.41	2.16	9	1
1:A:200:LEU:CD2	1:A:233:PHE:CE1	0.41	3.04	4	1
1:A:191:LEU:HD12	1:A:204:PHE:CE2	0.41	2.51	13	2
1:A:189:ILE:CB	1:A:204:PHE:CE1	0.41	3.04	18	1
1:A:239:SER:HA	1:A:242:ARG:HG2	0.41	1.92	3	2
1:A:241:ILE:CG2	1:A:245:ILE:HD11	0.41	2.42	14	1
1:A:200:LEU:C	1:A:200:LEU:HD22	0.40	2.41	8	1
1:A:187:VAL:CG2	1:A:208:MET:HG2	0.40	2.46	9	2
1:A:222:LEU:HD23	1:A:226:LEU:CD1	0.40	2.46	9	1
1:A:189:ILE:HG22	1:A:191:LEU:CD2	0.40	2.47	8	1
1:A:228:LEU:HD13	1:A:233:PHE:CD2	0.40	2.51	19	1
1:A:195:ILE:N	1:A:195:ILE:CD1	0.40	2.85	7	1
1:A:191:LEU:HB3	1:A:193:MET:HE2	0.40	1.93	12	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	64/78 (82%)	61±1 (95±1%)	3±1 (5±1%)	0±0 (0±0%)	100	100
All	All	1280/1560 (82%)	1213 (95%)	67 (5%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/71 (83%)	46±2 (78±3%)	13±2 (22±3%)	2	29
All	All	1180/1420 (83%)	918 (78%)	262 (22%)	2	29

All 29 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	222	LEU	20
1	A	193	MET	19
1	A	200	LEU	19
1	A	202	ASP	18
1	A	205	THR	14
1	A	210	GLU	11
1	A	243	GLN	11
1	A	199	LYS	10
1	A	216	GLU	10
1	A	187	VAL	10
1	A	207	ASN	10
1	A	209	ASN	9
1	A	246	GLU	9
1	A	194	GLU	8
1	A	227	ASP	8
1	A	201	ARG	8
1	A	213	MET	8
1	A	217	MET	7
1	A	190	ARG	7
1	A	211	LYS	7
1	A	242	ARG	7
1	A	224	ASP	5
1	A	208	MET	5
1	A	196	ASP	5
1	A	198	GLN	5
1	A	225	ASP	4
1	A	247	SER	4
1	A	219	SER	3
1	A	191	LEU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 87% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *snf5.bmr.b.data*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	918
Number of shifts mapped to atoms	918
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	76	-0.23 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	75	0.25 ± 0.18	None needed (< 0.5 ppm)
$^{13}\text{C}'$	76	0.74 ± 0.17	Should be applied
^{15}N	71	-0.12 ± 0.26	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 788 atoms were assigned a chemical shift out of a possible 903. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	313/313 (100%)	125/125 (100%)	128/128 (100%)	60/60 (100%)
Sidechain	450/539 (83%)	340/352 (97%)	105/170 (62%)	5/17 (29%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	25/51 (49%)	25/25 (100%)	0/25 (0%)	0/1 (0%)
Overall	788/903 (87%)	490/502 (98%)	233/323 (72%)	65/78 (83%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 914 atoms were assigned a chemical shift out of a possible 1043. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	370/371 (100%)	147/148 (99%)	152/152 (100%)	71/71 (100%)
Sidechain	519/621 (84%)	392/404 (97%)	121/199 (61%)	6/18 (33%)
Aromatic	25/51 (49%)	25/25 (100%)	0/25 (0%)	0/1 (0%)
Overall	914/1043 (88%)	564/577 (98%)	273/376 (73%)	77/90 (86%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	214	THR	HG1	6.04	0.08 – 2.19	23.2
1	A	232	THR	HG1	5.24	0.08 – 2.19	19.4

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

