



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:11 PM UTC

PDB ID : 5L7D / pdb_0000517d
Title : Structure of human Smoothed in complex with cholesterol
Authors : Byrne, E.F.X.; Sircar, R.; Miller, P.S.; Hedger, G.; Luchetti, G.; Nachtergaele, S.; Tully, M.D.; Mydock-McGrane, L.; Covey, D.F.; Rambo, R.P.; Sansom, M.S.P.; Newstead, S.; Rohatgi, R.; Siebold, C.
Deposited on : 2016-06-03
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	638	 82% 10% 8%
1	B	638	 82% 9% 9%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Smoothened homolog,Soluble cytochrome b562,Smoothened homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	588	Total	C	N	O	S	0	0	0
			4640	2985	792	830	33			
1	B	578	Total	C	N	O	S	0	0	0
			4567	2936	779	819	33			

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	329	PHE	VAL	conflict	UNP Q99835
A	1011	ALA	-	linker	UNP Q99835
A	1012	ARG	-	linker	UNP Q99835
A	1013	ARG	-	linker	UNP Q99835
A	1014	GLN	-	linker	UNP Q99835
A	1015	LEU	-	linker	UNP Q99835
A	1022	TRP	MET	conflict	UNP P0ABE7
A	1117	ILE	HIS	conflict	UNP P0ABE7
A	1121	LEU	-	linker	UNP P0ABE7
A	1122	GLU	-	linker	UNP P0ABE7
A	1123	ARG	-	linker	UNP P0ABE7
A	1124	ALA	-	linker	UNP P0ABE7
A	1125	ARG	-	linker	UNP P0ABE7
A	1126	SER	-	linker	UNP P0ABE7
A	1127	THR	-	linker	UNP P0ABE7
A	1128	LEU	-	linker	UNP P0ABE7
A	556	GLY	-	expression tag	UNP Q99835
A	557	THR	-	expression tag	UNP Q99835
A	558	GLU	-	expression tag	UNP Q99835
A	559	THR	-	expression tag	UNP Q99835
A	560	SER	-	expression tag	UNP Q99835
A	561	GLN	-	expression tag	UNP Q99835
A	562	VAL	-	expression tag	UNP Q99835
A	563	ALA	-	expression tag	UNP Q99835

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	564	PRO	-	expression tag	UNP Q99835
A	565	ALA	-	expression tag	UNP Q99835
B	329	PHE	VAL	conflict	UNP Q99835
B	1011	ALA	-	linker	UNP Q99835
B	1012	ARG	-	linker	UNP Q99835
B	1013	ARG	-	linker	UNP Q99835
B	1014	GLN	-	linker	UNP Q99835
B	1015	LEU	-	linker	UNP Q99835
B	1022	TRP	MET	conflict	UNP P0ABE7
B	1117	ILE	HIS	conflict	UNP P0ABE7
B	1121	LEU	-	linker	UNP P0ABE7
B	1122	GLU	-	linker	UNP P0ABE7
B	1123	ARG	-	linker	UNP P0ABE7
B	1124	ALA	-	linker	UNP P0ABE7
B	1125	ARG	-	linker	UNP P0ABE7
B	1126	SER	-	linker	UNP P0ABE7
B	1127	THR	-	linker	UNP P0ABE7
B	1128	LEU	-	linker	UNP P0ABE7
B	556	GLY	-	expression tag	UNP Q99835
B	557	THR	-	expression tag	UNP Q99835
B	558	GLU	-	expression tag	UNP Q99835
B	559	THR	-	expression tag	UNP Q99835
B	560	SER	-	expression tag	UNP Q99835
B	561	GLN	-	expression tag	UNP Q99835
B	562	VAL	-	expression tag	UNP Q99835
B	563	ALA	-	expression tag	UNP Q99835
B	564	PRO	-	expression tag	UNP Q99835
B	565	ALA	-	expression tag	UNP Q99835

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CHOLESTEROL (CCD ID: CLR) (formula: $C_{27}H_{46}O$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			28	27	1		


- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Na	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Smoothened homolog,Soluble cytochrome b562,Smoothened homolog



- Molecule 1: Smoothened homolog,Soluble cytochrome b562,Smoothened homolog



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.90Å 63.02Å 208.59Å 90.00° 96.64° 90.00°	Depositor
Resolution (Å)	29.33 – 3.20 29.33 – 3.20	Depositor EDS
% Data completeness (in resolution range)	94.6 (29.33-3.20) 94.4 (29.33-3.20)	Depositor EDS
R_{merge}	0.30	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 3.19Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.230 , 0.264 0.269 , 0.307	Depositor DCC
R_{free} test set	1306 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	93.0	Xtriage
Anisotropy	0.310	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 94.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9265	wwPDB-VP
Average B, all atoms (Å ²)	137.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NAG, CLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/4757	1.34	10/6467 (0.2%)
1	B	0.76	0/4683	1.34	7/6364 (0.1%)
All	All	0.76	0/9440	1.34	17/12831 (0.1%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	ASP	CA-C-N	5.84	128.42	120.54
1	A	255	ASP	C-N-CA	5.84	128.42	120.54
1	A	256	TRP	N-CA-C	5.61	117.84	111.11
1	A	113	ARG	CA-C-N	5.58	129.49	121.50
1	A	113	ARG	C-N-CA	5.58	129.49	121.50
1	A	187	PHE	CA-C-N	5.30	127.92	120.28
1	A	187	PHE	C-N-CA	5.30	127.92	120.28
1	A	1054	ASP	CA-CB-CG	5.26	117.86	112.60
1	B	187	PHE	CA-C-N	5.22	127.79	120.28
1	B	187	PHE	C-N-CA	5.22	127.79	120.28
1	B	446	ASN	CA-CB-CG	5.15	117.75	112.60
1	B	1040	GLN	CA-C-N	5.09	126.98	120.56
1	B	1040	GLN	C-N-CA	5.09	126.98	120.56
1	A	542	LEU	CA-C-N	5.04	127.04	120.28
1	A	542	LEU	C-N-CA	5.04	127.04	120.28
1	B	255	ASP	CA-C-N	5.02	127.01	120.28
1	B	255	ASP	C-N-CA	5.02	127.01	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4640	0	4584	26	0
1	B	4567	0	4494	26	0
2	A	28	0	26	0	0
3	A	28	0	46	1	0
4	A	2	0	0	0	0
All	All	9265	0	9150	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:382:ASP:HB2	1:B:394:TYR:HB2	1.79	0.65
1:B:78:CYS:HB2	1:B:83:LEU:HD21	1.77	0.64
1:B:258:ASN:HA	1:B:261:ARG:HH21	1.63	0.63
1:A:258:ASN:HA	1:A:261:ARG:HH21	1.63	0.63
1:A:382:ASP:HB2	1:A:394:TYR:HB2	1.83	0.61
1:A:339:TRP:HH2	1:A:423:VAL:HG13	1.66	0.60
1:B:252:PHE:HD1	1:B:259:SER:HB3	1.71	0.55
1:A:182:VAL:HA	1:A:185:ILE:HD12	1.88	0.54
1:B:397:TYR:HB3	1:B:477:GLN:HG2	1.89	0.54
1:B:164:PRO:HD2	1:B:167:LEU:HB2	1.90	0.54
1:B:182:VAL:HA	1:B:185:ILE:HD12	1.89	0.54
1:B:305:GLU:HG2	1:B:306:PRO:HA	1.90	0.53
1:A:248:THR:HA	1:A:536:VAL:HG21	1.90	0.52
1:A:164:PRO:HD2	1:A:167:LEU:HB2	1.91	0.51
1:A:329:PHE:HB3	1:A:411:VAL:HG11	1.93	0.50
1:A:74:ARG:HH21	1:A:99:GLN:HB2	1.76	0.50
1:B:329:PHE:HB3	1:B:411:VAL:HG11	1.93	0.49
1:B:539:LYS:HA	1:B:542:LEU:HD12	1.93	0.49
1:A:1015:LEU:HD22	1:A:1128:LEU:HD22	1.97	0.46
1:B:1067:SER:HB3	1:B:1070:SER:HB2	1.98	0.45
1:B:541:THR:HA	1:B:544:ILE:HD12	1.98	0.45
1:A:454:ILE:HD11	1:B:516:LEU:HD21	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:LEU:HD11	3:A:1203:CLR:H17	1.99	0.45
1:A:119:TRP:HA	1:A:122:ILE:HG22	1.99	0.45
1:B:1015:LEU:HD22	1:B:1128:LEU:HD22	1.99	0.44
1:B:536:VAL:CG1	1:B:536:VAL:O	2.65	0.44
1:A:158:GLU:HA	1:A:162:GLY:HA2	1.99	0.43
1:A:252:PHE:HD1	1:A:259:SER:HB2	1.83	0.43
1:B:119:TRP:HA	1:B:122:ILE:HG22	2.01	0.43
1:B:158:GLU:HA	1:B:162:GLY:HA2	1.99	0.43
1:B:221:LEU:HD22	1:B:509:ILE:HG21	2.00	0.42
1:A:362:LEU:O	1:A:366:SER:HB3	2.20	0.42
1:B:230:MET:HE1	1:B:519:LYS:HG3	2.01	0.42
1:A:408:ILE:HA	1:A:411:VAL:HG22	2.01	0.42
1:A:170:THR:HG22	1:A:172:ASP:H	1.83	0.41
1:B:408:ILE:HA	1:B:411:VAL:HG22	2.01	0.41
1:B:325:LEU:HD23	1:B:326:MET:HE2	2.03	0.41
1:A:1038:ALA:HB2	1:A:1099:VAL:HG22	2.01	0.41
1:A:114:ASN:HA	1:A:192:GLN:O	2.21	0.41
1:A:230:MET:HE1	1:A:519:LYS:HG3	2.02	0.41
1:A:1106:ALA:HA	1:A:1109:LEU:HG	2.02	0.41
1:A:325:LEU:HD23	1:A:326:MET:HE2	2.02	0.41
1:B:61:LEU:HB2	1:B:64:CYS:HB2	2.02	0.41
1:B:1098:LYS:HB3	1:B:1101:GLU:HB2	2.03	0.41
1:A:180:ASN:HB3	1:A:182:VAL:HG22	2.03	0.40
1:A:1098:LYS:HB3	1:A:1101:GLU:HB2	2.03	0.40
1:B:252:PHE:CD1	1:B:259:SER:HB3	2.53	0.40
1:A:252:PHE:CD1	1:A:259:SER:HB2	2.55	0.40
1:A:400:ARG:NH2	1:A:470:HIS:HD2	2.19	0.40
1:B:536:VAL:O	1:B:536:VAL:HG12	2.21	0.40
1:B:116:PRO:HA	1:B:191:GLY:HA3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	582/638 (91%)	561 (96%)	20 (3%)	1 (0%)	43	73
1	B	572/638 (90%)	553 (97%)	18 (3%)	1 (0%)	43	73
All	All	1154/1276 (90%)	1114 (96%)	38 (3%)	2 (0%)	43	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	176	GLU
1	A	115	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/529 (94%)	483 (98%)	12 (2%)	43	70
1	B	486/529 (92%)	477 (98%)	9 (2%)	50	73
All	All	981/1058 (93%)	960 (98%)	21 (2%)	47	71

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	ARG
1	A	161	ARG
1	A	176	GLU
1	A	251	THR
1	A	267	LEU
1	A	303	LEU
1	A	367	LEU
1	A	1018	LEU
1	A	1042	LYS
1	A	1111	THR
1	A	1121	LEU
1	A	1128	LEU
1	B	267	LEU
1	B	317	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	339	TRP
1	B	346	LEU
1	B	1042	LYS
1	B	1069	ASP
1	B	1092	LYS
1	B	1121	LEU
1	B	1128	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	HIS
1	A	114	ASN
1	A	184	ASN
1	A	192	GLN
1	A	219	ASN
1	A	260	ASN
1	A	380	GLN
1	A	1078	HIS
1	A	1103	GLN
1	A	470	HIS
1	A	521	ASN
1	B	63	HIS
1	B	219	ASN
1	B	260	ASN
1	B	380	GLN
1	B	1103	GLN
1	B	470	HIS
1	B	491	GLN
1	B	521	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1202	1	14,14,15	0.29	0	17,19,21	0.90	1 (5%)
3	CLR	A	1203	-	31,31,31	0.67	0	48,48,48	1.45	7 (14%)
2	NAG	A	1201	1	14,14,15	0.36	0	17,19,21	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1202	1	-	2/6/23/26	0/1/1/1
3	CLR	A	1203	-	-	5/10/68/68	0/4/4/4
2	NAG	A	1201	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1203	CLR	C4-C5-C10	3.94	121.47	116.42
3	A	1203	CLR	C4-C5-C6	-3.36	116.02	120.57
2	A	1202	NAG	C1-O5-C5	3.24	116.53	112.19
3	A	1203	CLR	C21-C20-C22	-3.07	105.58	110.34
3	A	1203	CLR	C1-C2-C3	3.00	114.46	110.48
3	A	1203	CLR	C7-C8-C14	2.67	114.72	110.93
3	A	1203	CLR	C12-C11-C9	2.35	117.13	113.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1203	CLR	C21-C20-C17	2.34	116.39	112.88

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1203	CLR	C17-C20-C22-C23
3	A	1203	CLR	C21-C20-C22-C23
2	A	1202	NAG	O5-C5-C6-O6
3	A	1203	CLR	C22-C23-C24-C25
3	A	1203	CLR	C23-C24-C25-C26
3	A	1203	CLR	C23-C24-C25-C27
2	A	1202	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1203	CLR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	588/638 (92%)	-0.09	2 (0%)	90 81	63, 116, 206, 272	0
1	B	578/638 (90%)	-0.08	4 (0%)	84 69	58, 138, 208, 221	0
All	All	1166/1276 (91%)	-0.08	6 (0%)	87 76	58, 126, 207, 272	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	128	ALA	3.2
1	B	142	PRO	2.9
1	A	1080	PHE	2.7
1	B	391	PHE	2.7
1	B	267	LEU	2.3
1	A	1106	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	1201	14/15	0.38	0.14	148,151,153,153	0
2	NAG	A	1202	14/15	0.68	0.14	171,172,174,175	0
4	NA	A	1204	1/1	0.78	0.11	105,105,105,105	0
4	NA	A	1205	1/1	0.91	0.16	71,71,71,71	0
3	CLR	A	1203	28/28	0.95	0.09	75,84,87,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.