



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 25, 2026 – 04:05 PM UTC

PDB ID : 5L7O / pdb_00005l7o
Title : X-ray structure of Triatoma virus empty capsid
Authors : Sanchez-Eugenio, R.
Deposited on : 2016-06-03
Resolution : 3.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1747 (3.70-3.50)
Clashscore	190562	1827 (3.70-3.50)
Ramachandran outliers	187476	1773 (3.70-3.50)
Sidechain outliers	187428	1772 (3.70-3.50)
RSRZ outliers	180081	1745 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	10% 68% 16% • 15%
2	B	255	5% 52% 17% • 30%
3	C	285	13% 80% 16% ••

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5398 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1785	1149	288	341	7			

- Molecule 2 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	179	Total	C	N	O	S	0	0	0
			1417	932	228	253	4			

- Molecule 3 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	276	Total	C	N	O	S	0	0	0
			2196	1421	356	415	4			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	54	MET	VAL	conflict	UNP Q9QEY5


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Capsid protein



• Molecule 2: Capsid protein



• Molecule 3: Capsid protein





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	305.59Å 305.59Å 796.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.91 – 3.60 49.91 – 3.60	Depositor EDS
% Data completeness (in resolution range)	98.3 (49.91-3.60) 98.3 (49.91-3.60)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 3.57Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.269 , 0.269 0.270 , 0.270	Depositor DCC
R_{free} test set	6433 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å ²)	112.7	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 80.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.048 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.46	EDS
Total number of atoms	5398	wwPDB-VP
Average B, all atoms (Å ²)	114.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1831	0.84	1/2497 (0.0%)
2	B	0.49	0/1459	0.89	4/1994 (0.2%)
3	C	0.51	0/2259	0.82	3/3095 (0.1%)
All	All	0.50	0/5549	0.84	8/7586 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	163	VAL	N-CA-C	6.70	118.37	108.65
2	B	142	HIS	CA-C-N	5.63	126.87	119.84
2	B	142	HIS	C-N-CA	5.63	126.87	119.84
1	A	194	ALA	N-CA-C	-5.45	105.73	112.38
2	B	199	SER	N-CA-C	5.33	112.92	108.13
3	C	166	VAL	N-CA-C	5.29	114.41	106.42
2	B	238	ALA	N-CA-C	5.07	116.77	108.76
3	C	72	SER	N-CA-C	5.00	115.65	108.74

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1785	0	1753	34	0
2	B	1417	0	1425	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	2196	0	2165	29	0
All	All	5398	0	5343	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:133:HIS:HB3	3:C:233:LEU:HD11	1.68	0.75
3:C:53:GLU:HA	3:C:58:TYR:CD2	2.31	0.66
1:A:198:SER:O	1:A:203:ASN:ND2	2.31	0.63
1:A:87:LYS:HG2	1:A:87:LYS:O	1.99	0.62
1:A:81:MET:HE2	3:C:275:ARG:HA	1.82	0.61
3:C:80:VAL:HG21	3:C:151:MET:HB3	1.83	0.59
1:A:68:LEU:HD11	1:A:147:ILE:HG23	1.88	0.56
3:C:96:THR:HG21	3:C:272:LEU:O	2.05	0.56
1:A:85:ALA:HB3	1:A:88:LEU:HD12	1.88	0.55
2:B:204:TYR:OH	2:B:209:LYS:HG2	2.07	0.54
2:B:183:LEU:C	2:B:183:LEU:HD23	2.32	0.54
1:A:133:VAL:HG22	1:A:165:PHE:CE1	2.43	0.54
3:C:37:GLY:O	3:C:38:ASN:C	2.53	0.52
1:A:118:ARG:HD3	3:C:27:ASP:CG	2.36	0.50
3:C:71:TYR:OH	3:C:231:THR:O	2.25	0.50
1:A:105:SER:HA	3:C:268:PRO:HG3	1.92	0.49
3:C:129:LYS:HE3	3:C:170:LEU:O	2.12	0.49
3:C:236:SER:OG	3:C:238:ALA:HB3	2.12	0.49
1:A:92:THR:HA	1:A:253:VAL:CG2	2.42	0.49
3:C:182:PHE:CE2	3:C:184:SER:HB3	2.49	0.48
1:A:81:MET:CE	3:C:275:ARG:HA	2.43	0.47
3:C:178:LEU:C	3:C:178:LEU:HD12	2.39	0.47
1:A:108:ALA:HB1	1:A:234:PRO:HB3	1.95	0.47
2:B:154:VAL:HG22	2:B:155:PRO:HD2	1.96	0.47
2:B:202:VAL:HG23	2:B:203:GLN:HG3	1.97	0.47
1:A:81:MET:O	1:A:91:TRP:HA	2.15	0.47
2:B:158:ASP:HB2	2:B:211:THR:HG23	1.95	0.47
2:B:156:VAL:O	2:B:157:PRO:C	2.58	0.47
3:C:118:ARG:NH1	3:C:187:PRO:O	2.44	0.46
2:B:247:PHE:CD2	2:B:248:PRO:HD2	2.50	0.46
3:C:52:ASP:OD1	3:C:54:MET:N	2.49	0.46
2:B:100:ILE:HD12	2:B:101:GLU:O	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:146:CYS:C	2:B:185:ILE:HD11	2.40	0.46
1:A:56:ARG:HG3	3:C:21:ASN:HA	1.97	0.46
1:A:92:THR:HA	1:A:253:VAL:HG21	1.97	0.46
1:A:101:THR:HG21	1:A:219:HIS:CE1	2.51	0.46
2:B:81:ILE:HG23	2:B:243:ILE:HD13	1.97	0.46
2:B:135:PHE:O	2:B:192:LEU:HA	2.15	0.46
2:B:211:THR:HG22	2:B:212:PRO:O	2.16	0.46
1:A:193:ASN:C	1:A:195:GLY:N	2.72	0.45
3:C:49:THR:HG22	3:C:51:SER:N	2.31	0.45
2:B:204:TYR:OH	2:B:209:LYS:CG	2.65	0.45
3:C:131:ASN:OD1	3:C:131:ASN:N	2.50	0.45
1:A:36:GLU:N	1:A:36:GLU:OE1	2.49	0.45
1:A:133:VAL:CG2	1:A:165:PHE:CE1	2.99	0.45
3:C:49:THR:HG22	3:C:51:SER:H	1.81	0.45
1:A:177:TYR:CE2	1:A:179:SER:HB3	2.52	0.45
3:C:52:ASP:OD1	3:C:52:ASP:C	2.59	0.45
1:A:233:ALA:HB2	3:C:106:LEU:HA	1.98	0.45
1:A:139:THR:O	1:A:140:MET:C	2.61	0.44
1:A:143:GLN:O	1:A:195:GLY:C	2.60	0.44
2:B:116:SER:O	2:B:120:ARG:HG3	2.17	0.44
1:A:121:ILE:HB	1:A:171:ALA:HB3	1.98	0.44
2:B:120:ARG:NH2	2:B:254:ALA:O	2.51	0.44
3:C:60:LEU:N	3:C:60:LEU:HD23	2.33	0.44
2:B:114:LEU:HD12	2:B:240:PHE:CZ	2.53	0.44
2:B:106:PHE:CD1	2:B:106:PHE:C	2.96	0.43
3:C:275:ARG:O	3:C:276:PRO:C	2.62	0.43
1:A:203:ASN:ND2	1:A:203:ASN:N	2.65	0.42
3:C:100:ALA:HB3	3:C:101:PRO:HD3	2.00	0.42
1:A:87:LYS:O	1:A:87:LYS:CG	2.67	0.42
1:A:176:PRO:HB2	1:A:178:TYR:CD2	2.55	0.42
2:B:87:PRO:HA	2:B:239:HIS:HB3	2.01	0.42
1:A:37:GLN:HB3	1:A:38:PRO:HD3	2.01	0.42
1:A:177:TYR:O	1:A:178:TYR:C	2.63	0.42
1:A:194:ALA:O	1:A:195:GLY:C	2.63	0.42
3:C:193:ASP:O	3:C:208:ARG:NH1	2.53	0.41
1:A:71:PRO:HB3	1:A:205:MET:HE3	2.03	0.41
1:A:97:SER:HA	1:A:147:ILE:CD1	2.51	0.41
2:B:94:LEU:HD11	2:B:231:SER:HB2	2.02	0.41
2:B:138:GLN:HB2	2:B:235:ASN:HB2	2.03	0.41
2:B:222:ALA:HB2	3:C:247:LEU:HD11	2.02	0.41
1:A:146:ILE:CG2	1:A:147:ILE:N	2.84	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:183:LEU:HD23	2:B:184:ASP:N	2.35	0.41
3:C:158:VAL:O	3:C:158:VAL:HG23	2.21	0.40
1:A:161:ILE:HB	3:C:30:LEU:HD22	2.04	0.40
2:B:111:PRO:O	2:B:112:ASP:C	2.64	0.40
2:B:162:LEU:O	2:B:163:HIS:C	2.65	0.40
1:A:128:GLN:HB3	1:A:129:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/271 (84%)	211 (92%)	17 (8%)	0	100	100
2	B	177/255 (69%)	163 (92%)	13 (7%)	1 (1%)	21	54
3	C	274/285 (96%)	246 (90%)	28 (10%)	0	100	100
All	All	679/811 (84%)	620 (91%)	58 (8%)	1 (0%)	48	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	211	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/231 (85%)	189 (96%)	7 (4%)	31	56
2	B	161/231 (70%)	157 (98%)	4 (2%)	42	63
3	C	251/258 (97%)	244 (97%)	7 (3%)	38	60
All	All	608/720 (84%)	590 (97%)	18 (3%)	36	59

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	SER
1	A	64	ASP
1	A	92	THR
1	A	161	ILE
1	A	163	GLU
1	A	219	HIS
1	A	262	TYR
2	B	133	THR
2	B	199	SER
2	B	207	VAL
2	B	217	LEU
3	C	12	VAL
3	C	50	SER
3	C	51	SER
3	C	131	ASN
3	C	183	THR
3	C	228	SER
3	C	270	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	174	ASN
2	B	233	GLN
2	B	235	ASN
3	C	28	GLN
3	C	87	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/271 (84%)	1.11	27 (11%) 9 8	87, 113, 146, 211	0
2	B	179/255 (70%)	0.84	12 (6%) 24 15	88, 111, 129, 170	0
3	C	276/285 (96%)	1.04	36 (13%) 7 6	89, 111, 133, 162	0
All	All	685/811 (84%)	1.01	75 (10%) 10 9	87, 111, 139, 211	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	TYR	5.7
1	A	193	ASN	4.6
1	A	141	GLN	3.9
1	A	160	ASN	3.6
1	A	264	SER	3.5
3	C	175	GLU	3.4
3	C	276	PRO	3.1
2	B	138	GLN	3.0
2	B	231	SER	3.0
1	A	194	ALA	3.0
3	C	164	TYR	3.0
3	C	165	ARG	3.0
3	C	193	ASP	3.0
1	A	256	PRO	3.0
3	C	119	GLY	3.0
3	C	27	ASP	2.9
3	C	242	SER	2.9
3	C	87	HIS	2.9
3	C	245	ASP	2.9
1	A	58	SER	2.9
3	C	269	VAL	2.8
3	C	205	TYR	2.7
3	C	153	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	67	SER	2.7
2	B	136	HIS	2.6
3	C	132	TYR	2.6
1	A	84	ALA	2.6
1	A	187	SER	2.6
3	C	131	ASN	2.5
3	C	173	GLN	2.5
1	A	240	THR	2.5
3	C	270	ASP	2.5
3	C	126	ARG	2.4
1	A	44	GLY	2.4
2	B	247	PHE	2.4
3	C	57	GLU	2.4
3	C	61	LYS	2.4
3	C	251	LYS	2.4
3	C	86	VAL	2.3
3	C	65	TYR	2.3
2	B	121	GLU	2.3
3	C	70	LYS	2.3
3	C	120	GLY	2.3
1	A	48	LEU	2.2
1	A	134	GLU	2.2
3	C	148	ASP	2.2
3	C	160	ASP	2.2
1	A	35	ASP	2.2
2	B	113	ALA	2.2
1	A	90	VAL	2.2
1	A	169	GLY	2.2
2	B	202	VAL	2.1
3	C	26	VAL	2.1
1	A	87	LYS	2.1
1	A	163	GLU	2.1
3	C	273	THR	2.1
3	C	68	ARG	2.1
3	C	191	CYS	2.1
2	B	101	GLU	2.1
2	B	77	THR	2.1
3	C	223	GLY	2.1
3	C	16	PRO	2.1
2	B	96	ASN	2.1
1	A	99	PHE	2.1
3	C	256	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	102	GLY	2.1
1	A	123	PRO	2.1
2	B	140	ASN	2.1
1	A	179	SER	2.0
1	A	96	ARG	2.0
2	B	143	PRO	2.0
1	A	59	TRP	2.0
1	A	221	ALA	2.0
1	A	66	LYS	2.0
1	A	250	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.