



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:48 AM UTC

PDB ID : 1L9S / pdb_00001l9s
Title : CRYSTAL STRUCTURE OF THE I257T VARIANT OF THE COPPER-CONTAINING NITRITE REDUCTASE FROM ALCALIGENES FAECALIS S-6
Authors : Boulanger, M.J.; Murphy, M.E.P.
Deposited on : 2002-03-26
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	82% 15% ..
1	B	341	82% 15% ..
1	C	341	79% 18% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COPPER-CONTAINING NITRITE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2559	1637	430	481	11			
1	B	336	Total	C	N	O	S	0	0	0
			2559	1637	430	481	11			
1	C	336	Total	C	N	O	S	0	0	0
			2559	1637	430	481	11			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	257	THR	ILE	engineered mutation	UNP P38501
A	341	ILE	-	cloning artifact	UNP P38501
A	342	GLU	-	cloning artifact	UNP P38501
A	343	GLY	-	cloning artifact	UNP P38501
A	344	ARG	-	cloning artifact	UNP P38501
B	257	THR	ILE	engineered mutation	UNP P38501
B	341	ILE	-	cloning artifact	UNP P38501
B	342	GLU	-	cloning artifact	UNP P38501
B	343	GLY	-	cloning artifact	UNP P38501
B	344	ARG	-	cloning artifact	UNP P38501
C	257	THR	ILE	engineered mutation	UNP P38501
C	341	ILE	-	cloning artifact	UNP P38501
C	342	GLU	-	cloning artifact	UNP P38501
C	343	GLY	-	cloning artifact	UNP P38501
C	344	ARG	-	cloning artifact	UNP P38501

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Cu	0	0
			2	2		
2	C	2	Total	Cu	0	0
			2	2		

- Molecule 3 is NITRITE ION (CCD ID: NO2) (formula: NO₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			3	1	2		
3	B	1	Total	N	O	0	0
			3	1	2		
3	C	1	Total	N	O	0	0
			3	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	495	Total	O	0	0
			495	495		
4	B	482	Total	O	0	0
			482	482		
4	C	453	Total	O	0	0
			453	453		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: COPPER-CONTAINING NITRITE REDUCTASE



• Molecule 1: COPPER-CONTAINING NITRITE REDUCTASE



• Molecule 1: COPPER-CONTAINING NITRITE REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.48Å 102.03Å 146.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.54 – 1.78 25.54 – 1.78	Depositor EDS
% Data completeness (in resolution range)	87.0 (25.54-1.78) 84.8 (25.54-1.78)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.56 (at 1.78Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.157 , 0.197 0.163 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9122	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, NO2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2630	1.07	20/3587 (0.6%)
1	B	0.52	0/2630	1.05	19/3587 (0.5%)
1	C	0.49	0/2630	1.07	23/3587 (0.6%)
All	All	0.51	0/7890	1.06	62/10761 (0.6%)

There are no bond length outliers.

All (62) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307	ASN	N-CA-C	-11.13	90.49	108.41
1	C	307	ASN	N-CA-C	-10.67	89.21	107.99
1	B	307	ASN	N-CA-C	-10.13	90.15	107.99
1	B	305	ASN	N-CA-C	-9.30	94.17	108.96
1	C	79	ASP	N-CA-C	-9.06	97.44	110.59
1	A	305	ASN	N-CA-C	-9.04	94.58	108.96
1	C	305	ASN	N-CA-C	-8.93	94.76	108.96
1	A	79	ASP	N-CA-C	-8.87	99.09	110.53
1	B	79	ASP	N-CA-C	-8.55	99.50	110.53
1	C	63	ALA	N-CA-C	8.07	121.67	108.52
1	B	63	ALA	N-CA-C	7.78	121.09	108.41
1	A	63	ALA	N-CA-C	7.30	120.30	108.41
1	B	222	GLY	N-CA-C	7.01	124.68	115.36
1	C	222	GLY	N-CA-C	6.94	124.59	115.36
1	C	185	VAL	N-CA-C	6.82	114.65	107.76
1	B	203	TYR	N-CA-C	6.81	119.33	111.02
1	A	222	GLY	N-CA-C	6.39	123.85	115.36
1	B	306	HIS	N-CA-C	6.26	120.41	113.21
1	B	47	GLU	N-CA-C	-6.18	99.13	108.96
1	B	148	SER	N-CA-C	-6.09	105.43	112.92
1	C	148	SER	N-CA-C	-6.09	105.43	112.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	270	PHE	N-CA-C	6.05	119.76	112.38
1	A	306	HIS	N-CA-C	6.04	120.15	113.21
1	A	309	ILE	N-CA-C	-6.00	104.66	110.72
1	A	52	ILE	N-CA-C	5.97	118.63	112.90
1	C	309	ILE	N-CA-C	-5.76	104.90	110.72
1	A	315	GLY	N-CA-C	5.73	123.25	115.00
1	C	275	ASP	N-CA-C	-5.70	102.86	110.55
1	B	185	VAL	N-CA-C	5.69	114.14	107.77
1	C	219	VAL	N-CA-C	5.65	117.05	108.80
1	A	185	VAL	N-CA-C	5.60	113.41	107.76
1	A	275	ASP	N-CA-C	-5.56	103.05	110.55
1	B	329	ASP	N-CA-C	-5.51	105.19	111.14
1	A	303	TYR	N-CA-C	-5.50	99.32	108.34
1	C	303	TYR	N-CA-C	-5.47	99.37	108.34
1	B	261	GLY	N-CA-C	-5.42	100.33	113.18
1	B	309	ILE	N-CA-C	-5.40	105.27	110.72
1	B	275	ASP	N-CA-C	-5.36	103.31	110.55
1	A	304	VAL	N-CA-C	5.34	116.39	108.65
1	C	104	GLY	N-CA-C	5.33	117.01	111.95
1	B	108	GLY	N-CA-C	-5.32	107.67	114.37
1	C	52	ILE	N-CA-C	5.32	119.76	113.22
1	C	302	ALA	N-CA-C	-5.31	101.88	110.32
1	B	304	VAL	N-CA-C	5.29	116.52	108.80
1	C	304	VAL	N-CA-C	5.25	116.27	108.65
1	A	158	PRO	N-CA-C	-5.24	103.05	111.38
1	A	302	ALA	N-CA-C	-5.20	101.80	110.17
1	C	216	THR	N-CA-C	-5.17	106.82	113.23
1	B	302	ALA	N-CA-C	-5.17	101.85	110.17
1	C	261	GLY	N-CA-C	-5.16	100.96	113.18
1	A	104	GLY	N-CA-C	5.11	116.81	111.95
1	A	134	TYR	N-CA-C	-5.10	101.44	109.50
1	C	86	ILE	N-CA-C	5.09	115.23	108.11
1	A	148	SER	N-CA-C	-5.09	106.66	112.92
1	A	261	GLY	N-CA-C	-5.03	101.27	113.18
1	C	313	GLU	N-CA-C	-5.03	107.22	113.55
1	A	265	TRP	N-CA-C	-5.03	99.81	108.56
1	C	87	ASN	CA-C-N	5.02	125.01	119.89
1	C	87	ASN	C-N-CA	5.02	125.01	119.89
1	C	158	PRO	N-CA-C	-5.02	103.22	111.21
1	B	315	GLY	N-CA-C	5.01	122.21	115.00
1	B	303	TYR	N-CA-C	-5.00	99.12	108.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2559	0	2481	26	0
1	B	2559	0	2481	28	0
1	C	2559	0	2481	36	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	3	0	0	1	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
4	A	495	0	0	5	0
4	B	482	0	0	8	0
4	C	453	0	0	10	0
All	All	9122	0	7443	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:LYS:HB2	4:B:5215:HOH:O	1.70	0.91
1:A:26:HIS:HE1	1:A:74:VAL:H	1.28	0.80
1:C:26:HIS:HE1	1:C:74:VAL:H	1.32	0.78
1:B:26:HIS:HE1	1:B:74:VAL:H	1.30	0.77
1:A:160:GLU:HB3	4:A:5354:HOH:O	1.88	0.74
1:A:228:THR:HG21	4:A:5449:HOH:O	1.88	0.73
1:B:13:PRO:HG2	1:B:37:LYS:HG2	1.70	0.73
1:C:26:HIS:CE1	1:C:74:VAL:H	2.08	0.70
1:B:26:HIS:CE1	1:B:74:VAL:H	2.10	0.70
1:B:141:MET:HG3	4:B:3145:HOH:O	1.91	0.69
1:C:18:GLU:H	1:C:18:GLU:CD	2.01	0.69
1:A:26:HIS:CE1	1:A:74:VAL:H	2.11	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:22:PRO:HB2	1:C:221:ASN:HD21	1.58	0.68
1:A:5:THR:OG1	1:A:8:GLU:HG3	1.97	0.64
1:C:88:PRO:HD2	1:C:91:ASN:ND2	2.13	0.64
1:C:77:GLN:O	1:C:159:ARG:HD3	1.98	0.63
1:C:231:LYS:HD3	4:C:5324:HOH:O	1.99	0.62
1:C:231:LYS:HA	4:C:5636:HOH:O	1.98	0.62
1:C:300:ILE:HG22	4:C:4420:HOH:O	2.00	0.61
1:C:26:HIS:HD2	1:C:27:ALA:O	1.84	0.61
1:C:192:LYS:HE3	4:C:5661:HOH:O	2.00	0.60
1:B:58:GLU:CD	1:B:195:LYS:HE3	2.26	0.59
1:C:163:HIS:HD2	1:C:167:GLY:O	1.85	0.59
1:C:141:MET:HE2	1:C:203:TYR:CE2	2.38	0.58
1:C:141:MET:SD	4:C:4145:HOH:O	2.57	0.57
1:B:296:GLN:NE2	4:B:2128:HOH:O	2.37	0.57
1:A:136:CYS:HB2	1:A:150:MET:HG2	1.88	0.55
1:B:22:PRO:HB2	1:B:221:ASN:HD21	1.71	0.55
1:B:49:LYS:NZ	4:B:5537:HOH:O	2.40	0.55
1:C:60:HIS:HD2	4:C:5013:HOH:O	1.89	0.55
1:C:166:LYS:N	1:C:166:LYS:HD2	2.22	0.54
1:B:18:GLU:HG3	4:B:5588:HOH:O	2.07	0.54
1:A:307:ASN:HA	1:C:249:ASN:O	2.07	0.54
1:A:300:ILE:HG22	4:A:2363:HOH:O	2.08	0.54
1:A:321:LYS:HD2	4:A:1321:HOH:O	2.06	0.53
1:A:22:PRO:HB2	1:A:221:ASN:HD21	1.73	0.53
1:C:95:HIS:HB3	1:C:150:MET:HE1	1.90	0.53
1:B:249:ASN:O	1:C:307:ASN:HA	2.10	0.52
1:B:168:LYS:N	1:B:168:LYS:HE3	2.25	0.52
1:B:143:PRO:O	1:B:147:VAL:HG22	2.11	0.51
1:A:26:HIS:HD2	1:A:27:ALA:O	1.94	0.51
1:C:18:GLU:HG2	4:C:5184:HOH:O	2.11	0.50
1:B:15:GLN:O	1:B:15:GLN:HG3	2.11	0.50
1:A:249:ASN:O	1:B:307:ASN:HA	2.12	0.50
1:A:4:ALA:HB1	1:A:8:GLU:OE1	2.11	0.49
1:C:18:GLU:CD	1:C:18:GLU:N	2.71	0.48
1:A:164:ASP:OD1	1:A:168:LYS:HB2	2.13	0.47
1:A:143:PRO:O	1:A:147:VAL:HG22	2.14	0.47
1:B:300:ILE:HG22	4:B:3425:HOH:O	2.13	0.47
1:C:130:GLY:HA2	1:C:270:PHE:CD1	2.49	0.47
1:C:37:LYS:HE2	4:C:5323:HOH:O	2.14	0.47
1:A:16:LYS:HE3	4:A:5515:HOH:O	2.15	0.45
1:C:141:MET:HE1	4:C:4200:HOH:O	2.15	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:GLU:OE1	1:B:195:LYS:HE3	2.15	0.45
1:A:235:ALA:O	1:A:322:VAL:HA	2.17	0.45
1:C:77:GLN:O	1:C:78:ASP:HB2	2.16	0.45
1:C:310:GLU:HA	1:C:314:LEU:HB2	1.97	0.45
1:A:132:PHE:CE1	1:A:269:LYS:HE3	2.52	0.44
1:A:196:TYR:CD2	1:A:202:ALA:HB2	2.52	0.44
3:A:704:NO2:O1	1:C:135:HIS:CE1	2.71	0.44
1:B:26:HIS:HD2	1:B:27:ALA:O	2.00	0.44
1:B:163:HIS:HD2	1:B:167:GLY:O	2.01	0.44
1:B:174:LYS:HE2	4:B:3174:HOH:O	2.17	0.43
1:A:174:LYS:HE2	1:A:176:TYR:CE2	2.53	0.43
1:A:255:HIS:ND1	1:A:279:GLU:O	2.51	0.43
1:B:95:HIS:HB3	1:B:150:MET:HE1	2.00	0.43
1:C:160:GLU:HG2	1:C:163:HIS:CE1	2.53	0.43
1:B:310:GLU:HA	1:B:314:LEU:HB2	2.01	0.43
1:C:64:PHE:CE2	1:C:150:MET:HG2	2.54	0.42
1:B:301:TYR:HB2	1:B:320:PHE:HB2	2.02	0.42
1:B:136:CYS:HB2	1:B:150:MET:HG2	2.01	0.42
1:C:5:THR:OG1	1:C:8:GLU:HG3	2.20	0.42
1:C:133:VAL:CG1	1:C:134:TYR:N	2.83	0.42
1:A:81:LEU:C	1:A:81:LEU:HD23	2.45	0.41
1:A:321:LYS:HD3	1:A:321:LYS:C	2.45	0.41
1:C:253:ARG:HA	1:C:281:TRP:O	2.20	0.41
1:B:132:PHE:CE1	1:B:269:LYS:HE3	2.56	0.41
1:B:231:LYS:HD3	4:B:5110:HOH:O	2.20	0.41
1:B:77:GLN:O	1:B:78:ASP:HB2	2.21	0.41
1:A:310:GLU:HA	1:A:314:LEU:HB2	2.01	0.41
1:A:313:GLU:O	1:C:211:ARG:NH2	2.53	0.41
1:C:235:ALA:O	1:C:322:VAL:HA	2.21	0.41
1:C:301:TYR:HB2	1:C:320:PHE:HB2	2.02	0.41
1:C:33:GLU:HG2	4:C:4031:HOH:O	2.20	0.40
1:A:321:LYS:C	1:A:321:LYS:CD	2.94	0.40
1:B:22:PRO:HB2	1:B:221:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/341 (98%)	329 (98%)	4 (1%)	1 (0%)	36	22
1	B	334/341 (98%)	328 (98%)	6 (2%)	0	100	100
1	C	334/341 (98%)	327 (98%)	7 (2%)	0	100	100
All	All	1002/1023 (98%)	984 (98%)	17 (2%)	1 (0%)	48	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	THR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/268 (98%)	256 (97%)	8 (3%)	36	15
1	B	264/268 (98%)	257 (97%)	7 (3%)	39	19
1	C	264/268 (98%)	257 (97%)	7 (3%)	39	19
All	All	792/804 (98%)	770 (97%)	22 (3%)	38	17

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	LYS
1	A	83	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	221	ASN
1	A	247	GLN
1	A	300	ILE
1	A	305	ASN
1	A	314	LEU
1	A	321	LYS
1	B	83	LEU
1	B	168	LYS
1	B	174	LYS
1	B	221	ASN
1	B	247	GLN
1	B	305	ASN
1	B	314	LEU
1	C	18	GLU
1	C	83	LEU
1	C	166	LYS
1	C	221	ASN
1	C	247	GLN
1	C	314	LEU
1	C	332	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	26	HIS
1	A	163	HIS
1	A	221	ASN
1	B	26	HIS
1	B	77	GLN
1	B	115	ASN
1	B	221	ASN
1	B	296	GLN
1	C	26	HIS
1	C	60	HIS
1	C	115	ASN
1	C	163	HIS
1	C	221	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NO2	B	504	2	1,2,2	3.64	1 (100%)	0,1,1	-	-
3	NO2	C	604	2	1,2,2	3.53	1 (100%)	0,1,1	-	-
3	NO2	A	704	2	1,2,2	3.68	1 (100%)	0,1,1	-	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	704	NO2	O1-N	3.68	1.40	1.22
3	B	504	NO2	O1-N	3.64	1.40	1.22
3	C	604	NO2	O1-N	3.53	1.40	1.22

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	704	NO2	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/341 (98%)	-0.35	2 (0%) 85 89	12, 18, 32, 47	0
1	B	336/341 (98%)	-0.38	1 (0%) 90 93	13, 18, 35, 48	0
1	C	336/341 (98%)	-0.14	2 (0%) 85 89	14, 23, 36, 46	0
All	All	1008/1023 (98%)	-0.29	5 (0%) 87 91	12, 19, 35, 48	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	GLN	3.2
1	C	229	GLY	3.1
1	A	4	ALA	2.7
1	C	168	LYS	2.5
1	A	339	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NO2	B	504	3/3	0.91	0.10	27,27,29,30	0
3	NO2	C	604	3/3	0.91	0.10	27,27,33,35	0
3	NO2	A	704	3/3	0.92	0.15	35,35,35,36	0
2	CU	C	502	1/1	0.99	0.03	22,22,22,22	0
2	CU	C	501	1/1	0.99	0.06	24,24,24,24	0
2	CU	A	502	1/1	1.00	0.03	17,17,17,17	0
2	CU	B	501	1/1	1.00	0.03	19,19,19,19	0
2	CU	B	502	1/1	1.00	0.04	20,20,20,20	0
2	CU	A	501	1/1	1.00	0.04	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.