



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 10:43 AM UTC

PDB ID : 6L9C / pdb_00006l9c
Title : Neutron structure of copper amine oxidase from *Arthrobacter glibiformis* at pD 7.4
Authors : Murakawa, T.; Kurihara, K.; Shoji, M.; Shibasaki, C.; Sunami, T.; Tamada, T.; Yano, N.; Yamada, T.; Kusaka, K.; Suzuki, M.; Shigeta, Y.; Kuroki, R.; Hayashi, H.; Yano, Y.; Tanizawa, K.; Adachi, M.; Okajima, T.
Deposited on : 2019-11-08
Resolution : 1.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	FAILED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION, NEUTRON DIFFRACTION

The reported resolution of this entry is 1.14 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1393 (1.16-1.12)
Ramachandran outliers	187476	1369 (1.16-1.12)
Sidechain outliers	187428	1369 (1.16-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	X	622	 95% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13687 atoms, of which 4809 are hydrogens and 2620 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylethylamine oxidase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	X	621	Total	C	D	H	N	O	S	0	587	0
			11015	3317	1013	4809	891	971	14			

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	1	Total	Cu	0	0
			1	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	X	1063	Total	D	O	0	1
			2670	1607	1063		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: Phenylethylamine oxidase

Chain X:  95% 5%



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	157.55Å 61.78Å 92.33Å 90.00° 112.13° 90.00°	Depositor
Resolution (Å)	45.03 – 1.14	Depositor
% Data completeness (in resolution range)	99.3 (45.03-1.14)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.14Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.168 , 0.181	Depositor
Wilson B-factor (Å ²)	15.7	Xtriage
Anisotropy	0.386	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13687	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASA, CU, TPQ, E9C, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.61	2/8789 (0.0%)	0.75	2/11953 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	151[B]	ARG	CB-CG	-5.47	1.36	1.52
1	X	151[A]	ARG	CB-CG	-5.47	1.36	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	151[B]	ARG	CG-CD-NE	-7.27	96.02	112.00
1	X	151[A]	ARG	CG-CD-NE	-7.27	96.02	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	6206	4809	1132	28	0
2	X	1	0	0	0	0
3	X	1	0	0	0	0
4	X	2670	0	0	20	8

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8878	4809	1132	33	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:393[B]:ASP:OD2	4:X:801:HOH:O	1.73	1.06
1:X:164[A]:GLU:OE2	4:X:802:HOH:O	1.76	1.02
1:X:614[B]:GLU:OE1	4:X:803:HOH:O	1.80	0.98
1:X:614[A]:GLU:OE2	4:X:804:HOH:O	1.84	0.96
1:X:561[A]:HIS:ND1	1:X:562[A]:SER:O	2.13	0.80
1:X:413[A]:GLU:OE2	4:X:805:HOH:O	2.00	0.80
1:X:296[A]:TYR:OH	4:X:806:HOH:O	2.00	0.69
1:X:610[B]:LYS:NZ	4:X:838:HOH:O	2.45	0.49
1:X:562[C]:SER:O	4:X:814:HOH:O	2.24	0.47
1:X:471[A]:SER:OG	4:X:811:HOH:O	2.23	0.47
1:X:603[B]:PRO:HG2	4:X:992:HOH:O	2.12	0.43
1:X:565[B]:ALA:CB	4:X:1417:HOH:O	2.66	0.43
1:X:565[B]:ALA:HB1	4:X:1417:HOH:O	2.15	0.42
1:X:103[A]:GLU:HG3	4:X:842:HOH:O	2.15	0.42
1:X:381[A]:ASN:HD22	1:X:381[A]:ASN:H	1.56	0.41
1:X:563[A]:GLY:HA2	4:X:871:HOH:O	2.15	0.41
1:X:564[B]:GLY:O	1:X:569[B]:SER:OG	2.37	0.41
1:X:296[A]:TYR:CG	1:X:382[A]:E9C:OAJ	2.75	0.40
1:X:614[B]:GLU:HB2	4:X:1039:HOH:O	2.15	0.40

All (8) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:X:1486:HOH:O	4:X:1486:HOH:O[2_556]	1.78	0.42
4:X:804:HOH:O	4:X:930:HOH:O[2_556]	1.86	0.34
4:X:930:HOH:O	4:X:1088:HOH:O[2_556]	2.00	0.20
4:X:1142:HOH:O	4:X:1538:HOH:O[2_556]	2.01	0.19
4:X:1287:HOH:O	4:X:1433:HOH:O[2_556]	2.11	0.09
4:X:808:HOH:O	4:X:1433:HOH:O[2_556]	2.12	0.08
4:X:1294:HOH:O	4:X:1521:HOH:O[2_556]	2.19	0.01
4:X:1810:HOH:O	4:X:1814:HOH:O[2_556]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	1084/622 (174%)	1049 (97%)	33 (3%)	2 (0%)	43 13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	55[A]	ALA
1	X	55[B]	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	907/513 (177%)	891 (98%)	16 (2%)	51 13

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	53[A]	SER
1	X	53[B]	SER
1	X	54[A]	GLU
1	X	54[B]	GLU
1	X	164[A]	GLU
1	X	164[B]	GLU
1	X	267[A]	ARG
1	X	267[B]	ARG
1	X	371[A]	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	371[B]	MET
1	X	376[A]	PHE
1	X	376[B]	PHE
1	X	444[A]	ASP
1	X	444[B]	ASP
1	X	541[A]	LEU
1	X	541[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	E9C	X	382[A]	1	12,14,15	1.96	3 (25%)	10,19,21	1.40	0
1	TPQ	X	382[B]	1	13,14,15	0.98	1 (7%)	13,19,21	1.36	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	E9C	X	382[A]	1	-	1/5/22/24	0/1/1/1
1	TPQ	X	382[B]	1	-	1/5/22/24	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	382[A]	E9C	CD2-CG	-3.73	1.45	1.49
1	X	382[A]	E9C	CB-CG	-3.62	1.44	1.50
1	X	382[B]	TPQ	C1-C2	2.57	1.52	1.49
1	X	382[A]	E9C	CD1-CG	2.57	1.41	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	382[B]	TPQ	C6-C1-C2	3.92	121.45	118.66

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	X	382[A]	E9C	O-C-CA-CB
1	X	382[B]	TPQ	O-C-CA-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	X	382[A]	E9C	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

6.4 Ligands

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers

EDS failed to run properly - this section is therefore empty.