



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:26 AM UTC

PDB ID : 9L92 / pdb_00009L92
Title : The crystal structure of human RyR3 Repeat12 domain
Authors : Hadiatullah, H.; Lin, L.; Yuchi, Z.
Deposited on : 2024-12-29
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	9% 87% 8% .
1	B	215	13% 83% 7% 8%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ryanodine receptor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	206	Total	C	N	O	S	0	0	0
			1598	1014	282	295	7			
1	B	198	Total	C	N	O	S	0	0	0
			1503	948	262	286	7			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	145	Total 145	O 145	0	0
3	B	158	Total 158	O 158	0	0

- Molecule 1: Ryanodine receptor 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.04Å 122.15Å 47.71Å 90.00° 93.27° 90.00°	Depositor
Resolution (Å)	47.60 – 1.97 47.60 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.60-1.97) 99.7 (47.60-1.97)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 1.97Å)	Xtriage
Refinement program	PHENIX 1.14	Depositor
R, R_{free}	0.228 , 0.279 0.229 , 0.278	Depositor DCC
R_{free} test set	2017 reflections (6.20%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3428	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	6/1636 (0.4%)	1.36	5/2230 (0.2%)
1	B	0.84	6/1534 (0.4%)	1.42	7/2093 (0.3%)
All	All	0.84	12/3170 (0.4%)	1.39	12/4323 (0.3%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	882	MET	SD-CE	6.20	1.95	1.79
1	B	958	MET	SD-CE	6.10	1.94	1.79
1	B	882	MET	SD-CE	6.07	1.94	1.79
1	A	958	MET	SD-CE	6.00	1.94	1.79
1	A	959	MET	SD-CE	5.99	1.94	1.79
1	B	959	MET	CG-SD	5.74	1.95	1.80
1	B	959	MET	SD-CE	5.72	1.93	1.79
1	B	922	MET	SD-CE	5.72	1.93	1.79
1	A	922	MET	SD-CE	5.70	1.93	1.79
1	B	922	MET	CG-SD	5.39	1.94	1.80
1	A	922	MET	CG-SD	5.24	1.93	1.80
1	A	959	MET	CG-SD	5.10	1.93	1.80

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1034	ARG	NE-CZ-NH2	7.20	125.68	119.20
1	B	852	CYS	CA-C-N	-6.14	113.36	119.92
1	B	852	CYS	C-N-CA	-6.14	113.36	119.92
1	A	1042	ARG	NE-CZ-NH2	5.64	124.27	119.20
1	A	1004	THR	O-C-N	-5.59	116.78	123.27
1	A	1003	TRP	CA-C-N	5.37	131.15	122.29
1	A	1003	TRP	C-N-CA	5.37	131.15	122.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1011	LEU	N-CA-C	5.37	122.23	110.80
1	B	972	ASP	CA-CB-CG	5.12	117.72	112.60
1	A	977	PRO	N-CA-C	5.07	116.89	110.70
1	B	971	SER	CA-C-N	5.06	131.45	121.58
1	B	971	SER	C-N-CA	5.06	131.45	121.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1598	0	1543	14	1
1	B	1503	0	1423	8	1
2	A	6	0	8	1	0
2	B	18	0	24	0	0
3	A	145	0	0	1	0
3	B	158	0	0	0	0
All	All	3428	0	2998	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1052:GLU:OE2	3:A:1201:HOH:O	2.02	0.77
1:A:850:ILE:HG13	1:A:851:PRO:CD	2.17	0.74
1:A:850:ILE:HG13	1:A:851:PRO:HD3	1.70	0.73
1:A:954:PRO:HG3	1:B:958:MET:HE1	1.72	0.72
1:A:906:VAL:HG12	1:A:961:ASN:ND2	2.04	0.71
1:A:1011:LEU:HD23	1:A:1013:ASN:H	1.60	0.66
1:A:1004:THR:CG2	1:A:1015:ARG:HD2	2.26	0.66
1:A:850:ILE:HD12	2:A:1101:GOL:H2	1.81	0.63
1:A:1011:LEU:HD23	1:A:1011:LEU:C	2.31	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:954:PRO:HG3	1:B:958:MET:CE	2.36	0.56
1:B:885:ILE:HG12	1:B:959:MET:CE	2.38	0.53
1:A:1004:THR:HG23	1:A:1015:ARG:HD2	1.92	0.50
1:A:850:ILE:HG13	1:A:851:PRO:HD2	1.90	0.50
1:A:1011:LEU:CD2	1:A:1013:ASN:H	2.23	0.50
1:B:1034:ARG:HH21	1:B:1038:ARG:HH11	1.61	0.47
1:B:882:MET:SD	1:B:953:LEU:HD11	2.54	0.47
1:B:852:CYS:HB3	1:B:853:PRO:HD3	1.96	0.47
1:B:1013:ASN:CG	1:B:1014:LYS:H	2.28	0.42
1:B:940:VAL:HG21	1:B:1050:ASN:HB3	2.02	0.41
1:A:906:VAL:HG12	1:A:961:ASN:HD21	1.83	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:852:CYS:SG	1:B:852:CYS:N[2_444]	2.11	0.09

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/215 (95%)	199 (98%)	5 (2%)	0	100	100
1	B	194/215 (90%)	181 (93%)	11 (6%)	2 (1%)	12	5
All	All	398/430 (93%)	380 (96%)	16 (4%)	2 (0%)	24	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1009	GLN
1	B	999	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/193 (86%)	167 (100%)	0	100	100
1	B	156/193 (81%)	156 (100%)	0	100	100
All	All	323/386 (84%)	323 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	979	GLN
1	A	1013	ASN
1	B	919	ASN
1	B	939	HIS
1	B	979	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	1101	-	5,5,5	0.51	0	5,5,5	0.35	0
2	GOL	B	1102	-	5,5,5	0.48	0	5,5,5	0.29	0
2	GOL	B	1101	-	5,5,5	0.50	0	5,5,5	0.40	0
2	GOL	B	1103	-	5,5,5	0.54	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1101	-	-	0/4/4/4	-
2	GOL	B	1102	-	-	1/4/4/4	-
2	GOL	B	1101	-	-	0/4/4/4	-
2	GOL	B	1103	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1103	GOL	C1-C2-C3-O3
2	B	1102	GOL	C1-C2-C3-O3
2	B	1103	GOL	O2-C2-C3-O3
2	B	1103	GOL	O1-C1-C2-O2
2	B	1103	GOL	O1-C1-C2-C3



There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/215 (95%)	0.70	19 (9%) 14 20	21, 32, 54, 71	0
1	B	198/215 (92%)	0.64	28 (14%) 6 9	20, 29, 54, 62	0
All	All	404/430 (93%)	0.67	47 (11%) 9 13	20, 31, 54, 71	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1011	LEU	7.9
1	B	999	ILE	5.9
1	B	1010	ASP	5.8
1	B	1006	GLY	4.6
1	B	1009	GLN	4.6
1	A	1007	ILE	4.5
1	A	1008	GLN	4.0
1	A	1013	ASN	3.9
1	A	850	ILE	3.5
1	B	1007	ILE	3.4
1	A	1011	LEU	3.3
1	B	1008	GLN	3.3
1	B	997	ASP	3.2
1	B	1013	ASN	3.2
1	A	1044	PHE	3.2
1	B	882	MET	3.1
1	B	998	ARG	3.1
1	B	995	ALA	3.0
1	A	1009	GLN	2.9
1	B	1012	LYS	2.8
1	A	1024	LEU	2.7
1	B	1014	LYS	2.7
1	A	944	ALA	2.7
1	B	1032	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	947	ASP	2.6
1	B	852	CYS	2.6
1	B	1029	THR	2.6
1	A	898	ASN	2.6
1	B	1015	ARG	2.5
1	B	959	MET	2.5
1	B	1000	LYS	2.5
1	A	1004	THR	2.5
1	A	1005	TYR	2.4
1	B	1054	SER	2.3
1	B	994	TRP	2.3
1	A	1003	TRP	2.2
1	A	951	VAL	2.2
1	A	1055	ASP	2.2
1	B	1017	PRO	2.2
1	B	1033	ASN	2.1
1	B	996	LYS	2.1
1	A	959	MET	2.1
1	B	958	MET	2.1
1	A	940	VAL	2.1
1	B	1026	ASP	2.1
1	B	887	LEU	2.0
1	A	941	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	1101	6/6	0.63	0.23	40,45,51,52	0
2	GOL	B	1102	6/6	0.71	0.17	41,46,52,53	0
2	GOL	B	1103	6/6	0.77	0.15	32,37,55,55	0
2	GOL	B	1101	6/6	0.84	0.13	36,36,50,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GOL A 1101:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GOL B 1102:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GOL B 1103:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.