



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 02:21 PM UTC

PDB ID : 1LAT / pdb_00001lat
Title : GLUCOCORTICOID RECEPTOR MUTANT/DNA COMPLEX
Authors : Gewirth, D.T.; Sigler, P.B.
Deposited on : 1995-12-18
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	19	79% 21%
1	D	19	58% 37% 5%
2	A	82	% 57% 23% 5% 13%
2	B	82	4% 60% 22% 7% 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*TP*CP*CP*AP*GP*AP*AP*CP*AP*TP*GP*TP*TP*CP*TP*G P*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	19	Total	C	N	O	P	0	0	0
			386	186	69	113	18			
1	D	19	Total	C	N	O	P	0	0	0
			386	186	69	113	18			

- Molecule 2 is a protein called GLUCOCORTICOID RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	71	Total	C	N	O	S	0	0	0
			557	344	105	97	11			
2	B	74	Total	C	N	O	S	0	0	0
			578	356	111	100	11			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	434	MET	-	expression tag	UNP P06536
A	435	LYS	-	expression tag	UNP P06536
A	436	PRO	-	expression tag	UNP P06536
A	437	ALA	-	expression tag	UNP P06536
A	438	ARG	-	expression tag	UNP P06536
A	439	PRO	-	expression tag	UNP P06536
A	458	GLU	GLY	engineered mutation	UNP P06536
A	459	GLY	SER	engineered mutation	UNP P06536
A	462	ALA	VAL	engineered mutation	UNP P06536
A	477	LYS	ALA	engineered mutation	UNP P06536
A	478	TYR	GLY	engineered mutation	UNP P06536
A	479	GLU	ARG	engineered mutation	UNP P06536
A	480	GLY	ASN	engineered mutation	UNP P06536
A	481	LYS	ASP	engineered mutation	UNP P06536
B	434	MET	-	expression tag	UNP P06536

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	435	LYS	-	expression tag	UNP P06536
B	436	PRO	-	expression tag	UNP P06536
B	437	ALA	-	expression tag	UNP P06536
B	438	ARG	-	expression tag	UNP P06536
B	439	PRO	-	expression tag	UNP P06536
B	458	GLU	GLY	engineered mutation	UNP P06536
B	459	GLY	SER	engineered mutation	UNP P06536
B	462	ALA	VAL	engineered mutation	UNP P06536
B	477	LYS	ALA	engineered mutation	UNP P06536
B	478	TYR	GLY	engineered mutation	UNP P06536
B	479	GLU	ARG	engineered mutation	UNP P06536
B	480	GLY	ASN	engineered mutation	UNP P06536
B	481	LYS	ASP	engineered mutation	UNP P06536

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	82	Total	O	0	0
			82	82		
4	D	74	Total	O	0	0
			74	74		
4	A	72	Total	O	0	0
			72	72		
4	B	88	Total	O	0	0
			88	88		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*TP*TP*CP*CP*AP*GP*AP*AP*CP*AP*TP*GP*TP*TP*CP*TP*G P*GP*A)-3')

Chain C: 



- Molecule 1: DNA (5'-D(*TP*TP*CP*CP*AP*GP*AP*AP*CP*AP*TP*GP*TP*TP*CP*TP*G P*GP*A)-3')

Chain D: 




- Molecule 2: GLUCOCORTICOID RECEPTOR

Chain A: 



- Molecule 2: GLUCOCORTICOID RECEPTOR

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.72Å 76.07Å 118.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.90 6.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.6 (6.00-1.90) 93.3 (6.00-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.89Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.195 , 0.281 0.222 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 157.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2227	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	1.18	0/432	1.23	1/665 (0.2%)
1	D	1.27	1/432 (0.2%)	1.23	0/665
2	A	0.68	0/566	1.23	6/754 (0.8%)
2	B	0.76	0/587	1.10	5/782 (0.6%)
All	All	0.97	1/2017 (0.0%)	1.20	12/2866 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3
1	D	0	2
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	6	DG	O3'-P	-5.24	1.53	1.61

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	438	ARG	CA-C-N	-7.58	112.40	120.66
2	B	438	ARG	C-N-CA	-7.58	112.40	120.66
2	A	492	CYS	CA-C-N	7.37	127.22	119.05
2	A	492	CYS	C-N-CA	7.37	127.22	119.05
2	B	445	ASP	N-CA-C	-6.29	102.67	110.41
2	A	445	ASP	N-CA-C	-6.06	102.67	110.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	472	HIS	N-CA-C	-6.03	97.96	110.80
2	B	442	VAL	N-CA-C	5.91	116.94	111.45
2	A	475	LEU	N-CA-C	-5.90	98.23	110.80
2	A	448	SER	N-CA-C	-5.65	106.55	113.50
2	B	469	GLU	N-CA-C	5.42	120.00	112.90
1	C	2	DT	C4-C5-C7	-5.37	114.34	122.40

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	13	DT	Sidechain
1	C	19	DA	Sidechain
1	C	6	DG	Sidechain
1	D	13	DT	Sidechain
1	D	3	DC	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	386	0	216	0	0
1	D	386	0	217	7	0
2	A	557	0	548	27	0
2	B	578	0	571	19	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	72	0	0	1	0
4	B	88	0	0	4	0
4	C	82	0	0	1	0
4	D	74	0	0	1	0
All	All	2227	0	1552	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:438:ARG:HB3	2:B:439:PRO:CD	2.03	0.89
2:A:478:TYR:HE1	2:A:481:LYS:HB3	1.37	0.88
2:B:437:ALA:HB3	2:B:455:LEU:HD21	1.61	0.83
2:B:475:LEU:HD13	4:B:289:HOH:O	1.79	0.82
2:A:474:TYR:HE1	2:A:493:PRO:HB3	1.54	0.71
2:B:506:ASN:HB3	4:B:157:HOH:O	1.90	0.71
2:B:438:ARG:NE	2:B:438:ARG:HA	2.06	0.70
1:D:4:DC:H2''	1:D:5:DA:N7	2.13	0.64
2:A:476:CYS:SG	2:A:479:GLU:N	2.71	0.63
2:B:438:ARG:HB3	2:B:439:PRO:HD3	1.82	0.62
2:A:470:GLY:O	2:A:471:GLN:HB2	1.99	0.61
2:A:498:ARG:HD2	4:A:312:HOH:O	1.99	0.61
2:B:438:ARG:HB3	2:B:439:PRO:HD2	1.84	0.60
2:A:473:ASN:O	2:A:474:TYR:CD1	2.56	0.58
2:B:441:LEU:HG	4:B:155:HOH:O	2.04	0.57
1:D:10:DA:OP1	2:A:472:HIS:CD2	2.58	0.56
1:D:4:DC:H2''	1:D:5:DA:C8	2.42	0.55
2:A:468:VAL:HG21	2:A:507:LEU:HD23	1.89	0.54
2:B:468:VAL:HG21	2:B:507:LEU:HD22	1.90	0.54
2:B:506:ASN:HD21	2:B:508:GLU:HB2	1.73	0.53
2:A:468:VAL:HG21	2:A:507:LEU:CD2	2.40	0.52
2:A:478:TYR:CE1	2:A:481:LYS:HB3	2.29	0.52
2:B:498:ARG:HG2	2:B:498:ARG:HH11	1.73	0.52
2:B:485:ASP:OD1	2:B:488:ARG:HB2	2.10	0.51
2:A:481:LYS:HG2	2:A:481:LYS:O	2.11	0.49
1:D:12:DG:OP1	2:A:490:LYS:HE3	2.12	0.49
1:D:5:DA:H2''	1:D:6:DG:O5'	2.13	0.48
2:A:472:HIS:HB3	2:A:474:TYR:OH	2.14	0.47
2:A:472:HIS:HD2	2:A:473:ASN:N	2.11	0.47
2:B:451:HIS:CD2	4:B:338:HOH:O	2.67	0.47
2:A:473:ASN:O	2:A:473:ASN:CG	2.57	0.47
2:B:438:ARG:HA	2:B:438:ARG:HE	1.78	0.47
2:A:474:TYR:HE1	2:A:493:PRO:CB	2.26	0.46
1:D:19:DA:H2''	4:D:48:HOH:O	2.15	0.46
2:A:472:HIS:O	2:A:473:ASN:C	2.57	0.46
2:A:480:GLY:HA2	2:A:498:ARG:NH1	2.30	0.46
2:B:449:GLY:O	2:B:455:LEU:HD12	2.16	0.46
2:A:477:LYS:N	2:A:477:LYS:HD2	2.31	0.46
2:B:443:CYS:O	2:B:485:ASP:HA	2.16	0.46
1:D:4:DC:H5''	2:B:450:CYS:O	2.16	0.45
2:B:498:ARG:HH11	2:B:502:GLN:NE2	2.14	0.45
2:A:472:HIS:CD2	2:A:472:HIS:C	2.95	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:79:HOH:O	2:A:465:LYS:HE3	2.17	0.44
2:A:477:LYS:N	2:A:477:LYS:CD	2.80	0.44
2:A:472:HIS:HB3	2:A:474:TYR:CZ	2.52	0.44
2:A:480:GLY:HA2	2:A:498:ARG:HH12	1.83	0.43
2:A:498:ARG:O	2:A:502:GLN:HG3	2.20	0.42
2:A:472:HIS:CD2	2:A:473:ASN:N	2.88	0.41
2:A:478:TYR:CD1	2:A:478:TYR:C	2.98	0.41
2:B:479:GLU:HG3	2:B:481:LYS:HG2	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	69/82 (84%)	59 (86%)	7 (10%)	3 (4%)	2	0
2	B	72/82 (88%)	63 (88%)	7 (10%)	2 (3%)	4	0
All	All	141/164 (86%)	122 (86%)	14 (10%)	5 (4%)	3	0

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	471	GLN
2	A	472	HIS
2	A	480	GLY
2	B	438	ARG
2	B	503	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	60/69 (87%)	55 (92%)	5 (8%)	10	4
2	B	61/69 (88%)	54 (88%)	7 (12%)	5	2
All	All	121/138 (88%)	109 (90%)	12 (10%)	7	3

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	446	GLU
2	A	472	HIS
2	A	473	ASN
2	A	477	LYS
2	A	488	ARG
2	B	441	LEU
2	B	468	VAL
2	B	475	LEU
2	B	490	LYS
2	B	502	GLN
2	B	506	ASN
2	B	507	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	491	ASN
2	B	502	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	19/19 (100%)	-0.87	0 100 100	22, 35, 50, 67	0
1	D	19/19 (100%)	-0.80	0 100 100	23, 35, 51, 51	0
2	A	71/82 (86%)	-0.10	1 (1%) 73 76	26, 43, 76, 85	0
2	B	74/82 (90%)	-0.25	3 (4%) 41 44	24, 40, 69, 82	0
All	All	183/202 (90%)	-0.31	4 (2%) 62 66	22, 40, 72, 85	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	472	HIS	3.2
2	B	510	ARG	2.5
2	B	509	ALA	2.4
2	B	437	ALA	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	1515	1/1	0.91	0.04	50,50,50,50	0
3	ZN	A	1514	1/1	0.99	0.03	34,34,34,34	0
3	ZN	B	1514	1/1	0.99	0.01	33,33,33,33	0
3	ZN	B	1515	1/1	0.99	0.01	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.