



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:14 AM UTC

PDB ID : 1LAW / pdb_00001law
Title : STABILIZATION OF ESCHERICHIA COLI RIBONUCLEASE HI BY
CAVITY-FILLING MUTATIONS WITHIN A HYDROPHOBIC CORE
Authors : Ishikawa, K.; Nakamura, H.; Morikawa, K.; Kanaya, S.
Deposited on : 1993-05-10
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	155	 61% 32% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	155	Total	C	N	O	S	0	0	0
			1239	777	227	228	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	ILE	VAL	conflict	UNP P0A7Y4

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	111	Total	O	0	0
			111	111		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: RIBONUCLEASE H

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.06Å 86.84Å 35.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.188 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1350	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.25	1/1267 (0.1%)	2.07	38/1711 (2.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	40	THR	CA-CB	5.87	1.62	1.53

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	138	ARG	CD-NE-CZ	11.68	140.75	124.40
1	A	153	VAL	CA-C-N	10.45	141.50	121.54
1	A	153	VAL	C-N-CA	10.45	141.50	121.54
1	A	76	GLN	OE1-CD-NE2	7.61	130.21	122.60
1	A	29	ARG	CD-NE-CZ	7.46	134.84	124.40
1	A	94	ASP	CA-C-N	7.29	132.97	122.40
1	A	94	ASP	C-N-CA	7.29	132.97	122.40
1	A	114	HIS	N-CA-C	7.09	119.70	109.14
1	A	51	ALA	O-C-N	7.07	129.35	122.07
1	A	127	HIS	CA-CB-CG	-6.96	106.84	113.80
1	A	109	ALA	CA-C-N	6.74	130.55	120.31
1	A	109	ALA	C-N-CA	6.74	130.55	120.31
1	A	143	ASN	O-C-N	6.73	126.10	121.19
1	A	46	ARG	NE-CZ-NH1	6.68	128.18	121.50
1	A	62	HIS	CA-CB-CG	-6.64	107.16	113.80
1	A	116	ILE	N-CA-C	6.48	117.25	108.17
1	A	105	GLN	CA-C-O	6.40	127.20	120.42
1	A	81	TRP	N-CA-C	6.35	121.13	112.68
1	A	148	ASP	O-C-N	6.27	130.28	122.63
1	A	138	ARG	CG-CD-NE	6.13	125.50	112.00
1	A	46	ARG	CD-NE-CZ	6.05	132.88	124.40
1	A	115	GLN	N-CA-CB	6.04	120.27	110.42
1	A	154	GLU	N-CA-C	5.59	122.71	110.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	44	ASN	CA-CB-CG	5.58	118.18	112.60
1	A	5	VAL	CA-C-O	-5.58	114.47	121.11
1	A	14	LEU	CB-CA-C	5.52	119.25	109.65
1	A	21	GLY	O-C-N	5.49	129.81	123.44
1	A	106	ARG	CD-NE-CZ	-5.45	116.76	124.40
1	A	42	THR	N-CA-CB	-5.44	102.48	110.97
1	A	96	LYS	N-CA-C	-5.42	103.29	110.40
1	A	54	VAL	N-CA-CB	-5.40	103.72	110.47
1	A	115	GLN	CA-C-O	-5.34	114.65	120.36
1	A	50	MET	CA-C-N	5.18	127.17	120.44
1	A	50	MET	C-N-CA	5.18	127.17	120.44
1	A	143	ASN	CA-CB-CG	5.16	117.76	112.60
1	A	72	GLN	CB-CG-CD	5.10	121.27	112.60
1	A	40	THR	N-CA-C	5.04	116.77	111.28
1	A	66	ILE	N-CA-CB	5.02	118.87	112.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1239	0	1219	35	0
2	A	111	0	0	6	0
All	All	1350	0	1219	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:CYS:H	1:A:114:HIS:HD2	1.07	0.95
1:A:152:GLN:HB3	1:A:154:GLU:HB2	1.59	0.83
1:A:152:GLN:HE22	1:A:155:VAL:HB	1.43	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:CYS:H	1:A:114:HIS:CD2	2.00	0.69
1:A:127:HIS:HB3	1:A:128:PRO:HD2	1.76	0.67
1:A:111:LEU:CD2	1:A:116:ILE:HD12	2.35	0.57
1:A:63:CYS:N	1:A:114:HIS:HD2	1.91	0.56
1:A:47:MET:HA	1:A:47:MET:HE2	1.88	0.55
1:A:77:GLY:HA3	1:A:104:TRP:CH2	2.42	0.55
1:A:153:VAL:HG23	1:A:154:GLU:OE1	2.08	0.54
1:A:152:GLN:NE2	1:A:155:VAL:HB	2.18	0.53
1:A:111:LEU:HD23	1:A:116:ILE:HD12	1.91	0.53
1:A:153:VAL:HG23	1:A:154:GLU:CD	2.34	0.53
1:A:149:THR:HG23	2:A:319:HOH:O	2.08	0.52
1:A:64:GLU:HB3	2:A:335:HOH:O	2.09	0.51
1:A:153:VAL:HG23	1:A:154:GLU:OE2	2.10	0.51
1:A:49:LEU:O	1:A:53:ILE:HG13	2.11	0.50
1:A:9:THR:HB	1:A:51:ALA:HB1	1.94	0.50
1:A:125:ALA:HA	2:A:304:HOH:O	2.12	0.49
1:A:127:HIS:CB	1:A:128:PRO:HD2	2.43	0.48
1:A:128:PRO:HA	1:A:131:GLU:OE2	2.13	0.48
1:A:91:LYS:HE2	2:A:311:HOH:O	2.15	0.46
1:A:9:THR:HB	1:A:51:ALA:CB	2.45	0.46
1:A:138:ARG:NH1	2:A:252:HOH:O	2.50	0.45
1:A:142:MET:HE3	1:A:142:MET:HB3	1.73	0.45
1:A:106:ARG:NH1	2:A:214:HOH:O	2.36	0.45
1:A:102:ASP:OD2	1:A:150:GLY:HA3	2.16	0.44
1:A:60:LYS:HD2	1:A:60:LYS:HA	1.72	0.44
1:A:65:VAL:HG12	1:A:66:ILE:N	2.33	0.43
1:A:68:SER:HA	1:A:119:GLU:O	2.18	0.43
1:A:149:THR:O	1:A:151:TYR:N	2.50	0.43
1:A:22:TYR:CD1	1:A:22:TYR:C	2.96	0.43
1:A:67:LEU:HD22	1:A:118:TRP:CZ3	2.54	0.42
1:A:41:ARG:HH11	1:A:41:ARG:HD3	1.55	0.40
1:A:92:THR:OG1	1:A:94:ASP:OD1	2.31	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	153/155 (99%)	144 (94%)	7 (5%)	2 (1%)	9 3

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	GLU
1	A	150	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	127/127 (100%)	120 (94%)	7 (6%)	19 8

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	66	ILE
1	A	99	LYS
1	A	116	ILE
1	A	149	THR
1	A	153	VAL
1	A	154	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	76	GLN
1	A	80	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	105	GLN
1	A	113	GLN
1	A	114	HIS
1	A	152	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.