



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:24 AM UTC

PDB ID : 5LA9 / pdb_00005la9
Title : HIF prolyl hydroxylase 2 (PHD2-R281C/V314C) cross-linked to HIF-1alpha
NODD-L397C/D412C and N-oxalylglycine (NOG) (complex-2)
Authors : Chowdhury, R.; Schofield, C.J.
Deposited on : 2016-06-14
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	
1	B	252	
2	C	19	
2	D	19	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6745 atoms, of which 3215 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Egl nine homolog 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	205	Total	C	H	N	O	S	0	0	0
			2983	982	1429	267	293	12			
1	B	212	Total	C	H	N	O	S	0	0	0
			3140	1024	1525	280	299	12			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	175	GLY	-	expression tag	UNP Q9GZT9
A	176	SER	-	expression tag	UNP Q9GZT9
A	177	HIS	-	expression tag	UNP Q9GZT9
A	178	MET	-	expression tag	UNP Q9GZT9
A	179	ALA	-	expression tag	UNP Q9GZT9
A	180	SER	-	expression tag	UNP Q9GZT9
A	201	ALA	CYS	engineered mutation	UNP Q9GZT9
A	281	CYS	ARG	engineered mutation	UNP Q9GZT9
A	314	CYS	VAL	engineered mutation	UNP Q9GZT9
A	398	ALA	ARG	engineered mutation	UNP Q9GZT9
B	175	GLY	-	expression tag	UNP Q9GZT9
B	176	SER	-	expression tag	UNP Q9GZT9
B	177	HIS	-	expression tag	UNP Q9GZT9
B	178	MET	-	expression tag	UNP Q9GZT9
B	179	ALA	-	expression tag	UNP Q9GZT9
B	180	SER	-	expression tag	UNP Q9GZT9
B	201	ALA	CYS	engineered mutation	UNP Q9GZT9
B	281	CYS	ARG	engineered mutation	UNP Q9GZT9
B	314	CYS	VAL	engineered mutation	UNP Q9GZT9
B	398	ALA	ARG	engineered mutation	UNP Q9GZT9

- Molecule 2 is a protein called Hypoxia-inducible factor 1-alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	19	Total	C	H	N	O	S	0	0	0
			257	83	127	19	26	2			
2	D	19	Total	C	H	N	O	S	0	0	0
			258	83	128	19	26	2			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	397	CYS	LEU	engineered mutation	UNP Q16665
C	412	CYS	ASP	engineered mutation	UNP Q16665
D	397	CYS	LEU	engineered mutation	UNP Q16665
D	412	CYS	ASP	engineered mutation	UNP Q16665

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	B	1	Total	Mn	0	0
			1	1		

- Molecule 4 is N-OXALYLGLYCINE (CCD ID: OGA) (formula: C₄H₅NO₅).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	0	0
			13	4	3	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	H	N	O	0	0
			13	4	3	1	5		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	30	Total	O	0	0
			30	30		
6	B	37	Total	O	0	0
			37	37		
6	C	3	Total	O	0	0
			3	3		
6	D	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Egl nine homolog 1




- Molecule 1: Egl nine homolog 1



- Molecule 2: Hypoxia-inducible factor 1-alpha



- Molecule 2: Hypoxia-inducible factor 1-alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.47Å 97.33Å 71.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.81 48.13 – 2.81	Depositor EDS
% Data completeness (in resolution range)	97.7 (48.13-2.81) 97.7 (48.13-2.81)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.251 , 0.279 0.260 , 0.280	Depositor DCC
R_{free} test set	719 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	6745	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OGA, MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/1591	0.37	0/2164
1	B	0.14	0/1652	0.35	0/2240
2	C	0.17	0/131	0.47	0/179
2	D	0.11	0/131	0.40	0/179
All	All	0.14	0/3505	0.36	0/4762

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1554	1429	1424	23	0
1	B	1615	1525	1525	19	0
2	C	130	127	128	4	0
2	D	130	128	128	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	3	3	0	0
4	B	10	3	3	0	0
5	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	30	0	0	5	0
6	B	37	0	0	2	0
6	C	3	0	0	0	0
6	D	4	0	0	0	0
All	All	3530	3215	3211	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:ALA:N	6:A:602:HOH:O	2.21	0.72
1:B:293:ASN:ND2	6:B:603:HOH:O	2.25	0.67
1:B:272:LEU:HG	1:B:276:MET:HE3	1.84	0.60
1:A:299:MET:SD	6:A:619:HOH:O	2.57	0.58
1:A:326:CYS:HG	1:A:390:TYR:HE2	1.54	0.53
1:A:194:ALA:HB1	1:A:283:CYS:SG	2.50	0.52
1:A:330:LEU:HD11	1:A:384:TYR:HB2	1.92	0.52
1:B:235:PHE:HB3	1:B:255:LYS:HB3	1.91	0.51
1:A:199:VAL:HB	1:A:200:PRO:HD3	1.93	0.51
1:B:229:LEU:O	1:B:232:THR:HG22	2.10	0.50
1:A:330:LEU:HD11	1:A:384:TYR:CB	2.42	0.50
2:C:406:ASP:OD1	2:C:406:ASP:N	2.29	0.49
1:B:251:ILE:HG23	1:B:311:VAL:HG22	1.94	0.48
1:A:377:GLN:OE1	1:B:344:ARG:NH1	2.46	0.48
1:B:370:ARG:HG2	2:D:397:CYS:SG	2.54	0.48
1:A:303:TYR:HB2	1:A:383:ARG:HB3	1.96	0.47
1:B:357:GLU:O	1:B:362:ARG:NH1	2.47	0.47
1:B:303:TYR:HB2	1:B:383:ARG:HB3	1.98	0.46
1:B:232:THR:HG21	1:B:234:LYS:CE	2.46	0.46
1:A:220:GLN:OE1	1:A:332:LYS:NZ	2.50	0.45
1:A:303:TYR:HD1	6:A:612:HOH:O	1.99	0.45
1:B:292:ILE:HA	1:B:391:PHE:O	2.17	0.45
1:B:212:ASP:OD1	6:B:601:HOH:O	2.21	0.45
1:B:254:ASP:OD1	1:B:254:ASP:N	2.50	0.44
1:B:241:VAL:HG11	2:D:401:ALA:HB3	1.99	0.44
1:A:326:CYS:HA	1:A:364:LEU:O	2.18	0.43
1:B:326:CYS:HA	1:B:364:LEU:O	2.19	0.43
1:A:241:VAL:N	2:C:401:ALA:O	2.52	0.42
1:A:315:ASP:N	6:A:601:HOH:O	2.15	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ALA:HB2	1:A:279:LEU:HD22	2.01	0.42
1:A:283:CYS:O	1:A:284:ASN:C	2.62	0.42
1:B:193:LEU:C	1:B:193:LEU:HD23	2.45	0.42
1:A:336:ALA:CB	1:A:342:ILE:HD11	2.49	0.42
1:A:280:ILE:HG13	1:A:390:TYR:CZ	2.54	0.42
1:B:338:VAL:HG13	1:B:339:SER:N	2.36	0.41
1:B:232:THR:HG21	1:B:234:LYS:HE2	2.02	0.41
1:B:251:ILE:N	1:B:251:ILE:HD12	2.36	0.40
1:A:330:LEU:CD1	1:A:384:TYR:HB2	2.51	0.40
1:A:345:ILE:HB	1:A:354:ALA:HB3	2.03	0.40
1:A:370:ARG:HA	2:C:397:CYS:SG	2.61	0.40
1:A:227:ARG:NH1	6:A:610:HOH:O	2.53	0.40
1:A:315:ASP:HA	2:C:402:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/252 (80%)	192 (96%)	9 (4%)	0	100	100
1	B	208/252 (82%)	199 (96%)	9 (4%)	0	100	100
2	C	17/19 (90%)	16 (94%)	1 (6%)	0	100	100
2	D	17/19 (90%)	14 (82%)	3 (18%)	0	100	100
All	All	443/542 (82%)	421 (95%)	22 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/209 (72%)	151 (100%)	0	100	100
1	B	160/209 (77%)	158 (99%)	2 (1%)	61	85
2	C	14/14 (100%)	13 (93%)	1 (7%)	13	38
2	D	14/14 (100%)	14 (100%)	0	100	100
All	All	339/446 (76%)	336 (99%)	3 (1%)	70	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	353	PHE
1	B	357	GLU
2	C	406	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	372	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	OGA	B	502	3	9,9,9	2.44	1 (11%)	8,11,11	1.29	0
4	OGA	A	502	3	9,9,9	2.49	2 (22%)	8,11,11	1.25	0
5	SO4	A	503	-	4,4,4	0.24	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OGA	B	502	3	-	0/8/9/9	-
4	OGA	A	502	3	-	0/8/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	OGA	O2'-C2	6.43	1.35	1.23
4	B	502	OGA	O2'-C2	6.40	1.35	1.23
4	A	502	OGA	C2-C1	-2.06	1.51	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	205/252 (81%)	0.95	18 (8%) 15 11	22, 38, 61, 85	0
1	B	212/252 (84%)	0.92	24 (11%) 10 7	19, 37, 58, 69	0
2	C	19/19 (100%)	1.14	1 (5%) 32 24	39, 44, 52, 55	0
2	D	19/19 (100%)	1.01	2 (10%) 11 8	27, 36, 50, 52	0
All	All	455/542 (83%)	0.95	45 (9%) 13 9	19, 38, 59, 85	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	VAL	6.0
1	A	188	LEU	4.6
1	B	249	LYS	4.4
1	B	241	VAL	4.0
1	A	351	ALA	3.7
1	B	278	ASP	3.7
1	B	266	CYS	3.6
1	A	189	PRO	3.6
1	A	352	GLN	3.4
1	A	251	ILE	3.4
1	B	250	ASP	3.3
1	A	354	ALA	3.2
1	B	188	LEU	3.1
1	A	401	VAL	3.1
1	A	340	GLY	3.0
1	B	191	LEU	2.9
1	A	284	ASN	2.9
1	B	284	ASN	2.8
1	B	293	ASN	2.7
1	B	338	VAL	2.7
1	B	352	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	314	CYS	2.6
1	B	351	ALA	2.6
1	B	190	ALA	2.5
1	A	319	GLY	2.4
1	A	262	LYS	2.4
1	B	406	GLY	2.4
1	A	191	LEU	2.3
1	B	282	HIS	2.3
1	B	370	ARG	2.3
1	B	304	PRO	2.3
1	B	262	LYS	2.2
1	B	353	PHE	2.2
1	B	189	PRO	2.2
1	A	281	CYS	2.2
1	A	357	GLU	2.2
2	D	412	CYS	2.1
1	B	331	ASN	2.1
1	A	288	GLY	2.1
2	D	413	PHE	2.1
2	C	407	THR	2.1
1	B	381	ALA	2.1
1	A	353	PHE	2.0
1	B	253	GLY	2.0
1	A	379	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	A	503	5/5	0.74	0.18	110,111,111,111	0
4	OGA	A	502	10/10	0.92	0.10	20,26,30,33	0
4	OGA	B	502	10/10	0.96	0.12	30,34,36,37	0
3	MN	B	501	1/1	0.99	0.07	17,17,17,17	0
3	MN	A	501	1/1	0.99	0.07	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.