



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:28 PM UTC

PDB ID : 7LA1 / pdb_00007la1
Title : Crystal structure of phosphoglycerate kinase from Mycobacterium avium
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2021-01-05
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	2% 92% 6% .
1	B	423	6% 91% 6% .
1	C	423	3% 94% . .
1	D	423	0% 95% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoglycerate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	412	Total	C	N	O	S	0	18	0
			3068	1941	527	594	6			
1	B	411	Total	C	N	O	S	0	7	0
			2996	1890	513	587	6			
1	C	412	Total	C	N	O	S	0	15	0
			3042	1925	523	588	6			
1	D	409	Total	C	N	O	S	0	12	0
			3004	1896	517	585	6			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP A0QHY4
A	1	VAL	-	expression tag	UNP A0QHY4
A	416	GLY	-	expression tag	UNP A0QHY4
A	417	HIS	-	expression tag	UNP A0QHY4
A	418	HIS	-	expression tag	UNP A0QHY4
A	419	HIS	-	expression tag	UNP A0QHY4
A	420	HIS	-	expression tag	UNP A0QHY4
A	421	HIS	-	expression tag	UNP A0QHY4
A	422	HIS	-	expression tag	UNP A0QHY4
B	0	MET	-	initiating methionine	UNP A0QHY4
B	1	VAL	-	expression tag	UNP A0QHY4
B	416	GLY	-	expression tag	UNP A0QHY4
B	417	HIS	-	expression tag	UNP A0QHY4
B	418	HIS	-	expression tag	UNP A0QHY4
B	419	HIS	-	expression tag	UNP A0QHY4
B	420	HIS	-	expression tag	UNP A0QHY4
B	421	HIS	-	expression tag	UNP A0QHY4
B	422	HIS	-	expression tag	UNP A0QHY4
C	0	MET	-	initiating methionine	UNP A0QHY4
C	1	VAL	-	expression tag	UNP A0QHY4
C	416	GLY	-	expression tag	UNP A0QHY4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	417	HIS	-	expression tag	UNP A0QHY4
C	418	HIS	-	expression tag	UNP A0QHY4
C	419	HIS	-	expression tag	UNP A0QHY4
C	420	HIS	-	expression tag	UNP A0QHY4
C	421	HIS	-	expression tag	UNP A0QHY4
C	422	HIS	-	expression tag	UNP A0QHY4
D	0	MET	-	initiating methionine	UNP A0QHY4
D	1	VAL	-	expression tag	UNP A0QHY4
D	416	GLY	-	expression tag	UNP A0QHY4
D	417	HIS	-	expression tag	UNP A0QHY4
D	418	HIS	-	expression tag	UNP A0QHY4
D	419	HIS	-	expression tag	UNP A0QHY4
D	420	HIS	-	expression tag	UNP A0QHY4
D	421	HIS	-	expression tag	UNP A0QHY4
D	422	HIS	-	expression tag	UNP A0QHY4

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	493	Total O 505 505	0	12
3	B	423	Total O 434 434	0	11
3	C	501	Total O 516 516	0	15
3	D	470	Total O 478 478	0	8

- Molecule 1: Phosphoglycerate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.84Å 111.17Å 99.27Å 90.00° 101.87° 90.00°	Depositor
Resolution (Å)	44.90 – 1.60 44.90 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.90-1.60) 99.7 (44.90-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.19rc4-4035	Depositor
R, R_{free}	0.164 , 0.194 0.164 , 0.194	Depositor DCC
R_{free} test set	2069 reflections (0.95%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14107	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/3156	0.45	0/4292
1	B	0.22	0/3065	0.43	0/4170
1	C	0.24	0/3135	0.44	0/4264
1	D	0.23	0/3088	0.41	0/4201
All	All	0.23	0/12444	0.43	0/16927

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3068	0	3142	17	0
1	B	2996	0	3035	17	0
1	C	3042	0	3112	9	0
1	D	3004	0	3054	7	0
2	A	16	0	24	4	0
2	B	20	0	30	3	0
2	C	24	0	36	1	0
2	D	4	0	6	1	0
3	A	505	0	0	4	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	434	0	0	4	0
3	C	516	0	0	0	1
3	D	478	0	0	4	0
All	All	14107	0	12439	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LEU:HD22	1:A:256:MET:HE1	1.66	0.77
1:A:188:GLU:OE2	3:A:601:HOH:O	2.03	0.77
1:B:69:ARG:H	2:B:504:EDO:H12	1.52	0.74
1:D:220:GLU:OE1	3:D:601:HOH:O	2.10	0.70
1:A:189:ILE:HD11	1:A:402[B]:ILE:HD12	1.74	0.69
1:A:379:GLY:HA2	2:A:504:EDO:H22	1.79	0.64
1:A:282:ARG:NH1	3:A:602:HOH:O	2.25	0.64
1:C:29:LEU:HD11	1:C:67:LEU:HD21	1.82	0.60
1:B:189:ILE:HD11	1:B:402:ILE:HD12	1.82	0.59
1:B:50:THR:OG1	1:B:183[B]:THR:HG22	2.03	0.59
1:B:69:ARG:H	2:B:504:EDO:C1	2.20	0.55
1:B:187[B]:GLU:HG3	3:B:789:HOH:O	2.08	0.54
1:A:354:GLY:HA2	2:A:504:EDO:H11	1.90	0.53
1:B:333:PHE:HB2	1:B:373:LEU:HD11	1.90	0.53
1:C:22[A]:ARG:HG2	1:C:170[A]:VAL:HG22	1.92	0.52
1:A:69[B]:ARG:HD2	1:B:77:ALA:HB2	1.92	0.51
1:A:218:VAL:HG21	1:A:391:LEU:HD22	1.93	0.51
1:B:111:GLU:HG3	3:B:793:HOH:O	2.12	0.50
1:D:29:LEU:HD11	1:D:67:LEU:HD21	1.94	0.50
1:A:194:GLN:O	1:A:199:THR:HB	2.11	0.50
1:A:385[A]:THR:HG21	3:A:620:HOH:O	2.11	0.49
1:D:32:ASP:OD1	3:D:602:HOH:O	2.20	0.46
1:B:49:PRO:HB2	1:B:183[B]:THR:HG21	1.96	0.46
1:B:147:GLU:HG3	3:B:791:HOH:O	2.16	0.46
1:A:125:ARG:O	2:A:503:EDO:H11	2.16	0.46
1:A:200:LYS:HE2	1:A:322:GLU:HG3	1.97	0.46
1:C:44:ILE:O	1:C:48:VAL:HG23	2.16	0.45
1:B:32:ASP:H	2:B:501:EDO:H12	1.80	0.45
1:B:364:GLY:HA2	1:B:385:THR:HA	1.99	0.45
1:B:216:LEU:HD22	1:B:256:MET:HE1	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:261:ARG:NH2	1:D:265:ASP:OD1	2.40	0.44
1:A:379:GLY:CA	2:A:504:EDO:H22	2.45	0.44
1:D:233:GLY:HA2	2:D:501:EDO:H21	1.99	0.44
1:C:160:PHE:O	1:C:163:VAL:HG22	2.17	0.43
1:B:81:ALA:HA	1:B:97:LEU:HD22	2.01	0.43
1:C:212[B]:VAL:HG12	1:C:235:MET:HB2	2.01	0.43
1:B:160:PHE:O	1:B:163:VAL:HG22	2.19	0.42
1:B:183[A]:THR:HG21	3:B:822:HOH:O	2.18	0.42
1:D:262[B]:ARG:HG3	3:D:797:HOH:O	2.20	0.42
1:A:359:SER:O	1:A:381:SER:HB2	2.20	0.42
1:D:183[B]:THR:HG22	3:D:905:HOH:O	2.18	0.42
1:A:206:VAL:HG11	1:A:345[B]:LEU:HD21	2.02	0.41
1:A:160:PHE:O	1:A:163:VAL:HG22	2.21	0.41
1:A:385[A]:THR:HB	3:A:809:HOH:O	2.20	0.41
1:C:24:ASP:OD2	1:C:43:ARG:NH1	2.53	0.41
1:C:74:PRO:HG3	2:C:502:EDO:H11	2.02	0.41
1:C:212[B]:VAL:CG1	1:C:235:MET:HB2	2.51	0.41
1:C:189:ILE:HD11	1:C:402[A]:ILE:HD12	2.03	0.40
1:B:193:GLU:OE2	1:B:393:TYR:OH	2.24	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:984:HOH:O	3:C:860:HOH:O[2_555]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	428/423 (101%)	419 (98%)	9 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	416/423 (98%)	405 (97%)	11 (3%)	0	100	100
1	C	425/423 (100%)	415 (98%)	10 (2%)	0	100	100
1	D	420/423 (99%)	411 (98%)	9 (2%)	0	100	100
All	All	1689/1692 (100%)	1650 (98%)	39 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/312 (103%)	318 (99%)	2 (1%)	78	66
1	B	310/312 (99%)	309 (100%)	1 (0%)	86	78
1	C	317/312 (102%)	316 (100%)	1 (0%)	86	78
1	D	313/312 (100%)	313 (100%)	0	100	100
All	All	1260/1248 (101%)	1256 (100%)	4 (0%)	87	78

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	383[A]	ILE
1	A	383[B]	ILE
1	B	411	THR
1	C	383	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	4	HIS
1	C	4	HIS
1	C	194	GLN
1	D	4	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	94	HIS
1	D	243	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	504	-	3,3,3	0.31	0	2,2,2	0.37	0
2	EDO	C	506	-	3,3,3	0.44	0	2,2,2	0.67	0
2	EDO	A	501	-	3,3,3	0.48	0	2,2,2	0.33	0
2	EDO	B	503	-	3,3,3	0.42	0	2,2,2	0.46	0
2	EDO	B	505	-	3,3,3	0.45	0	2,2,2	0.32	0
2	EDO	A	502	-	3,3,3	0.48	0	2,2,2	0.39	0
2	EDO	C	505	-	3,3,3	0.41	0	2,2,2	0.37	0
2	EDO	B	502	-	3,3,3	0.41	0	2,2,2	0.39	0
2	EDO	A	504	-	3,3,3	0.44	0	2,2,2	0.10	0
2	EDO	C	502	-	3,3,3	0.53	0	2,2,2	0.36	0
2	EDO	C	504	-	3,3,3	0.41	0	2,2,2	0.37	0
2	EDO	C	503	-	3,3,3	0.44	0	2,2,2	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	503	-	3,3,3	0.71	0	2,2,2	0.50	0
2	EDO	C	501	-	3,3,3	0.50	0	2,2,2	0.23	0
2	EDO	D	501	-	3,3,3	0.51	0	2,2,2	0.11	0
2	EDO	B	501	-	3,3,3	0.55	0	2,2,2	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	504	-	-	1/1/1/1	-
2	EDO	C	506	-	-	0/1/1/1	-
2	EDO	A	501	-	-	0/1/1/1	-
2	EDO	B	503	-	-	0/1/1/1	-
2	EDO	B	505	-	-	0/1/1/1	-
2	EDO	A	502	-	-	1/1/1/1	-
2	EDO	C	505	-	-	0/1/1/1	-
2	EDO	B	502	-	-	0/1/1/1	-
2	EDO	A	504	-	-	0/1/1/1	-
2	EDO	C	502	-	-	0/1/1/1	-
2	EDO	C	504	-	-	0/1/1/1	-
2	EDO	C	503	-	-	1/1/1/1	-
2	EDO	A	503	-	-	1/1/1/1	-
2	EDO	C	501	-	-	0/1/1/1	-
2	EDO	D	501	-	-	0/1/1/1	-
2	EDO	B	501	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	503	EDO	O1-C1-C2-O2
2	A	502	EDO	O1-C1-C2-O2
2	C	503	EDO	O1-C1-C2-O2
2	B	504	EDO	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	504	EDO	2	0
2	A	504	EDO	3	0
2	C	502	EDO	1	0
2	A	503	EDO	1	0
2	D	501	EDO	1	0
2	B	501	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	412/423 (97%)	-0.14	10 (2%) 59 63	10, 23, 40, 53	18 (4%)
1	B	411/423 (97%)	0.07	25 (6%) 27 27	12, 26, 51, 72	7 (1%)
1	C	412/423 (97%)	-0.15	11 (2%) 56 59	9, 23, 42, 69	15 (3%)
1	D	409/423 (96%)	-0.00	6 (1%) 72 76	10, 25, 48, 67	12 (2%)
All	All	1644/1692 (97%)	-0.06	52 (3%) 50 53	9, 24, 46, 72	52 (3%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	385[A]	THR	5.5
1	A	386[A]	GLY	4.9
1	B	368	ALA	4.8
1	C	0	MET	4.6
1	B	369	ALA	4.4
1	B	385	THR	4.2
1	B	2	ALA	4.0
1	B	371	ARG	3.9
1	B	1	VAL	3.8
1	B	364	GLY	3.8
1	B	375	ILE	3.8
1	D	198	SER	3.7
1	B	372	ALA	3.7
1	D	2	ALA	3.4
1	C	411	THR	3.3
1	C	1	VAL	3.3
1	A	383[A]	ILE	3.0
1	D	199	THR	3.0
1	B	284	ALA	2.9
1	B	373	LEU	2.9
1	B	285	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	198	SER	2.9
1	B	287	ALA	2.8
1	D	197	GLY	2.8
1	B	370	VAL	2.8
1	B	411	THR	2.6
1	A	197	GLY	2.6
1	B	386	GLY	2.6
1	C	33	GLY	2.6
1	A	246	SER	2.6
1	B	367	ALA	2.6
1	B	374	GLY	2.5
1	C	39	THR	2.5
1	C	196	THR	2.5
1	B	334	GLU	2.4
1	A	296	ALA	2.4
1	B	365	ASP	2.4
1	D	398	ALA	2.3
1	C	386	GLY	2.3
1	C	41	PRO	2.3
1	D	410	PRO	2.2
1	A	384[A]	SER	2.2
1	B	366	SER	2.2
1	C	38	ILE	2.2
1	B	3	VAL	2.2
1	C	365	ASP	2.2
1	A	0	MET	2.2
1	C	36	GLY	2.1
1	B	283	PHE	2.1
1	B	269	ASP	2.1
1	B	333	PHE	2.1
1	A	299	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	B	505	4/4	0.78	0.18	46,48,50,52	0
2	EDO	C	506	4/4	0.78	0.16	33,36,36,39	0
2	EDO	A	504	4/4	0.83	0.12	34,39,44,52	0
2	EDO	C	504	4/4	0.83	0.12	31,39,44,48	0
2	EDO	C	505	4/4	0.83	0.13	45,50,52,56	0
2	EDO	B	501	4/4	0.83	0.14	28,31,38,45	0
2	EDO	B	503	4/4	0.84	0.12	44,49,53,53	0
2	EDO	C	503	4/4	0.86	0.12	32,44,45,45	0
2	EDO	A	503	4/4	0.86	0.24	18,26,32,36	0
2	EDO	C	502	4/4	0.88	0.16	26,29,30,36	0
2	EDO	A	502	4/4	0.90	0.11	31,33,37,49	0
2	EDO	D	501	4/4	0.90	0.12	31,35,35,37	0
2	EDO	B	504	4/4	0.92	0.22	24,30,30,33	0
2	EDO	A	501	4/4	0.94	0.08	24,27,35,39	0
2	EDO	B	502	4/4	0.96	0.08	30,31,34,41	0
2	EDO	C	501	4/4	0.99	0.04	14,16,17,19	0

6.5 Other polymers

There are no such residues in this entry.