



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 07:39 PM UTC

PDB ID : 6LBR / pdb_00006lbr
Title : Crystal structure of yeast Cdc13 and ssDNA
Authors : Ge, Y.; Wu, Z.; Wu, J.; Lei, M.
Deposited on : 2019-11-14
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	
1	B	551	
2	C	25	
2	D	25	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KLLA0F20922p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	519	Total	C	N	O	S	0	0	0
			4190	2702	687	788	13			
1	B	208	Total	C	N	O	S	0	0	0
			1671	1078	280	307	6			

- Molecule 2 is a DNA chain called Telomere single-strand DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	22	Total	C	N	O	P	0	0	0
			460	219	84	135	22			
2	D	23	Total	C	N	O	P	0	0	0
			480	229	86	142	23			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	148	Total	O	0	0
			148	148		
3	B	35	Total	O	0	0
			35	35		
3	C	24	Total	O	0	0
			24	24		
3	D	20	Total	O	0	0
			20	20		

- Molecule 1: KLLA0F20922p




TYR	ASN
LYS	LEU
ARG	SER
GLU	ASP
ILE	ASP
LEU	ASP
THR	
ALA	
CYS	
ILE	
GLY	
SER	
ILE	
VAL	
ASP	
PHE	
THR	
LEU	
ILE	
PRO	
ARG	
VAL	
LEU	
ARG	
VAL	
ASN	
GLU	
TYR	
LEU	
TYR	
ILE	
LYS	
ILE	
TRP	
CYS	
PRO	
ILE	
TYR	
ALA	
THR	
LEU	
GLU	
SER	
LEU	
LEU	
ILE	
HIS	
SER	
ARG	
LEU	
GLU	
TYR	
ASN	
ASP	
ASP	
THR	
LEU	
LYS	
TYR	
GLU	

● Molecule 2: Telomere single-strand DNA



● Molecule 2: Telomere single-strand DNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.17Å 106.95Å 173.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.84 – 2.50 45.84 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.8 (45.84-2.50) 93.8 (45.84-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.15_3459	Depositor
R, R_{free}	0.202 , 0.252 0.203 , 0.252	Depositor DCC
R_{free} test set	2219 reflections (4.50%)	wwPDB-VP
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7028	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/4278	0.78	9/5802 (0.2%)
1	B	0.36	0/1706	0.61	2/2310 (0.1%)
2	C	0.36	0/516	0.58	0/797
2	D	0.38	0/538	0.59	0/831
All	All	0.35	0/7038	0.71	11/9740 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	702	ARG	N-CA-C	30.85	151.10	112.23
1	A	703	MET	CB-CA-C	13.12	134.92	110.11
1	A	704	SER	N-CA-CB	-10.92	93.40	110.44
1	A	704	SER	N-CA-C	10.66	126.44	113.23
1	A	703	MET	N-CA-C	-9.88	98.71	112.45

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	617	CYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4190	0	4252	53	0
1	B	1671	0	1719	23	0
2	C	460	0	251	1	0
2	D	480	0	263	6	0
3	A	148	0	0	2	0
3	B	35	0	0	0	0
3	C	24	0	0	0	0
3	D	20	0	0	0	0
All	All	7028	0	6485	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 76 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:656:THR:HG22	1:B:658:THR:H	1.31	0.94
1:A:759:THR:HG21	1:A:779:LYS:HA	1.64	0.78
1:A:562:ASP:HB3	1:A:564:THR:HB	1.68	0.76
1:A:656:THR:HG22	1:A:658:THR:H	1.51	0.74
1:A:860:ILE:HG23	1:A:878:ILE:HD11	1.70	0.72

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	515/551 (94%)	497 (96%)	18 (4%)	0	100	100
1	B	204/551 (37%)	198 (97%)	6 (3%)	0	100	100
All	All	719/1102 (65%)	695 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	477/507 (94%)	446 (94%)	31 (6%)	15	32
1	B	192/507 (38%)	184 (96%)	8 (4%)	26	52
All	All	669/1014 (66%)	630 (94%)	39 (6%)	18	38

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	847	LEU
1	B	612	MET
1	A	860	ILE
1	B	564	THR
1	B	650	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	810	GLN
1	B	674	ASN
1	B	686	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	519/551 (94%)	0.10	44 (8%) 16 14	17, 34, 81, 143	0
1	B	208/551 (37%)	0.27	11 (5%) 32 28	25, 40, 73, 107	0
2	C	22/25 (88%)	0.21	3 (13%) 7 5	30, 47, 100, 112	0
2	D	23/25 (92%)	0.46	3 (13%) 7 6	32, 60, 89, 120	0
All	All	772/1152 (67%)	0.16	61 (7%) 18 16	17, 37, 80, 143	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	756	LEU	8.1
1	B	707	PRO	4.9
1	A	741	GLY	4.8
1	A	808	GLY	4.7
1	A	369	GLY	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.