



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:35 PM UTC

PDB ID : 1LCV / pdb_00001lcv
Title : streptavidin-norbiotin complex
Authors : Livnah, O.; Pazy, Y.
Deposited on : 2002-04-07
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	121	
1	B	121	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called streptavidin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	121	Total	C	N	O	0	0	0
			904	565	157	182			
1	B	118	Total	C	N	O	0	0	0
			882	551	153	178			

- Molecule 2 is NORBIOTIN (CCD ID: SNR) (formula: $C_9H_{14}N_2O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	9	2	3	1		
2	B	1	Total	C	N	O	S	0	0
			15	9	2	3	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	45	Total 45	O 45	0	0
3	B	49	Total 49	O 49	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: streptavidin

Chain A:  78% 21% .



- Molecule 1: streptavidin

Chain B:  79% 16% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	47.29Å 95.32Å 105.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.188 , 0.233	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1910	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SNR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/927	0.81	2/1268 (0.2%)
1	B	0.41	0/904	0.86	3/1238 (0.2%)
All	All	0.40	0/1831	0.84	5/2506 (0.2%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	123	THR	N-CA-C	6.20	118.52	108.41
1	B	231	VAL	N-CA-C	5.68	117.60	108.85
1	B	323	THR	N-CA-C	5.42	117.23	108.34
1	B	231	VAL	CB-CA-C	-5.15	102.88	110.82
1	A	108	TRP	N-CA-C	5.13	117.21	109.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	904	0	848	23	1
1	B	882	0	823	17	0
2	A	15	0	13	0	0
2	B	15	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	45	0	0	0	0
3	B	49	0	0	0	0
All	All	1910	0	1697	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:GLN:HE22	1:B:314:THR:H	1.20	0.85
1:B:281:ASN:C	1:B:281:ASN:HD22	2.09	0.58
1:B:281:ASN:ND2	1:B:284:ARG:H	2.07	0.53
1:A:73:LEU:C	1:A:73:LEU:HD12	2.34	0.53
1:A:101:GLU:HB3	1:A:103:ARG:HH12	1.74	0.53
1:B:273:LEU:HD12	1:B:273:LEU:C	2.34	0.53
1:A:76:THR:HG22	1:A:77:VAL:N	2.24	0.52
1:A:91:THR:HB	1:B:291:THR:HB	1.92	0.52
1:B:256:LEU:C	1:B:256:LEU:HD12	2.35	0.52
1:A:64:PRO:HG3	1:A:71:THR:HG23	1.92	0.51
1:A:56:LEU:HD12	1:A:56:LEU:C	2.37	0.50
1:A:104:ILE:HB	1:A:130:PHE:HB2	1.94	0.49
1:B:281:ASN:C	1:B:281:ASN:ND2	2.70	0.49
1:A:29:PHE:HE1	1:A:56:LEU:CD2	2.28	0.47
1:A:95:GLN:HE22	1:B:314:THR:N	2.00	0.46
1:B:276:THR:HG22	1:B:277:VAL:N	2.30	0.45
1:A:120:TRP:CZ2	1:A:121:LYS:HD3	2.50	0.45
1:A:109:LEU:HG	1:B:309:LEU:HD11	1.98	0.45
1:A:73:LEU:C	1:A:73:LEU:CD1	2.90	0.45
1:A:63:ALA:N	1:A:64:PRO:HD3	2.32	0.44
1:A:29:PHE:CE1	1:A:56:LEU:HD21	2.53	0.44
1:B:273:LEU:HD12	1:B:273:LEU:O	2.18	0.43
1:A:73:LEU:HD12	1:A:73:LEU:O	2.18	0.43
1:A:101:GLU:HB3	1:A:103:ARG:NH1	2.33	0.43
1:B:281:ASN:ND2	1:B:283:TYR:H	2.17	0.42
1:A:118:ASN:OD1	1:A:121:LYS:HE2	2.19	0.42
1:A:20:THR:HG23	1:A:30:ILE:HD13	2.02	0.42
1:A:20:THR:O	1:A:133:VAL:HG12	2.19	0.42
1:A:33:ALA:HB1	1:A:60:TYR:CZ	2.56	0.41
1:B:271:THR:HG23	1:B:296:TYR:HB3	2.03	0.41
1:A:72:ALA:CB	1:B:288:SER:HA	2.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:263:ALA:N	1:B:264:PRO:HD3	2.36	0.40
1:A:29:PHE:CE1	1:A:56:LEU:CD2	3.05	0.40
1:B:230:ILE:O	1:B:241:GLY:HA3	2.20	0.40
1:B:253:ARG:HG3	1:B:253:ARG:HH11	1.87	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ASP:OD1	1:A:67:ASP:OD1[3_655]	1.79	0.41

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/121 (98%)	111 (93%)	8 (7%)	0	100	100
1	B	116/121 (96%)	112 (97%)	3 (3%)	1 (1%)	14	17
All	All	235/242 (97%)	223 (95%)	11 (5%)	1 (0%)	30	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	267	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/89 (100%)	88 (99%)	1 (1%)	65	81
1	B	87/89 (98%)	83 (95%)	4 (5%)	24	36
All	All	176/178 (99%)	171 (97%)	5 (3%)	38	56

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	VAL
1	B	231	VAL
1	B	271	THR
1	B	281	ASN
1	B	303	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	107	GLN
1	B	281	ASN
1	B	282	ASN
1	B	318	ASN
1	B	327	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SNR	A	1	-	16,16,16	1.49	2 (12%)	22,22,22	1.94	7 (31%)
2	SNR	B	2	-	16,16,16	1.47	2 (12%)	22,22,22	1.91	7 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SNR	A	1	-	-	4/6/27/27	0/2/2/2
2	SNR	B	2	-	-	4/6/27/27	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	SNR	O17-C15	4.17	1.35	1.22
2	A	1	SNR	O17-C15	4.14	1.35	1.22
2	A	1	SNR	O16-C15	-2.58	1.22	1.30
2	B	2	SNR	O16-C15	-2.57	1.22	1.30

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	SNR	C6-C4-N5	3.85	117.41	113.34
2	B	2	SNR	C3-C8-S7	-3.42	101.35	106.06
2	A	1	SNR	C8-C3-N2	-3.39	108.82	113.18
2	B	2	SNR	C6-C4-N5	3.36	116.90	113.34
2	B	2	SNR	C8-C3-N2	-3.35	108.86	113.18
2	A	1	SNR	C3-C8-S7	-2.94	102.01	106.06
2	B	2	SNR	C13-C14-C15	-2.77	107.28	114.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	SNR	N5-C1-N2	2.74	112.01	108.85
2	B	2	SNR	C4-C6-S7	-2.72	101.94	105.03
2	A	1	SNR	C13-C14-C15	-2.68	107.51	114.51
2	A	1	SNR	C8-C3-C4	2.65	113.26	109.06
2	B	2	SNR	N5-C1-N2	2.54	111.79	108.85
2	B	2	SNR	C8-C3-C4	2.31	112.73	109.06
2	A	1	SNR	C4-C6-S7	-2.15	102.59	105.03

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	SNR	C13-C12-C6-C4
2	A	1	SNR	C13-C12-C6-S7
2	B	2	SNR	C13-C12-C6-C4
2	B	2	SNR	C13-C12-C6-S7
2	A	1	SNR	C13-C14-C15-O16
2	A	1	SNR	C13-C14-C15-O17
2	B	2	SNR	C13-C14-C15-O17
2	B	2	SNR	C13-C14-C15-O16

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.