



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:34 PM UTC

PDB ID : 4LCD / pdb_00004lcd
Title : Structure of an Rsp5UbxSna3 complex: Mechanism of ubiquitin ligation and lysine prioritization by a HECT E3
Authors : Kamadurai, H.B.; Miller, D.; Schulman, B.A.
Deposited on : 2013-06-21
Resolution : 3.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 64%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 31%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 3%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">64% 31% . .</p>
1	B	432	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 61%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 32%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 3%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">61% 32% . .</p>
2	C	24	<div style="display: flex; align-items: center;"> <div style="width: 25%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">25% 8% 67%</p>
2	D	24	<div style="display: flex; align-items: center;"> <div style="width: 17%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 33%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 42%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">17% 33% 8% 42%</p>
3	E	83	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 54%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 28%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">54% 28% 8% 10%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	83	 <p>A horizontal bar chart showing the quality of chain. The bar is divided into four segments: 46% (green), 35% (yellow), 10% (orange), and 10% (grey). A small red square is at the beginning of the bar, and a '%' symbol is above it.</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase RSP5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	419	3389	2169	585	624	11	0	0	0
1	B	413	3289	2110	565	604	10	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	378	GLY	-	expression tag	UNP P39940
A	379	SER	-	expression tag	UNP P39940
A	380	GLY	-	expression tag	UNP P39940
A	381	GLY	-	expression tag	UNP P39940
A	382	SER	-	expression tag	UNP P39940
A	455	LEU	CYS	engineered mutation	UNP P39940
A	517	GLY	CYS	engineered mutation	UNP P39940
A	721	ALA	CYS	engineered mutation	UNP P39940
B	378	GLY	-	expression tag	UNP P39940
B	379	SER	-	expression tag	UNP P39940
B	380	GLY	-	expression tag	UNP P39940
B	381	GLY	-	expression tag	UNP P39940
B	382	SER	-	expression tag	UNP P39940
B	455	LEU	CYS	engineered mutation	UNP P39940
B	517	GLY	CYS	engineered mutation	UNP P39940
B	721	ALA	CYS	engineered mutation	UNP P39940

- Molecule 2 is a protein called Protein SNA3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	8	51	34	8	9	0	0	0
2	D	14	80	51	14	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	125	ALA	LYS	engineered mutation	UNP P14359
D	125	ALA	LYS	engineered mutation	UNP P14359

- Molecule 3 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
3	E	75	Total	C	N	O	S	0	0	0
			565	353	98	112	2			
3	F	75	Total	C	N	O	S	0	0	0
			570	359	96	113	2			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-7	MET	-	expression tag	UNP P0CG48
E	-6	GLY	-	expression tag	UNP P0CG48
E	-5	HIS	-	expression tag	UNP P0CG48
E	-4	HIS	-	expression tag	UNP P0CG48
E	-3	HIS	-	expression tag	UNP P0CG48
E	-2	HIS	-	expression tag	UNP P0CG48
E	-1	HIS	-	expression tag	UNP P0CG48
E	0	HIS	-	expression tag	UNP P0CG48
E	75	CYS	GLY	engineered mutation	UNP P0CG48
F	-7	MET	-	expression tag	UNP P0CG48
F	-6	GLY	-	expression tag	UNP P0CG48
F	-5	HIS	-	expression tag	UNP P0CG48
F	-4	HIS	-	expression tag	UNP P0CG48
F	-3	HIS	-	expression tag	UNP P0CG48
F	-2	HIS	-	expression tag	UNP P0CG48
F	-1	HIS	-	expression tag	UNP P0CG48
F	0	HIS	-	expression tag	UNP P0CG48
F	75	CYS	GLY	engineered mutation	UNP P0CG48


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: E3 ubiquitin-protein ligase RSP5




• Molecule 1: E3 ubiquitin-protein ligase RSP5





- Molecule 2: Protein SNA3



- Molecule 2: Protein SNA3



- Molecule 3: Ubiquitin



- Molecule 3: Ubiquitin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.05Å 78.92Å 96.72Å 90.00° 101.67° 90.00°	Depositor
Resolution (Å)	29.40 – 3.10 29.40 – 3.10	Depositor EDS
% Data completeness (in resolution range)	95.0 (29.40-3.10) 91.3 (29.40-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 3.11Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.251 , 0.299 0.252 , 0.297	Depositor DCC
R_{free} test set	1091 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	72.9	Xtrriage
Anisotropy	0.519	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 64.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7944	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/3472	0.84	1/4702 (0.0%)
1	B	0.56	0/3371	0.85	0/4574
2	C	0.73	0/53	1.51	3/74 (4.1%)
2	D	0.39	0/82	1.24	1/114 (0.9%)
3	E	0.50	0/571	0.91	1/774 (0.1%)
3	F	0.46	0/576	0.90	3/782 (0.4%)
All	All	0.56	0/8125	0.87	9/11020 (0.1%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	110	ASP	N-CA-C	-8.09	103.31	113.01
2	C	105	GLN	CA-C-N	6.76	127.34	120.38
2	C	105	GLN	C-N-CA	6.76	127.34	120.38
3	F	36	ILE	CA-C-N	6.68	124.53	119.66
3	F	36	ILE	C-N-CA	6.68	124.53	119.66

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3389	0	3222	88	0
1	B	3289	0	3067	108	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	51	0	35	0	0
2	D	80	0	57	10	0
3	E	565	0	551	21	0
3	F	570	0	565	32	0
All	All	7944	0	7497	243	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 243 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:ARG:CG	1:B:449:ARG:HH11	1.95	0.78
1:B:599:LEU:HD11	1:B:644:VAL:HG21	1.68	0.75
1:B:408:HIS:HD1	2:D:109:TYR:HH	1.36	0.73
3:F:71:LEU:HD12	3:F:71:LEU:N	2.05	0.71
1:B:449:ARG:HH11	1:B:449:ARG:HG2	1.57	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/432 (96%)	390 (94%)	25 (6%)	0	100	100
1	B	407/432 (94%)	366 (90%)	41 (10%)	0	100	100
2	C	6/24 (25%)	2 (33%)	4 (67%)	0	100	100
2	D	12/24 (50%)	6 (50%)	5 (42%)	1 (8%)	0	4
3	E	73/83 (88%)	63 (86%)	10 (14%)	0	100	100
3	F	73/83 (88%)	65 (89%)	8 (11%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	986/1078 (92%)	892 (90%)	93 (9%)	1 (0%)	48 78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	111	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	353/386 (92%)	329 (93%)	24 (7%)	14 42
1	B	333/386 (86%)	307 (92%)	26 (8%)	11 38
2	C	3/18 (17%)	3 (100%)	0	100 100
2	D	3/18 (17%)	3 (100%)	0	100 100
3	E	60/76 (79%)	50 (83%)	10 (17%)	2 10
3	F	62/76 (82%)	52 (84%)	10 (16%)	2 11
All	All	814/960 (85%)	744 (91%)	70 (9%)	10 34

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	72	ARG
3	E	75	CYS
3	F	55	THR
1	B	394	GLU
1	A	806	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	746	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	2	GLN
3	F	31	GLN
1	A	668	ASN
1	A	735	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/432 (96%)	-0.02	5 (1%) 76 58	43, 84, 147, 180	0
1	B	413/432 (95%)	0.13	4 (0%) 79 61	46, 92, 140, 206	0
2	C	8/24 (33%)	0.73	0 100 100	107, 128, 133, 134	0
2	D	14/24 (58%)	0.47	0 100 100	113, 143, 157, 161	0
3	E	75/83 (90%)	0.17	1 (1%) 75 56	84, 124, 183, 204	0
3	F	75/83 (90%)	0.43	1 (1%) 75 56	94, 122, 143, 157	0
All	All	1004/1078 (93%)	0.10	11 (1%) 78 59	43, 96, 149, 206	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	589	ALA	4.9
1	A	754	SER	2.9
1	B	603	ILE	2.8
1	A	401	ALA	2.6
3	E	33	LYS	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.