



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:01 AM UTC

PDB ID : 4LCQ / pdb_00004lcq
Title : The crystal structure of di-Zn dihydropyrimidinase in complex with NCBI
Authors : Hsieh, Y.C.; Chen, M.C.; Hsu, C.C.; Chan, S.I.; Yang, Y.S.; Chen, C.J.
Deposited on : 2013-06-22
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	URQ	A	603	-	X	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4018 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called dihydropyrimidinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	0	0
			3759	2365	666	704	24			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is (2S)-3-(carbamoylamino)-2-methylpropanoic acid (CCD ID: URQ) (formula: C₅H₁₀N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	5	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	247	Total 247	O 247	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: dihydropyrimidinase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	161.20Å 161.20Å 93.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.81 30.00 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-1.81) 99.8 (30.00-1.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.17 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.174 , 0.200 0.173 , 0.200	Depositor DCC
R_{free} test set	2871 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	22.6	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 29.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4018	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, ZN, URQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.55	29/3828 (0.8%)	1.31	12/5182 (0.2%)

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	460	HIS	CG-CD2	9.33	1.46	1.35
1	A	461	ARG	CD-NE	-8.52	1.34	1.46
1	A	213	HIS	CE1-NE2	7.90	1.40	1.32
1	A	182	GLY	N-CA	7.06	1.54	1.45
1	A	65	HIS	ND1-CE1	6.92	1.39	1.32
1	A	460	HIS	ND1-CE1	6.49	1.39	1.32
1	A	129	VAL	N-CA	6.29	1.54	1.46
1	A	163	LEU	CA-C	6.27	1.58	1.52
1	A	63	HIS	ND1-CE1	6.24	1.38	1.32
1	A	226	THR	CA-C	6.01	1.60	1.52
1	A	57	PRO	C-O	5.85	1.30	1.23
1	A	78	ASP	C-O	5.77	1.31	1.24
1	A	121	VAL	N-CA	5.75	1.52	1.46
1	A	282	HIS	CG-CD2	5.64	1.42	1.35
1	A	376	SER	CA-CB	5.64	1.62	1.53
1	A	14	VAL	N-CA	5.63	1.53	1.46
1	A	378	ALA	N-CA	5.58	1.53	1.46
1	A	361	HIS	ND1-CE1	5.42	1.38	1.32
1	A	128	HIS	ND1-CE1	5.36	1.38	1.32
1	A	42	PRO	CA-C	5.21	1.57	1.52
1	A	217	ARG	CZ-NH2	5.19	1.40	1.33
1	A	282	HIS	ND1-CE1	5.19	1.37	1.32
1	A	114	ARG	CA-C	5.14	1.59	1.52
1	A	460	HIS	CE1-NE2	5.14	1.37	1.32
1	A	367	GLU	N-CA	5.11	1.52	1.46
1	A	469	TYR	C-O	-5.11	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	285	TRP	CD2-CE2	5.09	1.50	1.41
1	A	239	PRO	C-O	5.08	1.29	1.23
1	A	30	GLY	N-CA	5.05	1.52	1.45

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	461	ARG	NE-CZ-NH2	-13.11	107.40	119.20
1	A	461	ARG	NE-CZ-NH1	10.28	131.78	121.50
1	A	461	ARG	CD-NE-CZ	8.09	135.72	124.40
1	A	461	ARG	CB-CG-CD	-6.72	95.84	111.30
1	A	40	ARG	CA-C-N	-6.69	118.05	122.60
1	A	40	ARG	C-N-CA	-6.69	118.05	122.60
1	A	46	ARG	CA-CB-CG	-6.64	100.81	114.10
1	A	165	MET	N-CA-C	6.44	119.32	110.24
1	A	472	ILE	N-CA-C	-6.23	104.44	110.42
1	A	473	ARG	NE-CZ-NH1	-6.01	115.48	121.50
1	A	219	GLU	N-CA-C	6.00	119.44	111.75
1	A	43	PRO	CB-CA-C	-5.13	105.12	111.64

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3759	0	3704	32	0
2	A	2	0	0	0	0
3	A	10	0	7	4	0
4	A	247	0	0	2	0
All	All	4018	0	3711	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:603:URQ:H5	3:A:603:URQ:C4	1.54	1.18
1:A:466:GLU:HG3	1:A:470:LYS:HE3	1.38	1.00
1:A:16:ASN:HD21	1:A:391:ILE:H	1.13	0.94
1:A:466:GLU:CG	1:A:470:LYS:HE3	2.04	0.88
3:A:603:URQ:C4	3:A:603:URQ:N3	2.39	0.85
1:A:415:HIS:HD2	1:A:422:ASN:HD22	1.26	0.82
1:A:382:PHE:O	1:A:461:ARG:CD	2.31	0.79
1:A:382:PHE:O	1:A:461:ARG:HD2	1.86	0.74
1:A:487:GLU:HG2	1:A:488:PRO:HD2	1.78	0.65
1:A:382:PHE:O	1:A:461:ARG:HD3	1.96	0.64
1:A:7:ILE:HG23	1:A:48:ILE:HD13	1.81	0.63
1:A:16:ASN:ND2	1:A:391:ILE:H	1.92	0.60
3:A:603:URQ:N3	3:A:603:URQ:O41	2.30	0.60
3:A:603:URQ:H5	3:A:603:URQ:C5	2.04	0.58
1:A:401:ILE:N	1:A:401:ILE:HD12	2.18	0.58
1:A:323:ASN:HD22	1:A:348:VAL:H	1.52	0.57
1:A:244:HIS:CE1	1:A:295:PRO:HD3	2.40	0.56
1:A:40:ARG:HG2	4:A:944:HOH:O	2.05	0.56
1:A:48:ILE:CD1	1:A:48:ILE:N	2.68	0.56
1:A:48:ILE:N	1:A:48:ILE:HD12	2.23	0.54
1:A:483:VAL:HG12	1:A:484:VAL:O	2.08	0.54
1:A:116:THR:O	1:A:120:LYS:HD3	2.07	0.54
1:A:285:TRP:CZ2	1:A:335:LYS:HA	2.46	0.50
1:A:279:ASN:OD1	1:A:287:HIS:HE1	1.95	0.50
1:A:198:GLY:C	1:A:216:CYS:HB2	2.37	0.49
1:A:100:THR:HG23	1:A:130:ALA:O	2.12	0.49
1:A:139:LYS:O	1:A:177:HIS:HE1	1.94	0.49
1:A:445:GLU:OE2	1:A:450:LYS:NZ	2.46	0.48
1:A:393:LYS:O	1:A:394:ASP:HB2	2.16	0.46
1:A:146:ALA:HA	1:A:151:VAL:O	2.15	0.46
1:A:436:THR:HB	1:A:444:TYR:HB3	2.02	0.42
1:A:401:ILE:HD13	1:A:435:VAL:HB	2.02	0.41
1:A:213:HIS:HE1	1:A:343:ASN:HD21	1.69	0.41
1:A:46:ARG:HD3	4:A:901:HOH:O	2.19	0.41
1:A:214:GLU:HG2	1:A:296:PRO:HB2	2.03	0.41
1:A:412:ALA:HA	1:A:415:HIS:CD2	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	482/520 (93%)	465 (96%)	13 (3%)	4 (1%)	16 6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	383	ASN
1	A	244	HIS
1	A	5	GLY
1	A	347	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	399/422 (94%)	390 (98%)	9 (2%)	44 28

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	48	ILE
1	A	100	THR
1	A	102	LYS
1	A	200	LYS
1	A	405	LYS
1	A	413	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	415	HIS
1	A	466	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	147	GLN
1	A	177	HIS
1	A	287	HIS
1	A	290	GLN
1	A	313	ASN
1	A	323	ASN
1	A	343	ASN
1	A	415	HIS
1	A	460	HIS
1	A	462	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	A	155	1,2	10,11,12	1.12	1 (10%)	6,12,14	1.37	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	155	1,2	-	1/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	155	KCX	CX-NZ	-2.24	1.31	1.35

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	KCX	CD-CG-CB	-2.27	105.08	113.62

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	155	KCX	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	URQ	A	603	2	9,9,9	4.01	8 (88%)	11,11,11	4.35	9 (81%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	URQ	A	603	2	-	5/9/9/9	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	URQ	C7-C5	-5.89	1.33	1.52
3	A	603	URQ	C2-N1	-5.18	1.27	1.34
3	A	603	URQ	O2-C2	-4.72	1.16	1.24
3	A	603	URQ	C6-C5	-4.43	1.38	1.52
3	A	603	URQ	C2-N3	-3.89	1.24	1.33
3	A	603	URQ	O41-C4	-3.54	1.11	1.22
3	A	603	URQ	C6-N1	-2.87	1.38	1.46
3	A	603	URQ	O42-C4	-2.34	1.23	1.30

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	603	URQ	C6-C5-C4	8.72	126.72	110.61
3	A	603	URQ	C7-C5-C4	5.96	124.88	109.16
3	A	603	URQ	O42-C4-C5	5.03	131.38	115.05
3	A	603	URQ	O42-C4-O41	-4.84	113.11	124.08
3	A	603	URQ	C6-N1-C2	4.14	128.15	122.91
3	A	603	URQ	C5-C6-N1	3.50	131.06	113.09
3	A	603	URQ	O41-C4-C5	-2.29	115.30	122.27
3	A	603	URQ	O2-C2-N1	2.28	125.28	121.69
3	A	603	URQ	O2-C2-N3	-2.21	118.74	123.18

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	603	URQ	C5-C6-N1-C2
3	A	603	URQ	O2-C2-N1-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	603	URQ	N3-C2-N1-C6
3	A	603	URQ	C7-C5-C6-N1
3	A	603	URQ	C4-C5-C6-N1

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	URQ	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	484/520 (93%)	-0.38	5 (1%) 79 84	16, 23, 37, 79	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	ALA	6.8
1	A	5	GLY	3.6
1	A	488	PRO	3.5
1	A	483	VAL	2.0
1	A	466	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	155	12/13	0.95	0.07	12,18,19,19	3

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	URQ	A	603	10/10	0.87	0.13	32,41,45,55	0
2	ZN	A	602	1/1	0.99	0.06	26,26,26,26	0
2	ZN	A	601	1/1	0.99	0.07	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.