



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 11:05 PM UTC

PDB ID : 5LDY / pdb_00005ldy
Title : Structure of Yersinia pseudotuberculosis InvD
Authors : Sadana, P.; Scrima, A.
Deposited on : 2016-06-28
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 85% 9% 6%
1	B	357	 88% 7% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9725 atoms, of which 4782 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig domain protein group 1 domain protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	335	Total	C	H	N	O	S	0	0	0
			4795	1485	2382	410	512	6			
1	B	339	Total	C	H	N	O	S	0	0	0
			4836	1497	2400	417	516	6			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1620	MET	-	initiating methionine	UNP A0A0H3B1G5
A	1621	GLY	-	expression tag	UNP A0A0H3B1G5
A	1622	HIS	-	expression tag	UNP A0A0H3B1G5
A	1623	HIS	-	expression tag	UNP A0A0H3B1G5
A	1624	HIS	-	expression tag	UNP A0A0H3B1G5
A	1625	HIS	-	expression tag	UNP A0A0H3B1G5
A	1626	HIS	-	expression tag	UNP A0A0H3B1G5
A	1627	HIS	-	expression tag	UNP A0A0H3B1G5
A	1628	VAL	-	expression tag	UNP A0A0H3B1G5
A	1629	ASP	-	expression tag	UNP A0A0H3B1G5
A	1630	GLU	-	expression tag	UNP A0A0H3B1G5
A	1631	ASN	-	expression tag	UNP A0A0H3B1G5
A	1632	LEU	-	expression tag	UNP A0A0H3B1G5
A	1633	TYR	-	expression tag	UNP A0A0H3B1G5
A	1634	PHE	-	expression tag	UNP A0A0H3B1G5
A	1635	GLN	-	expression tag	UNP A0A0H3B1G5
A	1636	GLY	-	expression tag	UNP A0A0H3B1G5
A	1637	GLY	-	expression tag	UNP A0A0H3B1G5
A	1638	GLY	-	expression tag	UNP A0A0H3B1G5
A	1639	ARG	-	expression tag	UNP A0A0H3B1G5
B	1620	MET	-	initiating methionine	UNP A0A0H3B1G5
B	1621	GLY	-	expression tag	UNP A0A0H3B1G5
B	1622	HIS	-	expression tag	UNP A0A0H3B1G5
B	1623	HIS	-	expression tag	UNP A0A0H3B1G5
B	1624	HIS	-	expression tag	UNP A0A0H3B1G5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1625	HIS	-	expression tag	UNP A0A0H3B1G5
B	1626	HIS	-	expression tag	UNP A0A0H3B1G5
B	1627	HIS	-	expression tag	UNP A0A0H3B1G5
B	1628	VAL	-	expression tag	UNP A0A0H3B1G5
B	1629	ASP	-	expression tag	UNP A0A0H3B1G5
B	1630	GLU	-	expression tag	UNP A0A0H3B1G5
B	1631	ASN	-	expression tag	UNP A0A0H3B1G5
B	1632	LEU	-	expression tag	UNP A0A0H3B1G5
B	1633	TYR	-	expression tag	UNP A0A0H3B1G5
B	1634	PHE	-	expression tag	UNP A0A0H3B1G5
B	1635	GLN	-	expression tag	UNP A0A0H3B1G5
B	1636	GLY	-	expression tag	UNP A0A0H3B1G5
B	1637	GLY	-	expression tag	UNP A0A0H3B1G5
B	1638	GLY	-	expression tag	UNP A0A0H3B1G5
B	1639	ARG	-	expression tag	UNP A0A0H3B1G5

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	48	Total 48	O 48	0	0
3	B	36	Total 36	O 36	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ig domain protein group 1 domain protein



- Molecule 1: Ig domain protein group 1 domain protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	58.90Å 58.90Å 274.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.70 – 2.60 44.70 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (44.70-2.60) 99.9 (44.70-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.71 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.230 , 0.253 (Not available) , 0.270	Depositor DCC
R_{free} test set	1427 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	52.0	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.480 for h,-k,-l	Xtriage
Reported twinning fraction	0.500 for h,-k,-l	Depositor
Outliers	0 of 28540 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9725	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/2442	0.26	0/3348
1	B	0.12	0/2465	0.27	0/3378
All	All	0.12	0/4907	0.27	0/6726

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2413	2382	2381	18	3
1	B	2436	2400	2400	13	3
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	48	0	0	3	0
3	B	36	0	0	3	0
All	All	4943	4782	4781	31	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 31 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1958:ASP:OD1	1:A:1959:ASP:N	2.15	0.80
1:A:1886:ILE:O	3:A:2101:HOH:O	2.03	0.76
1:B:1680:GLY:N	1:B:1697:GLU:OE2	2.19	0.74
1:A:1756:ALA:O	1:A:1759:ARG:NH2	2.20	0.73
1:A:1759:ARG:NH1	1:A:1808:ASN:OD1	2.22	0.73

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1662:ASP:OD1	1:B:1700:SER:OG[3_544]	1.98	0.22
1:A:1914:GLY:O	1:B:1639:ARG:NH1[1_455]	2.08	0.12
1:A:1918:ARG:NH2	1:B:1683:VAL:O[1_455]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/286 (94%)	268 (100%)	1 (0%)	84	93
1	B	270/286 (94%)	268 (99%)	2 (1%)	76	89
All	All	539/572 (94%)	536 (99%)	3 (1%)	78	91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1939	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1639	ARG
1	B	1759	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1681	GLN
1	A	1820	GLN
1	A	1881	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	2001	-	4,4,4	0.98	0	6,6,6	0.48	0
2	PO4	B	2001	-	4,4,4	1.01	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/357 (93%)	-0.95	0 100 100	37, 63, 104, 150	0
1	B	339/357 (94%)	-0.91	0 100 100	41, 63, 109, 153	0
All	All	674/714 (94%)	-0.93	0 100 100	37, 63, 107, 153	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	2001	5/5	0.98	0.06	79,80,81,84	0
2	PO4	B	2001	5/5	0.98	0.07	88,93,98,98	0

6.5 Other polymers [i](#)

There are no such residues in this entry.