



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:09 AM UTC

PDB ID : 7LD9 / pdb_00007ld9
Title : Structure of human GGT1 in complex with ABBA
Authors : Terzyan, S.S.; Hanigan, M.
Deposited on : 2021-01-12
Resolution : 1.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	
2	B	189	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5151 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione hydrolase 1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	343	2794	1765	500	515	14	0	21	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	272	ALA	VAL	engineered mutation	UNP P19440

- Molecule 2 is a protein called Glutathione hydrolase 1 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	189	1470	920	260	284	6	0	8	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
6	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
6	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 7 is (2R)-2-amino-4-boronobutanoic acid (CCD ID: XUJ) (formula: $C_4H_{10}BNO_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	B	C	N	O	0	1
			10	1	4	1	4		

- Molecule 8 is (2S)-2-amino-4-boronobutanoic acid (CCD ID: XU7) (formula: $C_4H_{10}BNO_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total	B	C	N	O	0	1
			10	1	4	1	4		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	463	Total	O	0	0
			463	463		
9	B	272	Total	O	0	0
			272	272		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutathione hydrolase 1 heavy chain



- Molecule 2: Glutathione hydrolase 1 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	105.32Å 125.57Å 104.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.69 – 1.42 80.69 – 1.42	Depositor EDS
% Data completeness (in resolution range)	99.5 (80.69-1.42) 99.5 (80.69-1.42)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 1.42Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.107 , 0.134 0.106 , 0.133	Depositor DCC
R_{free} test set	6480 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	15.5	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5151	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, XUJ, NA, NAG, XU7, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	2/2872 (0.1%)	0.81	2/3887 (0.1%)
2	B	0.70	0/1501	0.83	0/2049
All	All	0.72	2/4373 (0.0%)	0.82	2/5936 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	36	HIS	N-CA	5.57	1.53	1.46
1	A	266	ASN	N-CA	-5.08	1.39	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	36	HIS	CB-CA-C	-6.68	102.67	111.42
1	A	254	ASN	CA-CB-CG	5.22	117.82	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2794	0	2812	24	0
2	B	1470	0	1441	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	70	0	65	0	0
3	B	14	0	13	1	0
4	A	1	0	0	0	0
4	B	1	0	0	1	0
5	A	1	0	0	0	0
6	A	15	0	18	0	0
6	B	30	0	36	1	0
7	B	10	0	0	1	0
8	B	10	0	0	0	0
9	A	463	0	0	9	1
9	B	272	0	0	5	0
All	All	5151	0	4385	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139[A]:ARG:HG2	1:A:229:TYR:CE2	2.25	0.72
1:A:142:GLU:HG3	9:A:1081:HOH:O	1.91	0.70
2:B:511:ASN:O	9:B:1201:HOH:O	2.11	0.68
7:B:1103[A]:XUJ:NAF	9:B:1202:HOH:O	2.28	0.67
1:A:150[B]:ARG:CA	1:A:151:LEU:N	2.63	0.61
1:A:149:GLY:C	1:A:150[B]:ARG:CA	2.74	0.60
1:A:48:LYS:HE2	1:A:52:LYS:HE3	1.83	0.60
1:A:78[B]:MET:HE3	9:A:1106:HOH:O	2.02	0.59
1:A:47:ALA:HB3	1:A:50[B]:CYS:SG	2.44	0.57
1:A:126[A]:GLN:HG2	2:B:422:ASP:CG	2.31	0.56
1:A:187:GLN:CG	9:A:908:HOH:O	2.54	0.55
1:A:139[A]:ARG:NH2	9:A:701:HOH:O	2.11	0.53
2:B:402:LEU:HD22	6:B:1105:EPE:O2S	2.08	0.53
1:A:187:GLN:HG3	9:A:908:HOH:O	2.08	0.53
1:A:126[A]:GLN:HG2	2:B:422:ASP:OD2	2.10	0.51
1:A:99:ARG:HD3	9:A:889:HOH:O	2.10	0.51
1:A:326:LYS:NZ	9:A:706:HOH:O	2.41	0.49
1:A:49:GLN:HB3	1:A:78[A]:MET:SD	2.52	0.48
2:B:387[B]:VAL:HG22	2:B:547:ILE:HD13	1.97	0.47
2:B:444:GLN:HG3	9:B:1422:HOH:O	2.15	0.47
3:B:1101:NAG:H62	9:B:1204:HOH:O	2.14	0.46
1:A:340:GLU:HG2	1:A:343:ARG:HH21	1.80	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:382:ALA:HA	4:B:1102:CL:CL	2.53	0.46
1:A:146:GLN:HG2	9:A:1069:HOH:O	2.16	0.45
1:A:340:GLU:HG2	1:A:343:ARG:NH2	2.31	0.45
2:B:562:LYS:NZ	9:B:1203:HOH:O	2.32	0.45
1:A:294:LYS:NZ	1:A:348:GLU:OE2	2.39	0.45
1:A:52:LYS:HE2	9:A:1043:HOH:O	2.18	0.43
1:A:182[A]:ARG:HE	1:A:182[A]:ARG:HB2	1.36	0.43
1:A:68:ALA:HA	2:B:384[A]:LEU:HD21	2.01	0.42
2:B:409:ARG:HB2	2:B:416[B]:LEU:HD23	2.02	0.42
1:A:295:GLY:HA3	1:A:355[B]:ARG:CZ	2.50	0.42
1:A:288:LEU:HD23	1:A:323:ALA:HA	2.02	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:995:HOH:O	9:A:995:HOH:O[3_554]	2.05	0.15

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/353 (103%)	359 (99%)	4 (1%)	0	100	100
2	B	195/189 (103%)	187 (96%)	7 (4%)	1 (0%)	24	7
All	All	558/542 (103%)	546 (98%)	11 (2%)	1 (0%)	43	20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	401	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/282 (105%)	296 (100%)	0	100	100
2	B	159/151 (105%)	157 (99%)	2 (1%)	61	29
All	All	455/433 (105%)	453 (100%)	2 (0%)	87	67

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	384[A]	LEU
2	B	384[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
2	B	464	GLN
2	B	535	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 3 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EPE	B	1106	-	15,15,15	1.91	3 (20%)	19,20,20	1.65	4 (21%)
6	EPE	A	608	-	15,15,15	1.62	3 (20%)	19,20,20	3.99	8 (42%)
6	EPE	B	1105	-	15,15,15	1.21	1 (6%)	19,20,20	1.98	6 (31%)
3	NAG	A	605	1	14,14,15	0.67	0	17,19,21	1.20	2 (11%)
3	NAG	A	603	1	14,14,15	0.55	0	17,19,21	1.30	1 (5%)
3	NAG	A	604	1	14,14,15	0.61	0	17,19,21	2.47	5 (29%)
3	NAG	A	601	1	14,14,15	0.53	0	17,19,21	1.32	3 (17%)
7	XUJ	B	1103[A]	2	8,9,9	1.75	1 (12%)	6,11,11	2.88	2 (33%)
3	NAG	A	602	1	14,14,15	0.94	1 (7%)	17,19,21	1.32	1 (5%)
8	XU7	B	1104[B]	2	8,9,9	1.75	1 (12%)	6,11,11	2.85	2 (33%)
3	NAG	B	1101	2	14,14,15	0.67	0	17,19,21	1.55	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EPE	B	1106	-	-	0/9/19/19	0/1/1/1
6	EPE	A	608	-	-	1/9/19/19	0/1/1/1
6	EPE	B	1105	-	-	0/9/19/19	0/1/1/1
3	NAG	A	605	1	-	1/6/23/26	0/1/1/1
3	NAG	A	603	1	-	0/6/23/26	0/1/1/1
3	NAG	A	604	1	-	2/6/23/26	0/1/1/1
3	NAG	A	601	1	-	0/6/23/26	0/1/1/1
7	XUJ	B	1103[A]	2	-	1/7/9/9	-
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1
8	XU7	B	1104[B]	2	-	0/7/9/9	-
3	NAG	B	1101	2	-	2/6/23/26	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1106	EPE	O1S-S	4.75	1.58	1.45
6	A	608	EPE	C10-S	4.66	1.84	1.77
7	B	1103[A]	XUJ	BAI-CAH	3.80	1.63	1.56
8	B	1104[B]	XU7	BAI-CAH	3.80	1.63	1.56
6	B	1106	EPE	O2S-S	3.33	1.54	1.45
6	B	1106	EPE	C10-S	2.86	1.81	1.77
6	A	608	EPE	O2S-S	2.82	1.53	1.45
6	B	1105	EPE	O2S-S	2.58	1.52	1.45
3	A	602	NAG	O5-C1	-2.38	1.39	1.43
6	A	608	EPE	O1S-S	2.18	1.51	1.45

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	608	EPE	O2S-S-C10	-10.84	90.35	106.73
6	A	608	EPE	O3S-S-C10	-9.11	88.18	106.00
3	A	604	NAG	C1-O5-C5	8.00	122.90	112.19
6	B	1105	EPE	O1S-S-C10	6.16	116.04	106.73
7	B	1103[A]	XUJ	OAL-BAI-CAH	-5.82	108.14	121.06
8	B	1104[B]	XU7	OAL-BAI-CAH	-5.82	108.14	121.06
6	A	608	EPE	O1S-S-C10	-5.40	98.57	106.73
6	A	608	EPE	C6-N1-C2	5.28	120.22	108.84
3	A	603	NAG	C1-O5-C5	4.66	118.44	112.19
3	B	1101	NAG	O5-C1-C2	-4.30	104.64	111.29
3	A	602	NAG	C1-C2-N2	4.06	116.83	110.43
6	A	608	EPE	O3S-S-O1S	3.89	121.14	111.40
6	B	1106	EPE	O1S-S-C10	-3.89	100.85	106.73
3	A	605	NAG	C1-O5-C5	3.87	117.37	112.19
7	B	1103[A]	XUJ	OAK-BAI-CAH	-3.83	112.55	121.06
8	B	1104[B]	XU7	OAK-BAI-CAH	-3.83	112.55	121.06
3	B	1101	NAG	C1-C2-N2	3.41	115.80	110.43
6	B	1105	EPE	C7-N4-C5	-3.32	102.39	111.24
3	A	604	NAG	O5-C5-C4	3.28	118.81	110.83
6	A	608	EPE	O3S-S-O2S	3.25	119.53	111.40
6	B	1106	EPE	O3S-S-O1S	3.12	119.20	111.40
3	A	604	NAG	C3-C4-C5	2.78	115.28	110.23
6	B	1106	EPE	C5-N4-C3	2.68	114.62	108.84
3	A	601	NAG	O5-C1-C2	-2.68	107.15	111.29
3	A	601	NAG	C1-O5-C5	2.67	115.76	112.19
6	B	1105	EPE	C5-N4-C3	2.52	114.27	108.84
6	B	1106	EPE	O2S-S-O1S	-2.50	105.69	113.82
6	B	1105	EPE	C9-N1-C2	2.42	117.69	111.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	604	NAG	C4-C3-C2	2.36	114.47	111.02
3	A	604	NAG	O7-C7-C8	-2.29	117.98	122.05
6	B	1105	EPE	O3S-S-O1S	-2.25	105.78	111.40
3	A	601	NAG	C4-C3-C2	2.25	114.31	111.02
3	A	605	NAG	O5-C1-C2	-2.17	107.93	111.29
6	A	608	EPE	C10-C9-N1	-2.11	104.37	112.36
6	B	1105	EPE	C6-N1-C2	2.05	113.25	108.84
6	A	608	EPE	C3-C2-N1	2.02	114.72	110.65

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	604	NAG	O5-C5-C6-O6
3	A	604	NAG	C4-C5-C6-O6
3	A	605	NAG	O5-C5-C6-O6
3	B	1101	NAG	C1-C2-N2-C7
3	B	1101	NAG	C3-C2-N2-C7
6	A	608	EPE	C10-C9-N1-C2
7	B	1103[A]	XUJ	OAA-CAB-CAD-NAF

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1105	EPE	1	0
7	B	1103[A]	XUJ	1	0
3	B	1101	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/353 (97%)	-0.49	5 (1%) 72 76	6, 16, 35, 71	21 (6%)
2	B	189/189 (100%)	-0.63	2 (1%) 78 81	7, 14, 28, 40	8 (4%)
All	All	532/542 (98%)	-0.54	7 (1%) 75 79	6, 15, 33, 71	29 (5%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	375	THR	3.9
1	A	33	PRO	3.6
1	A	150[A]	ARG	3.0
1	A	374	TYR	2.4
1	A	34	ASP	2.3
2	B	431	ASN	2.2
2	B	533[A]	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	605	14/15	0.80	0.15	33,49,70,77	0
3	NAG	A	604	14/15	0.84	0.13	40,51,69,74	0
3	NAG	A	601	14/15	0.84	0.12	37,44,52,56	0
3	NAG	A	603	14/15	0.89	0.10	36,44,52,58	0
7	XUJ	B	1103[A]	10/10	0.90	0.10	13,13,16,18	10
6	EPE	B	1106	15/15	0.92	0.12	19,26,27,28	0
6	EPE	A	608	15/15	0.92	0.11	30,37,45,52	0
6	EPE	B	1105	15/15	0.93	0.11	22,31,45,55	0
3	NAG	A	602	14/15	0.95	0.07	20,22,32,40	0
3	NAG	B	1101	14/15	0.97	0.06	17,23,35,40	0
5	NA	A	607	1/1	0.99	0.06	28,28,28,28	0
8	XU7	B	1104[B]	10/10	0.99	0.04	13,13,16,16	0
4	CL	A	606	1/1	1.00	0.02	14,14,14,14	0
4	CL	B	1102	1/1	1.00	0.10	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.