



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:29 PM UTC

PDB ID : 1LEC / pdb_00001lec
Title : STRUCTURES OF THE LECTIN IV OF GRIFFONIA SIMPLICIFOLIA
AND ITS COMPLEX WITH THE LEWIS B HUMAN BLOOD GROUP DE-
TERMINANT AT 2.0 ANGSTROMS RESOLUTION
Authors : Delbaere, L.; Vandonseelaar, M.; Quail, J.
Deposited on : 1992-12-17
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	3% 66% 29% 5%
2	B	3	100%

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 2092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WEST-CENTRAL AFRICAN LEGUME LECTIN IV.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	0	0	0
			1904	1226	309	369			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	143	Total	O	0	0
			143	143		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: WEST-CENTRAL AFRICAN LEGUME LECTIN IV



• Molecule 2: alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.90Å 78.90Å 89.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.00 59.07 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 99.7 (59.07-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.00Å)	Xtriage
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.187 , (Not available) 0.179 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 102.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2092	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, SO4, FUC, MN, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	8/1953 (0.4%)	2.05	77/2670 (2.9%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	198	ILE	CA-CB	9.68	1.66	1.54
1	A	201	HIS	CD2-NE2	-7.56	1.29	1.37
1	A	233	HIS	CD2-NE2	-6.33	1.30	1.37
1	A	193	HIS	CD2-NE2	-6.06	1.31	1.37
1	A	175	HIS	CD2-NE2	-5.60	1.31	1.37
1	A	114	HIS	CD2-NE2	-5.20	1.32	1.37
1	A	175	HIS	CG-ND1	-5.17	1.32	1.38
1	A	215	ILE	CA-CB	5.11	1.62	1.55

All (77) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	163	ASN	OD1-CG-ND2	-11.65	110.95	122.60
1	A	108	PHE	CA-CB-CG	-9.91	103.89	113.80
1	A	221	VAL	O-C-N	-9.05	113.29	122.97
1	A	82	ASN	OD1-CG-ND2	-9.02	113.58	122.60
1	A	39	LYS	N-CA-C	8.09	122.25	110.28
1	A	135	ASN	OD1-CG-ND2	-8.03	114.58	122.60
1	A	37	LEU	N-CA-C	8.01	122.32	112.23
1	A	163	ASN	CB-CG-ND2	7.85	128.17	116.40
1	A	82	ASN	CB-CG-ND2	7.65	127.88	116.40
1	A	5	ASN	OD1-CG-ND2	-7.61	114.99	122.60
1	A	242	ASN	OD1-CG-ND2	-7.56	115.04	122.60
1	A	35	LEU	N-CA-C	-7.31	96.81	108.73
1	A	42	ALA	N-CA-C	7.21	120.98	111.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	211	GLN	OE1-CD-NE2	-7.08	115.52	122.60
1	A	53	GLN	OE1-CD-NE2	-7.06	115.54	122.60
1	A	151	ASN	OD1-CG-ND2	-7.05	115.55	122.60
1	A	192	GLU	N-CA-C	7.05	122.92	111.37
1	A	64	ASP	CA-CB-CG	6.96	119.56	112.60
1	A	213	VAL	N-CA-C	6.96	118.84	108.96
1	A	58	GLU	CA-CB-CG	6.95	127.99	114.10
1	A	41	ASP	CA-CB-CG	6.88	119.47	112.60
1	A	201	HIS	CB-CG-CD2	-6.79	122.37	131.20
1	A	30	ARG	O-C-N	-6.76	115.35	123.13
1	A	43	ASN	OD1-CG-ND2	-6.62	115.98	122.60
1	A	114	HIS	N-CA-C	6.51	119.21	111.33
1	A	224	ASP	N-CA-C	6.45	120.41	111.90
1	A	111	LEU	N-CA-C	6.42	121.40	113.50
1	A	133	TRP	N-CA-C	6.41	119.60	108.76
1	A	88	ALA	O-C-N	-6.40	116.10	123.33
1	A	54	ALA	O-C-N	-6.38	115.87	123.27
1	A	47	VAL	N-CA-CB	-6.37	101.53	110.26
1	A	13	SER	CA-C-O	6.33	130.99	122.51
1	A	228	TYR	O-C-N	-6.25	115.91	123.29
1	A	146	ILE	N-CA-C	-6.24	99.49	108.53
1	A	139	ASN	OD1-CG-ND2	-6.23	116.37	122.60
1	A	136	LYS	CA-C-N	6.22	128.90	120.38
1	A	136	LYS	C-N-CA	6.22	128.90	120.38
1	A	135	ASN	CB-CG-ND2	6.21	125.71	116.40
1	A	2	ASN	OD1-CG-ND2	-6.19	116.41	122.60
1	A	131	ASP	CA-CB-CG	6.15	118.75	112.60
1	A	159	THR	CA-CB-OG1	-6.15	100.37	109.60
1	A	15	SER	N-CA-C	5.96	120.71	113.38
1	A	241	THR	CA-C-O	5.96	129.03	120.51
1	A	215	ILE	O-C-N	-5.80	116.94	123.03
1	A	46	PRO	N-CA-C	5.75	119.89	111.03
1	A	104	ASP	N-CA-C	5.75	118.99	110.48
1	A	230	LEU	N-CA-C	5.72	120.33	113.23
1	A	115	GLU	N-CA-C	5.62	119.40	112.54
1	A	113	ARG	O-C-N	-5.60	116.66	123.10
1	A	93	PHE	N-CA-C	-5.52	100.70	109.76
1	A	123	ASN	OD1-CG-ND2	-5.44	117.16	122.60
1	A	93	PHE	O-C-N	-5.44	116.83	123.19
1	A	146	ILE	O-C-N	-5.38	117.65	123.14
1	A	192	GLU	CA-C-N	-5.38	114.15	123.25
1	A	192	GLU	C-N-CA	-5.38	114.15	123.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	151	ASN	CB-CG-ND2	5.37	124.46	116.40
1	A	154	VAL	N-CA-C	-5.35	99.39	107.73
1	A	117	ALA	N-CA-C	5.34	117.11	111.28
1	A	183	LYS	CB-CA-C	-5.33	104.72	111.86
1	A	235	PHE	O-C-N	-5.33	116.86	123.36
1	A	57	SER	N-CA-C	5.31	116.87	111.14
1	A	133	TRP	CE2-CD2-CE3	5.28	124.08	118.80
1	A	162	GLU	N-CA-C	5.27	117.87	110.23
1	A	114	HIS	CB-CG-CD2	-5.26	124.36	131.20
1	A	104	ASP	CA-CB-CG	5.22	117.82	112.60
1	A	211	GLN	CG-CD-NE2	5.22	124.23	116.40
1	A	21	GLU	CA-CB-CG	-5.20	103.71	114.10
1	A	230	LEU	CA-C-O	5.19	125.35	119.27
1	A	13	SER	CB-CA-C	-5.18	104.40	112.07
1	A	208	VAL	CA-C-O	5.18	123.46	118.90
1	A	70	ALA	N-CA-C	5.18	117.99	109.76
1	A	233	HIS	CB-CG-CD2	-5.17	124.47	131.20
1	A	133	TRP	CE2-CD2-CG	-5.10	101.08	107.20
1	A	12	TRP	CG-CD2-CE3	5.07	138.97	133.90
1	A	39	LYS	CG-CD-CE	-5.05	99.68	111.30
1	A	8	TYR	CA-C-N	5.04	124.65	119.56
1	A	8	TYR	C-N-CA	5.04	124.65	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1904	0	1811	11	0
2	B	38	0	34	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	5	0	0	0	0
6	A	143	0	0	0	0
All	All	2092	0	1845	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:LEU:HD23	1:A:243:LYS:HD2	1.55	0.85
1:A:243:LYS:HD3	1:A:243:LYS:C	2.30	0.56
1:A:14:TYR:HA	1:A:30:ARG:HD3	1.93	0.51
1:A:24:PHE:CZ	1:A:30:ARG:HB2	2.50	0.46
1:A:205:LEU:HG	1:A:209:LEU:HD12	2.00	0.43
1:A:115:GLU:HG2	1:A:116:THR:HG23	2.01	0.43
1:A:235:PHE:CE2	1:A:237:THR:HB	2.54	0.43
1:A:37:LEU:O	1:A:52:GLY:HA3	2.18	0.42
1:A:143:TYR:HB3	1:A:160:ARG:NH2	2.34	0.42
1:A:80:LEU:HD21	1:A:132:THR:HG21	2.02	0.40
1:A:97:PRO:HD3	1:A:124:GLN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/243 (99%)	230 (95%)	11 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/203 (100%)	192 (95%)	11 (5%)	20	17

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	SER
1	A	16	LEU
1	A	43	ASN
1	A	47	VAL
1	A	65	SER
1	A	91	LEU
1	A	113	ARG
1	A	137	ASP
1	A	192	GLU
1	A	205	LEU
1	A	238	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	163	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	7,8,9	0.75	0	9,10,12	1.81	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	OE-CD-N	3.56	132.66	124.96
1	A	1	PCA	OE-CD-CG	-3.07	121.24	126.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.94	1 (7%)	17,19,21	2.31	6 (35%)
2	FUC	B	2	2	10,10,11	0.86	0	14,14,16	1.23	2 (14%)
2	NAG	B	3	2	14,14,15	1.18	2 (14%)	17,19,21	1.28	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	FUC	B	2	2	-	-	0/1/1/1
2	NAG	B	3	2	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	NAG	C1-C2	-2.58	1.48	1.52
2	B	3	NAG	O5-C1	-2.18	1.40	1.43
2	B	1	NAG	C4-C3	-2.04	1.47	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C8-C7-N2	6.11	126.26	116.12
2	B	1	NAG	O7-C7-N2	-4.78	113.54	121.98
2	B	3	NAG	C4-C3-C2	-3.60	105.74	111.02
2	B	1	NAG	C4-C3-C2	-2.45	107.43	111.02
2	B	1	NAG	C1-O5-C5	2.43	115.45	112.19
2	B	2	FUC	O3-C3-C2	-2.36	105.23	110.05
2	B	1	NAG	O3-C3-C2	2.31	114.19	109.40
2	B	1	NAG	C1-C2-N2	-2.29	106.83	110.43
2	B	2	FUC	C2-C3-C4	2.17	114.68	110.86

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	500	-	4,4,4	0.40	0	6,6,6	0.30	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	242/243 (99%)	-0.18	8 (3%) 49 48	14, 24, 48, 85	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	TYR	3.3
1	A	13	SER	2.7
1	A	44	GLY	2.6
1	A	42	ALA	2.3
1	A	15	SER	2.2
1	A	242	ASN	2.1
1	A	138	TRP	2.1
1	A	243	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.48	0.17	83,91,98,103	0

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	1	14/15	0.86	0.13	30,40,53,57	0
2	NAG	B	3	14/15	0.87	0.10	38,42,55,56	0
2	FUC	B	2	10/11	0.93	0.07	30,32,36,39	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	A	500	5/5	0.59	0.35	47,49,52,52	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	A	251	1/1	0.99	0.02	21,21,21,21	0
3	MN	A	250	1/1	0.99	0.02	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.