



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:23 PM UTC

PDB ID : 4LEV / pdb_00004lev
Title : Structure of human cGAS
Authors : Li, P.
Deposited on : 2013-06-26
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	 12% 83% 15% ..
1	B	369	 11% 75% 21% .

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic GMP-AMP synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	369	3057	1955	525	561	10	6	0	3	0
1	B	369	3057	1954	525	561	10	7	0	3	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	154	PHE	-	expression tag	UNP Q8N884
A	155	GLU	-	expression tag	UNP Q8N884
A	156	LEU	-	expression tag	UNP Q8N884
B	154	PHE	-	expression tag	UNP Q8N884
B	155	GLU	-	expression tag	UNP Q8N884
B	156	LEU	-	expression tag	UNP Q8N884

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	241	Total	O	0	0
			241	241		
3	B	210	Total	O	0	0
			210	210		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cyclic GMP-AMP synthase



- Molecule 1: Cyclic GMP-AMP synthase



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	212.98Å 47.72Å 86.87Å 90.00° 113.89° 90.00°	Depositor
Resolution (Å)	38.45 – 1.95 38.45 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.8 (38.45-1.95) 93.6 (38.45-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.95Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.167 , 0.202 0.168 , 0.201	Depositor DCC
R_{free} test set	1999 reflections (3.48%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtrriage
Anisotropy	0.373	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6567	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/3114	0.99	8/4169 (0.2%)
1	B	0.58	0/3114	0.92	8/4168 (0.2%)
All	All	0.60	0/6228	0.95	16/8337 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	305	SER	CA-C-N	-16.73	99.04	120.79
1	A	305	SER	C-N-CA	-16.73	99.04	120.79
1	B	214	TYR	N-CA-C	-8.30	102.18	113.30
1	B	256	ASN	CA-C-N	7.30	128.97	119.84
1	B	256	ASN	C-N-CA	7.30	128.97	119.84
1	A	234	VAL	CA-C-N	5.98	127.31	119.84
1	A	234	VAL	C-N-CA	5.98	127.31	119.84
1	B	316	ILE	N-CA-C	5.93	116.55	107.77
1	B	260	ASN	CA-C-N	5.67	126.92	119.84
1	B	260	ASN	C-N-CA	5.67	126.92	119.84
1	A	214	TYR	CB-CA-C	-5.65	104.00	111.74
1	B	427	LYS	N-CA-C	-5.64	103.09	110.53
1	A	260	ASN	CA-C-N	5.56	126.79	119.84
1	A	260	ASN	C-N-CA	5.56	126.79	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	471	PHE	N-CA-C	-5.16	105.84	111.82
1	A	179	ILE	N-CA-C	5.12	116.21	111.81

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	305	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3057	0	3088	38	2
1	B	3057	0	3086	54	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	241	0	0	9	0
3	B	210	0	0	6	0
All	All	6567	0	6174	92	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:LYS:NZ	3:B:836:HOH:O	2.08	0.85
1:A:458:LYS:NZ	3:A:875:HOH:O	2.16	0.79
1:B:418:GLU:OE2	3:B:811:HOH:O	2.03	0.75
1:A:236:ARG:HB2	1:A:254:LYS:HD2	1.69	0.73
1:A:365:LYS:HB3	1:A:370:PHE:HB3	1.72	0.71
1:A:271:LEU:HD21	1:A:276:MSE:HE1	1.73	0.70
1:B:236:ARG:HB2	1:B:254:LYS:HG3	1.74	0.70
1:A:204:ARG:HH21	1:A:261:PRO:HB3	1.60	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:GLU:OE1	3:A:869:HOH:O	2.12	0.67
1:B:294:THR:OG1	1:B:295:ASP:N	2.22	0.67
1:A:157:ASP:OD2	3:A:837:HOH:O	2.13	0.67
1:B:193:LEU:HD11	1:B:310:LEU:HD11	1.78	0.66
1:A:507:GLN:NE2	3:A:818:HOH:O	2.21	0.66
1:B:191:ASP:OD1	1:B:214:TYR:OH	2.14	0.65
1:B:255:ARG:HD2	1:B:259:GLU:OE2	1.98	0.64
1:A:273:ALA:HB1	1:A:361:PRO:HB3	1.83	0.60
1:B:164:LYS:HE3	1:B:520:ASP:HB3	1.83	0.60
1:B:365:LYS:HB3	1:B:370:PHE:HB3	1.83	0.60
1:A:448:GLN:NE2	3:A:913:HOH:O	2.32	0.60
1:B:305:SER:OG	1:B:361:PRO:O	2.19	0.60
1:A:209:LEU:O	1:A:211:THR:N	2.37	0.58
1:B:168:VAL:HG21	1:B:519:PHE:HB3	1.85	0.58
1:A:423:ARG:NH1	3:A:818:HOH:O	2.37	0.57
1:B:194:LEU:HD11	1:B:208:LEU:HG	1.88	0.55
1:B:186:VAL:HG23	1:B:316:ILE:HD11	1.90	0.54
1:A:202:ALA:HA	1:A:261:PRO:HB2	1.89	0.54
1:A:305:SER:OG	1:A:361:PRO:O	2.23	0.54
1:B:288:ILE:HD11	1:B:310:LEU:HD22	1.89	0.54
1:B:240:GLU:OE1	1:B:252:LYS:NZ	2.25	0.53
1:A:427:LYS:O	1:A:429:HIS:N	2.42	0.53
1:A:339:ARG:NH1	3:A:825:HOH:O	2.41	0.52
1:B:273:ALA:HB1	1:B:361:PRO:HB3	1.90	0.52
1:A:281:ARG:HH12	1:A:298:MSE:HE3	1.74	0.52
1:B:257:PRO:O	1:B:259:GLU:N	2.43	0.51
1:A:176:ARG:HG3	1:A:176:ARG:HH11	1.75	0.51
1:A:192:HIS:NE2	1:A:287:GLU:OE2	2.43	0.50
1:B:428:LYS:HA	3:B:839:HOH:O	2.12	0.50
1:B:260:ASN:C	1:B:262:LEU:H	2.19	0.50
1:A:299:LYS:O	3:A:762:HOH:O	2.20	0.49
1:B:187:LYS:HA	1:B:214:TYR:CE1	2.48	0.49
1:B:363:HIS:CE1	1:B:372:GLU:HG3	2.47	0.49
1:B:178:ASP:O	1:B:181:THR:HG23	2.13	0.48
1:B:413[A]:MSE:SE	1:B:467:CYS:HB3	2.63	0.48
1:A:409:CYS:O	1:A:413[A]:MSE:HG3	2.14	0.48
1:B:196:ARG:HE	1:B:287:GLU:CD	2.22	0.48
1:A:176:ARG:HH22	1:A:220:ILE:HA	1.79	0.48
1:B:265:PHE:HZ	1:B:279:LYS:HD3	1.79	0.47
1:B:498:LYS:HE3	1:B:502:GLU:OE2	2.13	0.47
1:B:281:ARG:HG3	1:B:308:VAL:HG22	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:ARG:HG2	1:A:236:ARG:HH11	1.79	0.47
1:B:385:GLU:HG2	3:B:767:HOH:O	2.15	0.47
1:A:281:ARG:NH1	1:A:298:MSE:HE3	2.31	0.46
1:B:305:SER:HB2	1:B:362:LYS:HZ3	1.80	0.46
1:B:403:LYS:HG2	1:B:457:ARG:HD2	1.97	0.46
1:A:423:ARG:NH2	1:A:521:GLU:OE1	2.50	0.45
1:A:236:ARG:HG2	1:A:236:ARG:NH1	2.32	0.45
1:B:305:SER:HB2	1:B:362:LYS:NZ	2.30	0.45
1:A:413[A]:MSE:SE	1:A:467:CYS:HB3	2.66	0.45
1:A:176:ARG:HG3	1:A:176:ARG:NH1	2.32	0.44
1:B:181:THR:OG1	1:B:182:ALA:N	2.50	0.44
1:B:260:ASN:O	1:B:262:LEU:N	2.50	0.44
1:A:282:LYS:O	1:A:286:GLU:HG3	2.18	0.44
1:A:258:LYS:H	1:A:258:LYS:HG3	1.42	0.43
1:B:181:THR:O	1:B:185[A]:MSE:HG3	2.17	0.43
1:B:209:LEU:HG	1:B:210:ASN:N	2.34	0.43
1:A:307:ALA:HB2	3:A:849:HOH:O	2.18	0.43
1:B:186:VAL:CG2	1:B:316:ILE:HD11	2.49	0.43
1:B:197:LEU:HD12	1:B:197:LEU:HA	1.85	0.43
1:A:355:LYS:HE3	1:A:385:GLU:OE2	2.19	0.43
1:B:171:LYS:NZ	3:B:763:HOH:O	2.52	0.43
1:B:344:LEU:CD1	1:B:386:ILE:HD12	2.49	0.42
1:A:176:ARG:NH2	1:A:220:ILE:HD12	2.34	0.42
1:B:423:ARG:NH1	3:B:794:HOH:O	2.52	0.42
1:B:305:SER:OG	1:B:362:LYS:HG2	2.20	0.42
1:B:209:LEU:HD13	1:B:358:TYR:CZ	2.53	0.42
1:A:277:LEU:O	1:A:281:ARG:HG3	2.18	0.42
1:B:366:GLU:HB3	1:B:367:GLY:H	1.67	0.42
1:B:511:GLU:OE1	1:B:518:VAL:HB	2.20	0.42
1:A:403:LYS:HD2	1:A:403:LYS:HA	1.80	0.41
1:B:159:ALA:HA	1:B:160:PRO:HD3	1.90	0.41
1:B:271:LEU:HD21	1:B:276:MSE:HE1	2.01	0.41
1:B:462:LEU:HA	1:B:462:LEU:HD12	1.75	0.41
1:B:197:LEU:HB3	1:B:206:VAL:HG11	2.02	0.41
1:A:260:ASN:C	1:A:262:LEU:H	2.29	0.41
1:B:256:ASN:CB	1:B:257:PRO:HD2	2.51	0.41
1:A:179:ILE:O	1:A:181:THR:N	2.53	0.40
1:A:156:LEU:HD23	1:A:156:LEU:HA	1.94	0.40
1:B:168:VAL:CG2	1:B:519:PHE:HB3	2.51	0.40
1:B:190:VAL:HB	1:B:214:TYR:HE1	1.85	0.40
1:B:210:ASN:HD22	1:B:210:ASN:HA	1.70	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:288:ILE:C	1:B:290:ASP:H	2.30	0.40
1:B:344:LEU:HD11	1:B:386:ILE:HG23	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:GLU:O	1:A:506:LYS:NZ[4_655]	2.16	0.04
1:A:456:ASP:OD2	1:B:400:LYS:NZ[4_656]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	370/369 (100%)	348 (94%)	17 (5%)	5 (1%)	9 3
1	B	370/369 (100%)	342 (92%)	20 (5%)	8 (2%)	5 1
All	All	740/738 (100%)	690 (93%)	37 (5%)	13 (2%)	6 1

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	428	LYS
1	B	257	PRO
1	B	258	LYS
1	B	307	ALA
1	A	180	SER
1	A	210	ASN
1	A	216	GLU
1	A	345	SER
1	B	345	SER
1	B	366	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	304	GLY
1	B	306	PRO
1	B	261	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	342/334 (102%)	329 (96%)	13 (4%)	29 19
1	B	342/334 (102%)	321 (94%)	21 (6%)	17 7
All	All	684/668 (102%)	650 (95%)	34 (5%)	22 11

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	179	ILE
1	A	227	ASP
1	A	231	LYS
1	A	258	LYS
1	A	269	GLU
1	A	305	SER
1	A	311	LEU
1	A	363	HIS
1	A	365	LYS
1	A	371	GLN
1	A	423	ARG
1	A	427	LYS
1	A	469	THR
1	B	171	LYS
1	B	178	ASP
1	B	180	SER
1	B	181	THR
1	B	195	LEU
1	B	197	LEU
1	B	211	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	231	LYS
1	B	234	VAL
1	B	237	ILE
1	B	281	ARG
1	B	294	THR
1	B	297	ILE
1	B	316	ILE
1	B	317	SER
1	B	362	LYS
1	B	403	LYS
1	B	462	LEU
1	B	469	THR
1	B	492	SER
1	B	518	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	ASN
1	A	224	ASN
1	B	210	ASN
1	B	244	ASN
1	B	419	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	364/369 (98%)	0.39	43 (11%) 9 10	13, 40, 118, 165	2 (0%)
1	B	364/369 (98%)	0.51	42 (11%) 9 11	15, 46, 126, 174	1 (0%)
All	All	728/738 (98%)	0.45	85 (11%) 9 11	13, 42, 123, 174	3 (0%)

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	367	GLY	7.0
1	A	370	PHE	5.8
1	B	214	TYR	5.5
1	A	156	LEU	5.3
1	B	215	TYR	5.1
1	A	367	GLY	5.1
1	B	370	PHE	5.1
1	A	157	ASP	5.1
1	A	154	PHE	4.7
1	B	305	SER	4.7
1	A	305	SER	4.3
1	A	294	THR	4.3
1	A	257	PRO	4.2
1	B	522	PHE	4.0
1	A	179	ILE	4.0
1	A	368	ASN	4.0
1	B	296	VAL	4.0
1	A	301	LYS	3.9
1	B	175	SER	3.9
1	A	214	TYR	3.8
1	B	154	PHE	3.8
1	A	522	PHE	3.7
1	A	369	GLY	3.7
1	B	306	PRO	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	235	PRO	3.5
1	A	364	ALA	3.5
1	B	364	ALA	3.5
1	A	159	ALA	3.3
1	B	177	ASP	3.3
1	A	366	GLU	3.3
1	A	158	ALA	3.3
1	B	307	ALA	3.3
1	A	215	TYR	3.2
1	A	365	LYS	3.2
1	B	212	GLY	3.1
1	B	369	GLY	3.1
1	B	176	ARG	3.1
1	B	297	ILE	3.1
1	B	174	LEU	3.0
1	B	365	LYS	3.0
1	B	260	ASN	3.0
1	A	213	SER	2.9
1	B	261	PRO	2.8
1	A	182	ALA	2.8
1	A	314	GLU	2.8
1	B	520	ASP	2.8
1	B	213	SER	2.8
1	A	363	HIS	2.8
1	B	211	THR	2.8
1	A	211	THR	2.7
1	B	363	HIS	2.7
1	B	158	ALA	2.6
1	B	258	LYS	2.6
1	B	368	ASN	2.6
1	B	366	GLU	2.5
1	A	174	LEU	2.5
1	B	159	ALA	2.5
1	A	260	ASN	2.5
1	B	253	PHE	2.4
1	B	257	PRO	2.4
1	A	161	GLY	2.4
1	A	306	PRO	2.4
1	A	255	ARG	2.3
1	B	294	THR	2.3
1	A	210	ASN	2.3
1	A	177	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	181	THR	2.3
1	B	179	ILE	2.3
1	B	291	ILE	2.3
1	A	176	ARG	2.3
1	B	313	SER	2.2
1	A	212	GLY	2.2
1	B	210	ASN	2.2
1	A	371	GLN	2.1
1	B	304	GLY	2.1
1	A	216	GLU	2.1
1	A	521	GLU	2.1
1	B	372	GLU	2.1
1	B	195	LEU	2.1
1	A	253	PHE	2.1
1	A	262	LEU	2.1
1	A	303	GLY	2.0
1	B	256	ASN	2.0
1	A	160	PRO	2.0
1	B	161	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	601	1/1	0.98	0.04	23,23,23,23	0
2	ZN	B	601	1/1	0.99	0.02	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.