



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:40 PM UTC

PDB ID : 7LEE / pdb_00007lee
Title : HIV-1 Protease WT (NL4-3) in Complex with PU5 (LR4-47)
Authors : Lockbaum, G.J.; Rusere, L.N.; Henes, M.; Kosovrasti, K.; Lee, S.K.; Spielvogel, E.; Nalivaika, E.A.; Swannstrom, R.; KurtYilmaz, N.; Schiffer, C.A.; Ali, A.
Deposited on : 2021-01-14
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	99	4% 95% 5%
1	B	99	3% 91% 9%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3157 atoms, of which 1537 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protease.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	99	Total	C	H	N	O	S	0	0	0
			1476	472	746	123	131	4			
1	B	99	Total	C	H	N	O	S	0	0	0
			1469	470	743	123	129	4			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	7	LYS	GLN	engineered mutation	UNP Q8ULI9
B	7	LYS	GLN	engineered mutation	UNP Q8ULI9

- Molecule 2 is diethyl [(4-{(2S,3R)-4-[(1,3-benzothiazol-6-yl)sulfonyl][(2S)-2-methylbutyl]amino}-2-[(3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl]oxy}carbonyl)amino]-3-hydroxybutyl}phenoxy)methyl]phosphonate (CCD ID: XVJ) (formula: C₃₄H₄₈N₃O₁₁PS₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	S	0	0
			99	34	48	3	11	1	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	56	Total	O	0	0
			56	56		
4	B	52	Total	O	0	0
			52	52		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protease



- Molecule 1: Protease



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.33Å 59.12Å 61.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.71 – 1.79 42.71 – 1.79	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.71-1.79) 98.4 (42.71-1.79)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.191 , 0.231 0.200 , 0.240	Depositor DCC
R_{free} test set	907 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3157	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XVJ, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/743	0.62	0/1012
1	B	0.53	0/739	0.63	0/1007
All	All	0.53	0/1482	0.63	0/2019

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	730	746	747	4	0
1	B	726	743	743	8	0
2	A	51	48	0	2	0
3	A	5	0	0	0	0
4	A	56	0	0	1	0
4	B	52	0	0	1	0
All	All	1620	1537	1490	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:CYS:SG	4:A:242:HOH:O	2.51	0.68
1:A:95:CYS:SG	1:B:97:LEU:HD22	2.38	0.62
1:B:15:ILE:HD13	1:B:33:LEU:CD1	2.31	0.60
2:A:101:XVJ:O40	1:B:25:ASP:OD2	2.30	0.49
2:A:101:XVJ:O39	1:B:49:GLY:HA3	2.13	0.47
1:A:99:PHE:HB3	1:B:95:CYS:HA	1.98	0.46
1:A:64:ILE:HG21	1:A:89:LEU:HD13	1.98	0.45
1:B:30:ASP:HB3	4:B:116:HOH:O	2.17	0.43
1:B:15:ILE:HD13	1:B:33:LEU:HD12	2.02	0.41
1:B:63:LEU:HD12	1:B:71:ALA:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/99 (98%)	97 (100%)	0	0	100	100
1	B	97/99 (98%)	96 (99%)	1 (1%)	0	100	100
All	All	194/198 (98%)	193 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/83 (92%)	76 (100%)	0	100	100
1	B	75/83 (90%)	75 (100%)	0	100	100
All	All	151/166 (91%)	151 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	69	HIS
1	A	88	ASN
1	B	88	ASN
1	B	98	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	102	-	4,4,4	0.27	0	6,6,6	0.22	0
2	XVJ	A	101	-	55,55,55	6.58	36 (65%)	68,78,78	2.51	18 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XVJ	A	101	-	-	7/52/71/71	0/5/5/5

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	101	XVJ	O38-S50	19.09	1.63	1.43
2	A	101	XVJ	O39-S50	17.60	1.62	1.43
2	A	101	XVJ	C13-C12	-11.89	1.30	1.52
2	A	101	XVJ	C34-N37	10.45	1.43	1.29
2	A	101	XVJ	C34-S51	-10.45	1.58	1.73
2	A	101	XVJ	C01-N37	10.29	1.58	1.39
2	A	101	XVJ	C23-C24	9.61	1.54	1.38
2	A	101	XVJ	C06-C01	9.52	1.54	1.40
2	A	101	XVJ	O44-C15	9.31	1.60	1.41
2	A	101	XVJ	C23-C22	9.01	1.55	1.38
2	A	101	XVJ	S50-N35	8.94	1.76	1.63
2	A	101	XVJ	C20-C19	8.73	1.56	1.38
2	A	101	XVJ	O43-C13	8.69	1.62	1.43
2	A	101	XVJ	C21-C20	8.14	1.52	1.38
2	A	101	XVJ	C02-S51	-7.48	1.54	1.73
2	A	101	XVJ	C16-C14	-7.27	1.35	1.54
2	A	101	XVJ	C06-C05	6.95	1.50	1.38
2	A	101	XVJ	C05-C04	6.58	1.49	1.38
2	A	101	XVJ	C03-C04	6.54	1.50	1.39
2	A	101	XVJ	C21-C22	6.30	1.50	1.38
2	A	101	XVJ	O43-C15	-5.31	1.31	1.41
2	A	101	XVJ	C24-C19	5.21	1.49	1.38
2	A	101	XVJ	P49-O46	5.13	1.69	1.57
2	A	101	XVJ	C16-C17	5.07	1.64	1.51
2	A	101	XVJ	C11-N36	5.02	1.47	1.34
2	A	101	XVJ	C03-C02	4.96	1.47	1.39
2	A	101	XVJ	C04-S50	4.61	1.82	1.76
2	A	101	XVJ	C01-C02	4.18	1.46	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	101	XVJ	P49-O47	4.15	1.67	1.57
2	A	101	XVJ	O42-C11	3.99	1.42	1.35
2	A	101	XVJ	C08-C09	3.89	1.58	1.52
2	A	101	XVJ	P49-C29	3.72	1.90	1.80
2	A	101	XVJ	O44-C17	-3.43	1.33	1.43
2	A	101	XVJ	O40-C09	-2.92	1.37	1.43
2	A	101	XVJ	C18-C19	2.36	1.56	1.51
2	A	101	XVJ	C15-C14	2.33	1.56	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	101	XVJ	S51-C34-N37	-12.02	109.46	117.01
2	A	101	XVJ	O38-S50-N35	7.09	113.36	106.69
2	A	101	XVJ	C29-O45-C22	-5.87	109.71	117.85
2	A	101	XVJ	O39-S50-N35	-4.48	102.47	106.69
2	A	101	XVJ	C04-S50-N35	-4.11	102.32	107.28
2	A	101	XVJ	C17-C16-C14	4.04	108.90	103.38
2	A	101	XVJ	O38-S50-C04	-3.29	104.02	108.10
2	A	101	XVJ	O47-P49-O46	2.95	113.62	104.09
2	A	101	XVJ	C07-N35-S50	-2.89	111.01	117.32
2	A	101	XVJ	C19-C18-C10	-2.88	108.52	113.40
2	A	101	XVJ	C18-C10-N36	-2.82	106.08	110.08
2	A	101	XVJ	O39-S50-C04	2.76	111.53	108.10
2	A	101	XVJ	C26-C07-N35	-2.68	109.31	112.41
2	A	101	XVJ	C03-C02-C01	-2.47	120.15	121.99
2	A	101	XVJ	C12-O42-C11	-2.33	113.88	117.09
2	A	101	XVJ	C09-C10-N36	-2.23	105.78	109.97
2	A	101	XVJ	C06-C01-N37	2.12	133.53	129.62
2	A	101	XVJ	C05-C04-C03	2.07	123.24	120.68

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	101	XVJ	C30-O46-P49-O47
2	A	101	XVJ	C30-O46-P49-O48
2	A	101	XVJ	C21-C22-O45-C29
2	A	101	XVJ	C23-C22-O45-C29
2	A	101	XVJ	O45-C29-P49-O48
2	A	101	XVJ	C13-C12-O42-C11
2	A	101	XVJ	N35-C07-C26-C27

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	101	XVJ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/99 (100%)	0.40	4 (4%) 42 41	22, 33, 57, 81	0
1	B	99/99 (100%)	0.44	3 (3%) 52 52	21, 34, 57, 69	0
All	All	198/198 (100%)	0.42	7 (3%) 47 47	21, 34, 57, 81	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	99	PHE	4.1
1	B	99	PHE	3.8
1	A	1	PRO	2.8
1	B	15	ILE	2.7
1	B	16	GLY	2.6
1	A	68	GLY	2.5
1	A	10	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	102	5/5	0.75	0.11	54,65,88,113	0
2	XVJ	A	101	51/51	0.93	0.10	20,35,63,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.