



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:01 PM UTC

PDB ID : 7LES / pdb\_00007les  
Title : Acanthamoeba castellanii CYP51 (AcCYP51)-Imidazole complex  
Authors : Sharma, V.; Podust, L.M.  
Deposited on : 2021-01-15  
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	 88% 9% .
1	B	460	 86% 11% ..
1	C	460	 88% 10% .
1	D	460	 85% 11% .

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Obtusifoliol 14alphademethylase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	449	3594	2322	599	655	18	0	2	0
1	B	448	3571	2312	596	645	18	0	1	0
1	C	449	3577	2313	598	648	18	0	1	0
1	D	447	3550	2298	590	644	18	0	1	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	MET	-	initiating methionine	UNP L8GJB3
A	34	ALA	-	expression tag	UNP L8GJB3
A	35	LYS	-	expression tag	UNP L8GJB3
A	36	LYS	-	expression tag	UNP L8GJB3
A	37	THR	-	expression tag	UNP L8GJB3
A	38	SER	-	expression tag	UNP L8GJB3
A	39	SER	-	expression tag	UNP L8GJB3
A	40	LYS	-	expression tag	UNP L8GJB3
A	41	GLY	-	expression tag	UNP L8GJB3
A	42	LYS	-	expression tag	UNP L8GJB3
A	487	HIS	-	expression tag	UNP L8GJB3
A	488	HIS	-	expression tag	UNP L8GJB3
A	489	HIS	-	expression tag	UNP L8GJB3
A	490	HIS	-	expression tag	UNP L8GJB3
A	491	HIS	-	expression tag	UNP L8GJB3
A	492	HIS	-	expression tag	UNP L8GJB3
B	33	MET	-	initiating methionine	UNP L8GJB3
B	34	ALA	-	expression tag	UNP L8GJB3
B	35	LYS	-	expression tag	UNP L8GJB3
B	36	LYS	-	expression tag	UNP L8GJB3
B	37	THR	-	expression tag	UNP L8GJB3

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	38	SER	-	expression tag	UNP L8GJB3
B	39	SER	-	expression tag	UNP L8GJB3
B	40	LYS	-	expression tag	UNP L8GJB3
B	41	GLY	-	expression tag	UNP L8GJB3
B	42	LYS	-	expression tag	UNP L8GJB3
B	487	HIS	-	expression tag	UNP L8GJB3
B	488	HIS	-	expression tag	UNP L8GJB3
B	489	HIS	-	expression tag	UNP L8GJB3
B	490	HIS	-	expression tag	UNP L8GJB3
B	491	HIS	-	expression tag	UNP L8GJB3
B	492	HIS	-	expression tag	UNP L8GJB3
C	33	MET	-	initiating methionine	UNP L8GJB3
C	34	ALA	-	expression tag	UNP L8GJB3
C	35	LYS	-	expression tag	UNP L8GJB3
C	36	LYS	-	expression tag	UNP L8GJB3
C	37	THR	-	expression tag	UNP L8GJB3
C	38	SER	-	expression tag	UNP L8GJB3
C	39	SER	-	expression tag	UNP L8GJB3
C	40	LYS	-	expression tag	UNP L8GJB3
C	41	GLY	-	expression tag	UNP L8GJB3
C	42	LYS	-	expression tag	UNP L8GJB3
C	487	HIS	-	expression tag	UNP L8GJB3
C	488	HIS	-	expression tag	UNP L8GJB3
C	489	HIS	-	expression tag	UNP L8GJB3
C	490	HIS	-	expression tag	UNP L8GJB3
C	491	HIS	-	expression tag	UNP L8GJB3
C	492	HIS	-	expression tag	UNP L8GJB3
D	33	MET	-	initiating methionine	UNP L8GJB3
D	34	ALA	-	expression tag	UNP L8GJB3
D	35	LYS	-	expression tag	UNP L8GJB3
D	36	LYS	-	expression tag	UNP L8GJB3
D	37	THR	-	expression tag	UNP L8GJB3
D	38	SER	-	expression tag	UNP L8GJB3
D	39	SER	-	expression tag	UNP L8GJB3
D	40	LYS	-	expression tag	UNP L8GJB3
D	41	GLY	-	expression tag	UNP L8GJB3
D	42	LYS	-	expression tag	UNP L8GJB3
D	487	HIS	-	expression tag	UNP L8GJB3
D	488	HIS	-	expression tag	UNP L8GJB3
D	489	HIS	-	expression tag	UNP L8GJB3
D	490	HIS	-	expression tag	UNP L8GJB3
D	491	HIS	-	expression tag	UNP L8GJB3

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	492	HIS	-	expression tag	UNP L8GJB3

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0
3	B	1	Total C O 7 4 3	0	0
3	B	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0

- Molecule 4 is IMIDAZOLE (CCD ID: IMD) (formula:  $C_3H_5N_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		
4	B	1	Total	C	N	0	0
			5	3	2		
4	C	1	Total	C	N	0	0
			5	3	2		
4	D	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is FORMIC ACID (CCD ID: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0
5	C	1	Total C O 3 1 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	16	Total O 16 16	0	0
6	B	8	Total O 8 8	0	0
6	C	9	Total O 9 9	0	0
6	D	4	Total O 4 4	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.56Å 125.58Å 100.51Å 90.00° 90.04° 90.00°	Depositor
Resolution (Å)	100.71 – 2.65 100.51 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.9 (100.71-2.65) 99.8 (100.51-2.65)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 2.65Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.192 , 0.275 0.196 , 0.271	Depositor DCC
$R_{free}$ test set	3178 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	78.1	Xtrriage
Anisotropy	0.102	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for l,k,-h 0.459 for h,-k,-l 0.001 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14572	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	81.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, FMT, HEM, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/3684	0.99	0/4988
1	B	0.75	0/3660	1.02	4/4953 (0.1%)
1	C	0.75	0/3667	0.98	2/4965 (0.0%)
1	D	0.77	0/3638	1.00	5/4928 (0.1%)
All	All	0.76	0/14649	1.00	11/19834 (0.1%)

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	44	PRO	O-C-N	6.93	124.41	121.15
1	D	44	PRO	O-C-N	5.83	123.89	121.15
1	B	176	ILE	N-CA-CB	5.65	120.30	110.65
1	B	232	LYS	N-CA-C	-5.59	105.10	111.14
1	B	454	ASN	N-CA-C	5.48	118.18	111.82

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3594	0	3551	11	0
1	B	3571	0	3546	15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3577	0	3541	14	0
1	D	3550	0	3501	15	0
2	A	43	0	30	2	0
2	B	43	0	30	3	0
2	C	43	0	30	1	0
2	D	43	0	30	3	0
3	A	7	0	10	0	0
3	B	14	0	20	0	0
3	D	21	0	30	0	0
4	A	5	0	5	0	0
4	B	5	0	5	0	0
4	C	5	0	5	1	0
4	D	5	0	5	0	0
5	A	3	0	1	0	0
5	B	3	0	1	0	0
5	C	3	0	1	0	0
6	A	16	0	0	0	0
6	B	8	0	0	0	0
6	C	9	0	0	0	0
6	D	4	0	0	0	0
All	All	14572	0	14342	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 61 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:125:VAL:HG12	1:D:126:VAL:HG12	1.84	0.60
1:D:193:MET:HE1	1:D:292:LEU:HD11	1.83	0.60
1:B:109:SER:HB2	1:B:371:ILE:HD11	1.86	0.57
2:B:502:HEM:HBC2	2:B:502:HEM:HMC2	1.87	0.57
2:D:502:HEM:HMC2	2:D:502:HEM:HBC2	1.86	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/460 (98%)	427 (95%)	19 (4%)	3 (1%)	18	30
1	B	447/460 (97%)	422 (94%)	22 (5%)	3 (1%)	18	30
1	C	448/460 (97%)	415 (93%)	29 (6%)	4 (1%)	14	24
1	D	446/460 (97%)	416 (93%)	24 (5%)	6 (1%)	9	16
All	All	1790/1840 (97%)	1680 (94%)	94 (5%)	16 (1%)	14	24

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	126	VAL
1	D	78	ILE
1	D	257	GLU
1	D	277	GLN
1	A	296	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/397 (96%)	359 (94%)	23 (6%)	17	30
1	B	378/397 (95%)	356 (94%)	22 (6%)	18	31
1	C	380/397 (96%)	361 (95%)	19 (5%)	22	37
1	D	373/397 (94%)	349 (94%)	24 (6%)	16	28
All	All	1513/1588 (95%)	1425 (94%)	88 (6%)	18	31

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	375	SER
1	D	203	THR
1	C	431	ARG
1	D	82	LYS
1	D	330	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	200	ASN
1	D	66	GLN
1	C	314	ASN
1	D	110	GLN
1	B	200	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEG	A	502	-	6,6,6	0.64	0	5,5,5	0.45	0
3	PEG	D	505	-	6,6,6	0.60	0	5,5,5	0.38	0
4	IMD	D	504	2	5,5,5	0.67	0	5,5,5	0.52	0
3	PEG	D	501	-	6,6,6	0.58	0	5,5,5	0.58	0
3	PEG	D	503	-	6,6,6	0.47	0	5,5,5	0.14	0
4	IMD	C	502	2	5,5,5	0.57	0	5,5,5	0.66	0
3	PEG	B	501	-	6,6,6	0.54	0	5,5,5	0.66	0
2	HEM	B	502	4,1	50,50,50	1.54	6 (12%)	67,82,82	1.52	14 (20%)
5	FMT	C	503	-	2,2,2	0.70	0	1,1,1	0.74	0
2	HEM	A	501	4,1	50,50,50	1.54	6 (12%)	67,82,82	1.68	12 (17%)
5	FMT	A	504	-	2,2,2	0.75	0	1,1,1	0.69	0
3	PEG	B	503	-	6,6,6	0.54	0	5,5,5	0.22	0
5	FMT	B	505	-	2,2,2	0.71	0	1,1,1	0.69	0
4	IMD	A	503	2	5,5,5	0.59	0	5,5,5	0.46	0
2	HEM	D	502	4,1	50,50,50	1.55	7 (14%)	67,82,82	1.59	13 (19%)
2	HEM	C	501	4,1	50,50,50	1.53	6 (12%)	67,82,82	1.77	14 (20%)
4	IMD	B	504	2	5,5,5	0.51	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	502	-	-	2/4/4/4	-
3	PEG	D	505	-	-	3/4/4/4	-
4	IMD	D	504	2	-	-	0/1/1/1
3	PEG	D	501	-	-	2/4/4/4	-
3	PEG	D	503	-	-	2/4/4/4	-
4	IMD	C	502	2	-	-	0/1/1/1
3	PEG	B	501	-	-	2/4/4/4	-
2	HEM	B	502	4,1	-	0/14/54/54	-
2	HEM	A	501	4,1	-	0/14/54/54	-
3	PEG	B	503	-	-	3/4/4/4	-
4	IMD	A	503	2	-	-	0/1/1/1
2	HEM	D	502	4,1	-	0/14/54/54	-
2	HEM	C	501	4,1	-	0/14/54/54	-
4	IMD	B	504	2	-	-	0/1/1/1

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	C1B-NB	-4.79	1.32	1.40
2	C	501	HEM	FE-NC	4.78	2.10	1.95
2	C	501	HEM	C1B-NB	-4.74	1.32	1.40
2	A	501	HEM	FE-NC	4.47	2.09	1.95
2	D	502	HEM	FE-NB	4.39	2.08	1.94

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	HEM	CHC-C4B-NB	5.62	130.47	124.42
2	A	501	HEM	CHC-C4B-NB	5.27	130.09	124.42
2	A	501	HEM	CHD-C1D-ND	4.51	129.28	124.42
2	C	501	HEM	C1B-NB-C4B	4.49	110.52	105.21
2	A	501	HEM	C1B-NB-C4B	4.47	110.51	105.21

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	PEG	O1-C1-C2-O2
3	A	502	PEG	O2-C3-C4-O4
3	B	501	PEG	O1-C1-C2-O2
3	B	503	PEG	O2-C3-C4-O4
3	D	503	PEG	O1-C1-C2-O2


There are no ring outliers.


5 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	502	IMD	1	0
2	B	502	HEM	3	0
2	A	501	HEM	2	0
2	D	502	HEM	3	0
2	C	501	HEM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring


in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	449/460 (97%)	-1.21	0	100   100	34, 78, 110, 156	2 (0%)
1	B	448/460 (97%)	-1.18	0	100   100	41, 80, 117, 153	1 (0%)
1	C	449/460 (97%)	-1.18	0	100   100	35, 77, 110, 153	1 (0%)
1	D	447/460 (97%)	-1.18	1 (0%)	91   90	41, 80, 114, 168	1 (0%)
All	All	1793/1840 (97%)	-1.19	1 (0%)	92   91	34, 79, 113, 168	5 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	277	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PEG	D	501	7/7	0.98	0.10	89,91,98,101	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PEG	D	505	7/7	0.98	0.10	77,93,109,113	0
5	FMT	B	505	3/3	0.98	0.07	81,81,86,87	0
5	FMT	C	503	3/3	0.98	0.07	78,78,85,85	0
2	HEM	C	501	43/43	0.99	0.04	53,62,69,74	0
3	PEG	D	503	7/7	0.99	0.05	71,73,85,91	0
3	PEG	A	502	7/7	0.99	0.06	69,75,82,85	0
4	IMD	A	503	5/5	0.99	0.06	70,70,78,82	0
4	IMD	B	504	5/5	0.99	0.05	70,79,90,93	0
4	IMD	C	502	5/5	0.99	0.10	68,73,76,80	0
4	IMD	D	504	5/5	0.99	0.06	72,80,91,97	0
5	FMT	A	504	3/3	0.99	0.04	79,79,83,84	0
3	PEG	B	501	7/7	0.99	0.07	72,96,114,115	0
3	PEG	B	503	7/7	0.99	0.06	74,77,82,82	0
2	HEM	A	501	43/43	1.00	0.03	51,60,69,86	0
2	HEM	D	502	43/43	1.00	0.03	43,57,81,97	0
2	HEM	B	502	43/43	1.00	0.04	44,57,82,94	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around IMD B 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around IMD C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around IMD D 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM A 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM D 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.