



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 12:29 PM UTC

PDB ID : 6LFC / pdb_00006lfc
Title : E. coli Thioesterase I mutant DG
Authors : Deng, X.; Chen, L.; Yang, G.
Deposited on : 2019-12-01
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	 3% 71% 22% . . .
1	B	182	 2% 75% 16% . .
1	C	182	 3% 76% 15% . .
1	D	182	 2% 73% 21% . . .
1	E	182	 % 67% 25% 5% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	182	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into segments: a small red segment at the start, followed by a large green segment labeled '78%', then a yellow segment labeled '15%', and finally a small grey segment at the end. Above the bar, a '%' symbol is positioned. Below the bar, there are two small black dots.</p>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA thioesterase I also functions as protease I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	175	1382	883	242	253	4	0	0	0
1	B	174	1376	880	241	251	4	0	0	0
1	C	174	1376	880	241	251	4	0	0	0
1	D	176	1391	888	244	255	4	0	0	0
1	E	176	1391	888	244	255	4	0	0	0
1	F	174	1376	880	241	251	4	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP J7QCR7
A	141	LEU	MET	conflict	UNP J7QCR7
A	142	ASP	GLU	engineered mutation	UNP J7QCR7
A	145	GLY	TYR	engineered mutation	UNP J7QCR7
B	1	MET	-	expression tag	UNP J7QCR7
B	141	LEU	MET	conflict	UNP J7QCR7
B	142	ASP	GLU	engineered mutation	UNP J7QCR7
B	145	GLY	TYR	engineered mutation	UNP J7QCR7
C	1	MET	-	expression tag	UNP J7QCR7
C	141	LEU	MET	conflict	UNP J7QCR7
C	142	ASP	GLU	engineered mutation	UNP J7QCR7
C	145	GLY	TYR	engineered mutation	UNP J7QCR7
D	1	MET	-	expression tag	UNP J7QCR7
D	141	LEU	MET	conflict	UNP J7QCR7
D	142	ASP	GLU	engineered mutation	UNP J7QCR7
D	145	GLY	TYR	engineered mutation	UNP J7QCR7
E	1	MET	-	expression tag	UNP J7QCR7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	141	LEU	MET	conflict	UNP J7QCR7
E	142	ASP	GLU	engineered mutation	UNP J7QCR7
E	145	GLY	TYR	engineered mutation	UNP J7QCR7
F	1	MET	-	expression tag	UNP J7QCR7
F	141	LEU	MET	conflict	UNP J7QCR7
F	142	ASP	GLU	engineered mutation	UNP J7QCR7
F	145	GLY	TYR	engineered mutation	UNP J7QCR7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total O 4 4	0	0
2	B	6	Total O 6 6	0	0
2	C	3	Total O 3 3	0	0
2	D	3	Total O 3 3	0	0
2	E	3	Total O 3 3	0	0
2	F	4	Total O 4 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acyl-CoA thioesterase I also functions as protease I




- Molecule 1: Acyl-CoA thioesterase I also functions as protease I




- Molecule 1: Acyl-CoA thioesterase I also functions as protease I




- Molecule 1: Acyl-CoA thioesterase I also functions as protease I







- Molecule 1: Acyl-CoA thioesterase I also functions as protease I



- Molecule 1: Acyl-CoA thioesterase I also functions as protease I



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.93Å 118.23Å 121.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.00 – 2.70 63.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (63.00-2.70) 99.9 (63.00-2.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.69Å)	Xtrriage
Refinement program	REFMAC v1.0	Depositor
R, R_{free}	0.227 , 0.261 0.265 , 0.284	Depositor DCC
R_{free} test set	969 reflections (3.22%)	wwPDB-VP
Wilson B-factor (Å ²)	73.1	Xtrriage
Anisotropy	0.213	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.014 for -h,l,k	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8315	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/1415	0.97	2/1923 (0.1%)
1	B	0.58	0/1409	1.02	6/1915 (0.3%)
1	C	0.73	1/1409 (0.1%)	1.09	3/1915 (0.2%)
1	D	0.71	1/1425 (0.1%)	1.17	3/1938 (0.2%)
1	E	0.83	2/1425 (0.1%)	1.14	4/1938 (0.2%)
1	F	0.58	0/1409	0.93	1/1915 (0.1%)
All	All	0.68	4/8492 (0.0%)	1.06	19/11544 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	102	PRO	N-CA	17.75	1.70	1.47
1	C	147	LYS	C-N	12.92	1.50	1.33
1	D	157	HIS	C-N	12.72	1.50	1.33
1	E	16	ARG	CZ-NH2	-5.91	1.25	1.33

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	157	HIS	CA-C-N	16.26	136.51	119.76
1	D	157	HIS	C-N-CA	16.26	136.51	119.76
1	E	101	GLU	CA-C-N	15.34	139.02	119.84
1	E	101	GLU	C-N-CA	15.34	139.02	119.84
1	C	147	LYS	CA-C-N	14.11	134.80	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1382	0	1377	51	0
1	B	1376	0	1372	46	0
1	C	1376	0	1372	32	0
1	D	1391	0	1386	47	0
1	E	1391	0	1386	53	0
1	F	1376	0	1372	27	0
2	A	4	0	0	0	0
2	B	6	0	0	0	0
2	C	3	0	0	0	0
2	D	3	0	0	0	0
2	E	3	0	0	0	0
2	F	4	0	0	0	0
All	All	8315	0	8265	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:146:LEU:CD1	1:D:151:MET:HE3	1.40	1.49
1:E:102:PRO:N	1:E:102:PRO:CA	1.70	1.44
1:D:146:LEU:HD11	1:D:151:MET:CE	1.65	1.27
1:A:141:LEU:CD2	1:A:169:TRP:CG	2.30	1.14
1:A:141:LEU:HD23	1:A:169:TRP:CG	1.80	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/182 (94%)	165 (96%)	5 (3%)	1 (1%)	21	44
1	B	170/182 (93%)	165 (97%)	5 (3%)	0	100	100
1	C	170/182 (93%)	163 (96%)	6 (4%)	1 (1%)	21	44
1	D	174/182 (96%)	165 (95%)	5 (3%)	4 (2%)	5	13
1	E	174/182 (96%)	162 (93%)	11 (6%)	1 (1%)	21	44
1	F	170/182 (93%)	160 (94%)	9 (5%)	1 (1%)	21	44
All	All	1029/1092 (94%)	980 (95%)	41 (4%)	8 (1%)	16	37

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	32	GLN
1	F	142	ASP
1	A	140	PHE
1	C	143	GLU
1	D	32	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/153 (95%)	137 (94%)	9 (6%)	16	39
1	B	145/153 (95%)	139 (96%)	6 (4%)	27	56
1	C	145/153 (95%)	137 (94%)	8 (6%)	19	45
1	D	147/153 (96%)	140 (95%)	7 (5%)	23	50
1	E	147/153 (96%)	137 (93%)	10 (7%)	14	35
1	F	145/153 (95%)	137 (94%)	8 (6%)	19	45
All	All	875/918 (95%)	827 (94%)	48 (6%)	19	45

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	149	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	99	ASN
1	D	156	ILE
1	E	36	SER
1	E	146	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	157	HIS
1	F	90	GLN
1	D	90	GLN
1	F	118	ASN
1	F	48	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/182 (96%)	0.28	5 (2%) 53 50	36, 56, 73, 90	0
1	B	174/182 (95%)	0.16	4 (2%) 61 58	39, 57, 74, 91	0
1	C	174/182 (95%)	0.33	6 (3%) 48 44	30, 56, 80, 99	0
1	D	176/182 (96%)	0.19	3 (1%) 69 67	40, 57, 77, 96	0
1	E	176/182 (96%)	0.25	2 (1%) 78 76	40, 59, 75, 88	0
1	F	174/182 (95%)	0.26	2 (1%) 78 76	45, 58, 74, 86	0
All	All	1049/1092 (96%)	0.25	22 (2%) 63 61	30, 57, 76, 99	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	102	PRO	4.4
1	C	112	ASN	4.1
1	D	156	ILE	3.3
1	F	153	ASP	3.2
1	B	112	ASN	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.