



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:23 PM UTC

PDB ID : 7LFV / pdb_00007lfv
Title : Crystal structure of the SARS CoV-1 Papain-like protease in complex with peptide inhibitor VIR251
Authors : Olsen, S.K.; Lv, Z.
Deposited on : 2021-01-18
Resolution : 2.23 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	 2% 87% 10%
1	B	325	 10% 79% 17%
2	D	5	 80% 20%
2	F	5	 80% 20%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	2005	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called papain-like protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	312	2456	1558	409	472	17	0	0	0
1	A	314	2473	1570	412	474	17	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP P0C6U8
B	318	LEU	-	expression tag	UNP P0C6U8
B	319	GLU	-	expression tag	UNP P0C6U8
B	320	HIS	-	expression tag	UNP P0C6U8
B	321	HIS	-	expression tag	UNP P0C6U8
B	322	HIS	-	expression tag	UNP P0C6U8
B	323	HIS	-	expression tag	UNP P0C6U8
B	324	HIS	-	expression tag	UNP P0C6U8
B	325	HIS	-	expression tag	UNP P0C6U8
A	1	MET	-	initiating methionine	UNP P0C6U8
A	318	LEU	-	expression tag	UNP P0C6U8
A	319	GLU	-	expression tag	UNP P0C6U8
A	320	HIS	-	expression tag	UNP P0C6U8
A	321	HIS	-	expression tag	UNP P0C6U8
A	322	HIS	-	expression tag	UNP P0C6U8
A	323	HIS	-	expression tag	UNP P0C6U8
A	324	HIS	-	expression tag	UNP P0C6U8
A	325	HIS	-	expression tag	UNP P0C6U8

- Molecule 2 is a protein called Papain-like protease peptide inhibitor VIR251.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	5	34	22	5	7	0	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	F	5	34	22	5	7	0	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0

- Molecule 6 is AMMONIUM ION (CCD ID: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total N 1 1	0	0

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	74	Total O 74 74	0	0
8	A	46	Total O 46 46	0	0
8	F	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: papain-like protease




- Molecule 1: papain-like protease



- Molecule 2: Papain-like protease peptide inhibitor VIR251



- Molecule 2: Papain-like protease peptide inhibitor VIR251



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	103.61Å 103.61Å 265.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.34 – 2.23 53.34 – 2.23	Depositor EDS
% Data completeness (in resolution range)	99.9 (53.34-2.23) 95.1 (53.34-2.23)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 2.22Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.197 , 0.235 0.213 , 0.244	Depositor DCC
R_{free} test set	2000 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	51.2	Xtrriage
Anisotropy	0.134	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5221	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, SO4, CL, 73O, ACY, ZN, DPP, EDO, GVE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2529	0.47	0/3432
1	B	0.46	1/2511 (0.0%)	0.63	4/3408 (0.1%)
2	D	1.41	0/3	0.37	0/2
2	F	1.33	0/3	1.13	0/2
All	All	0.41	1/5046 (0.0%)	0.56	4/6844 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	76	LEU	CG-CD1	-5.16	1.35	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	76	LEU	CA-CB-CG	10.38	152.62	116.30
1	B	76	LEU	CA-C-N	6.20	131.02	122.77
1	B	76	LEU	C-N-CA	6.20	131.02	122.77
1	B	76	LEU	CB-CG-CD1	5.42	126.97	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2473	0	2417	25	0
1	B	2456	0	2397	36	1
2	D	34	0	12	0	0
2	F	34	0	11	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	36	0	54	14	0
4	B	48	0	72	6	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	46	0	0	4	0
8	B	74	0	0	1	0
8	F	1	0	0	0	0
All	All	5221	0	4963	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ASP:OD1	8:A:2101:HOH:O	1.89	0.89
1:A:234:TYR:OH	8:A:2102:HOH:O	2.00	0.78
1:A:292:THR:HG23	4:A:2005:EDO:H11	1.72	0.72
1:A:293:LYS:H	4:A:2005:EDO:H22	1.55	0.71
1:A:212:LEU:H	4:A:2009:EDO:H12	1.56	0.71

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:GLN:NE2	1:B:311:TYR:OH[9_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/325 (96%)	301 (96%)	11 (4%)	0	100	100
1	B	310/325 (95%)	301 (97%)	9 (3%)	0	100	100
2	D	1/5 (20%)	1 (100%)	0	0	100	100
2	F	1/5 (20%)	1 (100%)	0	0	100	100
All	All	624/660 (94%)	604 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/282 (96%)	271 (100%)	0	100	100
1	B	269/282 (95%)	269 (100%)	0	100	100
All	All	540/564 (96%)	540 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	74	HIS
1	B	172	HIS
1	A	147	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	195	GLN
1	A	237	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	73O	D	2	2	12,13,14	0.58	0	11,16,18	0.75	0
2	73O	F	2	2	12,13,14	0.75	0	11,16,18	0.61	0
2	GVE	F	5	1,2	7,7,7	1.04	0	7,7,7	1.74	3 (42%)
2	DPP	D	3	2	4,5,6	0.60	0	1,5,7	0.25	0
2	DPP	F	3	2	4,5,6	0.44	0	1,5,7	0.01	0
2	GVE	D	5	1,2	7,7,7	1.33	1 (14%)	7,7,7	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	73O	D	2	2	-	3/6/7/9	0/1/1/1
2	73O	F	2	2	-	1/6/7/9	0/1/1/1
2	GVE	F	5	1,2	-	4/6/6/6	-
2	DPP	D	3	2	-	0/2/4/6	-
2	DPP	F	3	2	-	0/2/4/6	-
2	GVE	D	5	1,2	-	5/6/6/6	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	5	GVE	OXT-C	2.51	1.40	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	5	GVE	CB-CG-C	2.78	123.89	113.69
2	F	5	GVE	C1-CB-CG	2.70	120.71	112.81
2	F	5	GVE	CH3-OXT-C	-2.22	108.94	116.07

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2	73O	C54-C52-CA-N
2	D	2	73O	C54-C52-CA-C
2	D	5	GVE	C1-CB-CG-C
2	F	5	GVE	C1-CB-CG-C
2	F	5	GVE	CG-C-OXT-CH3

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 3 are monoatomic and 1 is modelled with single atom - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	406	-	3,3,3	0.40	0	2,2,2	0.61	0
4	EDO	A	2003	-	3,3,3	0.46	0	2,2,2	0.35	0
4	EDO	A	2007	-	3,3,3	0.38	0	2,2,2	0.49	0
4	EDO	B	404	-	3,3,3	0.39	0	2,2,2	0.57	0
4	EDO	B	412	-	3,3,3	0.41	0	2,2,2	0.38	0
4	EDO	B	409	-	3,3,3	0.47	0	2,2,2	0.40	0
4	EDO	A	2010	-	3,3,3	0.40	0	2,2,2	0.30	0
4	EDO	B	413	-	3,3,3	0.52	0	2,2,2	0.22	0
4	EDO	B	405	-	3,3,3	0.52	0	2,2,2	0.37	0
4	EDO	B	411	-	3,3,3	0.59	0	2,2,2	0.29	0
4	EDO	B	410	-	3,3,3	0.45	0	2,2,2	0.38	0
4	EDO	B	403	-	3,3,3	0.63	0	2,2,2	0.45	0
5	SO4	B	414	-	4,4,4	0.24	0	6,6,6	0.18	0
4	EDO	A	2005	-	3,3,3	0.49	0	2,2,2	0.20	0
4	EDO	A	2001	-	3,3,3	0.45	0	2,2,2	0.46	0
4	EDO	A	2004	-	3,3,3	0.47	0	2,2,2	0.42	0
4	EDO	B	402	-	3,3,3	0.58	0	2,2,2	0.20	0
4	EDO	B	407	-	3,3,3	0.43	0	2,2,2	0.50	0
4	EDO	A	2008	-	3,3,3	0.40	0	2,2,2	0.39	0
4	EDO	A	2006	-	3,3,3	0.39	0	2,2,2	0.36	0
4	EDO	A	2009	-	3,3,3	0.49	0	2,2,2	0.25	0
5	SO4	A	2012	-	4,4,4	0.27	0	6,6,6	0.49	0
5	SO4	B	415	-	4,4,4	0.30	0	6,6,6	0.58	0
4	EDO	B	408	-	3,3,3	0.51	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	406	-	-	0/1/1/1	-
4	EDO	A	2003	-	-	0/1/1/1	-
4	EDO	A	2007	-	-	0/1/1/1	-
4	EDO	B	404	-	-	0/1/1/1	-
4	EDO	B	412	-	-	0/1/1/1	-
4	EDO	B	409	-	-	0/1/1/1	-
4	EDO	A	2010	-	-	1/1/1/1	-
4	EDO	B	413	-	-	1/1/1/1	-
4	EDO	B	405	-	-	1/1/1/1	-
4	EDO	B	411	-	-	1/1/1/1	-
4	EDO	B	410	-	-	0/1/1/1	-
4	EDO	B	403	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	2005	-	-	1/1/1/1	-
4	EDO	A	2001	-	-	0/1/1/1	-
4	EDO	A	2004	-	-	0/1/1/1	-
4	EDO	B	402	-	-	0/1/1/1	-
4	EDO	B	407	-	-	1/1/1/1	-
4	EDO	A	2008	-	-	0/1/1/1	-
4	EDO	A	2006	-	-	0/1/1/1	-
4	EDO	A	2009	-	-	0/1/1/1	-
4	EDO	B	408	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	405	EDO	O1-C1-C2-O2
4	A	2005	EDO	O1-C1-C2-O2
4	B	411	EDO	O1-C1-C2-O2
4	B	413	EDO	O1-C1-C2-O2
4	A	2010	EDO	O1-C1-C2-O2

There are no ring outliers.

11 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	404	EDO	1	0
4	B	412	EDO	1	0
4	B	409	EDO	1	0
4	A	2010	EDO	3	0
4	B	405	EDO	2	0
4	B	403	EDO	1	0
4	A	2005	EDO	5	0
4	A	2008	EDO	2	0
4	A	2006	EDO	3	0
4	A	2009	EDO	1	0
4	B	408	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/325 (96%)	0.10	5 (1%) 70 72	43, 62, 95, 147	0
1	B	312/325 (96%)	0.26	31 (9%) 13 12	38, 56, 132, 177	0
2	D	1/5 (20%)	1.02	0 100 100	63, 63, 63, 63	0
2	F	1/5 (20%)	-1.18	0 100 100	39, 39, 39, 39	0
All	All	628/660 (95%)	0.17	36 (5%) 29 27	38, 60, 112, 177	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	234	TYR	5.4
1	B	221	VAL	5.1
1	B	224	PRO	4.4
1	B	226	VAL	4.4
1	B	314	THR	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GVE	D	5	8/8	0.82	0.13	50,65,73,78	0
2	73O	D	2	13/14	0.84	0.16	69,90,107,112	0
2	GVE	F	5	8/8	0.92	0.12	43,47,60,76	0
2	DPP	D	3	6/7	0.93	0.08	66,70,72,74	0
2	DPP	F	3	6/7	0.96	0.06	46,48,49,49	0
2	73O	F	2	13/14	0.97	0.09	47,56,71,76	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	NH4	A	2011	1/1	0.69	0.23	77,77,77,77	0
4	EDO	B	402	4/4	0.72	0.20	75,75,76,76	0
4	EDO	B	410	4/4	0.74	0.18	71,75,75,76	0
4	EDO	B	408	4/4	0.74	0.14	83,87,87,88	0
4	EDO	A	2001	4/4	0.75	0.17	76,78,78,78	0
4	EDO	A	2003	4/4	0.77	0.15	69,73,74,79	0
4	EDO	A	2007	4/4	0.80	0.20	87,87,91,92	0
4	EDO	B	405	4/4	0.81	0.15	65,69,72,74	0
4	EDO	B	411	4/4	0.82	0.25	71,75,75,75	0
4	EDO	A	2010	4/4	0.83	0.16	86,87,88,88	0
4	EDO	A	2006	4/4	0.83	0.25	67,70,73,75	0
4	EDO	A	2004	4/4	0.85	0.16	71,75,79,79	0
4	EDO	A	2005	4/4	0.85	0.15	78,83,83,83	0
4	EDO	B	407	4/4	0.86	0.19	83,83,85,90	0
4	EDO	B	403	4/4	0.86	0.15	54,55,58,64	0
4	EDO	B	413	4/4	0.89	0.17	79,82,86,90	0
4	EDO	B	409	4/4	0.89	0.15	63,67,70,70	0
4	EDO	B	412	4/4	0.89	0.17	82,85,90,92	0
4	EDO	A	2009	4/4	0.90	0.10	78,79,82,82	0
4	EDO	A	2008	4/4	0.92	0.17	71,72,73,73	0
3	ZN	B	401	1/1	0.92	0.13	122,122,122,122	0
4	EDO	B	406	4/4	0.92	0.15	70,73,73,76	0
4	EDO	B	404	4/4	0.92	0.18	84,85,86,89	0
7	CL	A	2013	1/1	0.93	0.10	85,85,85,85	0
5	SO4	A	2012	5/5	0.94	0.07	73,74,76,82	0
5	SO4	B	414	5/5	0.96	0.06	66,72,74,77	0
5	SO4	B	415	5/5	0.97	0.07	56,59,65,65	0
3	ZN	A	2002	1/1	0.99	0.03	92,92,92,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.