



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 01:22 AM UTC

PDB ID : 9LFV / pdb\_00009lfv  
Title : Crystal structure of mouse RIP3 kinase domain complexed with LK01004  
Authors : Xie, H.; Su, H.X.; Li, M.J.; Xu, Y.C.  
Deposited on : 2025-01-09  
Resolution : 2.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	 2% (red), 75% (green), 20% (grey)
1	B	325	 3% (red), 76% (green), 21% (grey)
1	C	325	 8% (red), 74% (green), 23% (grey)
1	D	325	 11% (red), 73% (green), 6% (yellow), 22% (grey)

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor-interacting serine/threonine-protein kinase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	Total 1955	C 1261	N 329	O 357	S 8	0	0	0
1	B	256	Total 1916	C 1232	N 331	O 345	S 8	0	0	0
1	C	250	Total 1753	C 1123	N 302	O 320	S 8	0	0	0
1	D	255	Total 1772	C 1138	N 305	O 323	S 6	0	0	0

There are 60 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	111	ALA	CYS	conflict	UNP Q9QZL0
A	136	ASP	ASN	conflict	UNP Q9QZL0
A	194	LYS	ASP	conflict	UNP Q9QZL0
A	310	LEU	-	expression tag	UNP Q9QZL0
A	311	GLU	-	expression tag	UNP Q9QZL0
A	312	HIS	-	expression tag	UNP Q9QZL0
A	313	HIS	-	expression tag	UNP Q9QZL0
A	314	HIS	-	expression tag	UNP Q9QZL0
A	315	HIS	-	expression tag	UNP Q9QZL0
A	316	HIS	-	expression tag	UNP Q9QZL0
A	317	HIS	-	expression tag	UNP Q9QZL0
A	318	HIS	-	expression tag	UNP Q9QZL0
A	319	HIS	-	expression tag	UNP Q9QZL0
A	320	HIS	-	expression tag	UNP Q9QZL0
A	321	HIS	-	expression tag	UNP Q9QZL0
B	111	ALA	CYS	conflict	UNP Q9QZL0
B	136	ASP	ASN	conflict	UNP Q9QZL0
B	194	LYS	ASP	conflict	UNP Q9QZL0
B	310	LEU	-	expression tag	UNP Q9QZL0
B	311	GLU	-	expression tag	UNP Q9QZL0
B	312	HIS	-	expression tag	UNP Q9QZL0

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	313	HIS	-	expression tag	UNP Q9QZL0
B	314	HIS	-	expression tag	UNP Q9QZL0
B	315	HIS	-	expression tag	UNP Q9QZL0
B	316	HIS	-	expression tag	UNP Q9QZL0
B	317	HIS	-	expression tag	UNP Q9QZL0
B	318	HIS	-	expression tag	UNP Q9QZL0
B	319	HIS	-	expression tag	UNP Q9QZL0
B	320	HIS	-	expression tag	UNP Q9QZL0
B	321	HIS	-	expression tag	UNP Q9QZL0
C	111	ALA	CYS	conflict	UNP Q9QZL0
C	136	ASP	ASN	conflict	UNP Q9QZL0
C	194	LYS	ASP	conflict	UNP Q9QZL0
C	310	LEU	-	expression tag	UNP Q9QZL0
C	311	GLU	-	expression tag	UNP Q9QZL0
C	312	HIS	-	expression tag	UNP Q9QZL0
C	313	HIS	-	expression tag	UNP Q9QZL0
C	314	HIS	-	expression tag	UNP Q9QZL0
C	315	HIS	-	expression tag	UNP Q9QZL0
C	316	HIS	-	expression tag	UNP Q9QZL0
C	317	HIS	-	expression tag	UNP Q9QZL0
C	318	HIS	-	expression tag	UNP Q9QZL0
C	319	HIS	-	expression tag	UNP Q9QZL0
C	320	HIS	-	expression tag	UNP Q9QZL0
C	321	HIS	-	expression tag	UNP Q9QZL0
D	111	ALA	CYS	conflict	UNP Q9QZL0
D	136	ASP	ASN	conflict	UNP Q9QZL0
D	194	LYS	ASP	conflict	UNP Q9QZL0
D	310	LEU	-	expression tag	UNP Q9QZL0
D	311	GLU	-	expression tag	UNP Q9QZL0
D	312	HIS	-	expression tag	UNP Q9QZL0
D	313	HIS	-	expression tag	UNP Q9QZL0
D	314	HIS	-	expression tag	UNP Q9QZL0
D	315	HIS	-	expression tag	UNP Q9QZL0
D	316	HIS	-	expression tag	UNP Q9QZL0
D	317	HIS	-	expression tag	UNP Q9QZL0
D	318	HIS	-	expression tag	UNP Q9QZL0
D	319	HIS	-	expression tag	UNP Q9QZL0
D	320	HIS	-	expression tag	UNP Q9QZL0
D	321	HIS	-	expression tag	UNP Q9QZL0

- Molecule 2 is 3-[2-[(2 {S})-butan-2-yl]-1,3-benzothiazol-5-yl]-1- {tert}-butyl-pyrazolo[3,4-d]pyrimidin-4-amine (CCD ID: A1EJO) (formula: C<sub>20</sub>H<sub>24</sub>N<sub>6</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
2	A	1	27	20	6	1	0	1

- Molecule 3 is 3-[2-[(2 {R})-butan-2-yl]-1,3-benzothiazol-5-yl]-1- {tert}-butyl-pyrazolo[3,4-d]pyrimidin-4-amine (CCD ID: A1EJP) (formula: C<sub>20</sub>H<sub>24</sub>N<sub>6</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
3	A	1	27	20	6	1	0	1
3	B	1	27	20	6	1	0	0

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	S	0	0
			27	20	6	1		
3	D	1	Total	C	N	S	0	0
			27	20	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	O	0	0
			7	7		
4	B	5	Total	O	0	0
			5	5		
4	D	2	Total	O	0	0
			2	2		



Chain D:  11% 73% 6% 22%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.66Å 94.66Å 122.21Å 90.00° 90.98° 90.00°	Depositor
Resolution (Å)	61.10 – 2.78 61.10 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.0 (61.10-2.78) 99.0 (61.10-2.78)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.50 (at 2.77Å)	Xtrriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, $R_{free}$	0.247 , 0.280 0.248 , 0.285	Depositor DCC
$R_{free}$ test set	1523 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.9	Xtrriage
Anisotropy	0.344	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 73.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.013 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7545	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EJO, A1EJP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2002	0.61	0/2733
1	B	0.32	1/1960 (0.1%)	0.57	0/2676
1	C	0.29	0/1794	0.55	0/2465
1	D	0.28	0/1816	0.55	0/2501
All	All	0.31	1/7572 (0.0%)	0.57	0/10375

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	152	ASP	C-N	5.33	1.39	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	ASP	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	B	136	ASP	Peptide
1	C	136	ASP	Peptide
1	D	136	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1955	0	1892	6	0
1	B	1916	0	1857	4	0
1	C	1753	0	1567	4	0
1	D	1772	0	1569	9	0
2	A	27	0	0	0	0
3	A	27	0	0	0	0
3	B	27	0	0	0	0
3	C	27	0	0	0	0
3	D	27	0	0	1	0
4	A	7	0	0	0	0
4	B	5	0	0	0	0
4	D	2	0	0	0	0
All	All	7545	0	6885	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:120:ARG:HD3	1:D:295:VAL:HG11	1.87	0.57
1:C:120:ARG:HD3	1:C:295:VAL:HG11	1.86	0.56
1:A:113:ARG:HB3	1:A:118:LEU:HD11	1.88	0.54
1:C:116:PRO:HB2	1:C:295:VAL:HG22	1.89	0.54
1:D:35:VAL:HG21	1:D:88:VAL:HG11	1.95	0.47

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/325 (77%)	238 (95%)	12 (5%)	1 (0%)	30	57
1	B	248/325 (76%)	237 (96%)	10 (4%)	1 (0%)	30	57
1	C	240/325 (74%)	229 (95%)	10 (4%)	1 (0%)	30	57
1	D	247/325 (76%)	238 (96%)	8 (3%)	1 (0%)	30	57
All	All	986/1300 (76%)	942 (96%)	40 (4%)	4 (0%)	30	57

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	ASP
1	B	86	ASP
1	D	86	ASP
1	C	86	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/283 (72%)	202 (99%)	2 (1%)	68	86
1	B	198/283 (70%)	197 (100%)	1 (0%)	81	92
1	C	162/283 (57%)	160 (99%)	2 (1%)	63	84
1	D	160/283 (56%)	159 (99%)	1 (1%)	78	91
All	All	724/1132 (64%)	718 (99%)	6 (1%)	73	89

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	35	VAL
1	C	152	ASP
1	D	152	ASP
1	A	152	ASP
1	A	25	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	A1EJP	D	401	-	30,30,30	0.65	1 (3%)	40,46,46	1.40	9 (22%)
3	A1EJP	A	402[B]	-	30,30,30	0.73	1 (3%)	40,46,46	1.75	11 (27%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	A1EJP	B	401	-	30,30,30	0.69	1 (3%)	40,46,46	1.68	10 (25%)
3	A1EJP	C	401	-	30,30,30	0.63	1 (3%)	40,46,46	1.35	8 (20%)
2	A1EJO	A	401[A]	-	30,30,30	0.63	1 (3%)	40,46,46	1.31	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1EJP	D	401	-	-	4/16/16/16	0/4/4/4
3	A1EJP	A	402[B]	-	-	5/16/16/16	0/4/4/4
3	A1EJP	B	401	-	-	4/16/16/16	0/4/4/4
3	A1EJP	C	401	-	-	5/16/16/16	0/4/4/4
2	A1EJO	A	401[A]	-	-	6/16/16/16	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402[B]	A1EJP	C4-C5	-2.54	1.37	1.44
2	A	401[A]	A1EJO	C4-C5	-2.32	1.38	1.44
3	D	401	A1EJP	C4-C5	-2.19	1.38	1.44
3	C	401	A1EJP	C4-C5	-2.18	1.38	1.44
3	B	401	A1EJP	C4-C5	-2.13	1.38	1.44

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	A1EJP	C6-C5-N4	4.28	124.47	118.91
3	A	402[B]	A1EJP	C6-C5-N4	4.01	124.13	118.91
3	B	401	A1EJP	C4-C5-C6	-3.63	126.97	131.28
3	A	402[B]	A1EJP	C4-C5-C6	-3.61	126.99	131.28
3	A	402[B]	A1EJP	C11-C6-C5	-3.53	116.27	120.64

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	A1EJP	C16-C13-C14-C15
3	D	401	A1EJP	C16-C13-C14-C15

*Continued on next page...*

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	402[B]	A1EJP	N4-C5-C6-C7
3	C	401	A1EJP	N4-C5-C6-C7
3	B	401	A1EJP	N4-C5-C6-C7



There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	401	A1EJP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/325 (79%)	0.36	7 (2%) 56 48	33, 59, 88, 112	0
1	B	256/325 (78%)	0.53	9 (3%) 47 40	37, 64, 93, 109	0
1	C	250/325 (76%)	1.02	25 (10%) 12 10	52, 84, 113, 129	0
1	D	255/325 (78%)	1.19	37 (14%) 6 5	51, 89, 118, 137	0
All	All	1020/1300 (78%)	0.77	78 (7%) 20 16	33, 72, 110, 137	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	46	HIS	4.4
1	D	251	PRO	4.3
1	D	254	PRO	3.7
1	D	290	LEU	3.6
1	C	221	ALA	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	A1EJP	D	401	27/27	0.73	0.23	41,70,83,92	0
2	A1EJO	A	401[A]	27/27	0.78	0.27	44,51,55,57	27
3	A1EJP	B	401	27/27	0.79	0.19	26,55,66,73	0
3	A1EJP	A	402[B]	27/27	0.79	0.21	43,51,56,58	27
3	A1EJP	C	401	27/27	0.81	0.20	55,73,83,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around A1EJO A 401 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around A1EJP B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A1EJP A 402 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A1EJP C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.