



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:06 PM UTC

PDB ID : 3LH9 / pdb_00003lh9
Title : Crystal structure of mouse VPS26B(L197S/R199E) in spacegroup P41 21 2
Authors : Collins, B.; Shaw, D.; Norwood, S.
Deposited on : 2010-01-21
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	8% 63% 18% • 16%
1	B	340	6% 63% 17% • • 15%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein sorting-associated protein 26B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	1	0
			2376	1521	402	447	6			
1	B	288	Total	C	N	O	S	0	1	0
			2393	1532	405	450	6			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	MET	-	expression tag	UNP Q8C0E2
A	-2	GLY	-	expression tag	UNP Q8C0E2
A	-1	SER	-	expression tag	UNP Q8C0E2
A	0	HIS	-	expression tag	UNP Q8C0E2
A	1	HIS	-	expression tag	UNP Q8C0E2
A	2	HIS	-	expression tag	UNP Q8C0E2
A	3	HIS	-	expression tag	UNP Q8C0E2
A	4	HIS	-	expression tag	UNP Q8C0E2
A	5	HIS	-	expression tag	UNP Q8C0E2
A	6	MET	-	expression tag	UNP Q8C0E2
A	197	SER	LEU	engineered mutation	UNP Q8C0E2
A	199	GLU	ARG	engineered mutation	UNP Q8C0E2
B	-3	MET	-	expression tag	UNP Q8C0E2
B	-2	GLY	-	expression tag	UNP Q8C0E2
B	-1	SER	-	expression tag	UNP Q8C0E2
B	0	HIS	-	expression tag	UNP Q8C0E2
B	1	HIS	-	expression tag	UNP Q8C0E2
B	2	HIS	-	expression tag	UNP Q8C0E2
B	3	HIS	-	expression tag	UNP Q8C0E2
B	4	HIS	-	expression tag	UNP Q8C0E2
B	5	HIS	-	expression tag	UNP Q8C0E2
B	6	MET	-	expression tag	UNP Q8C0E2
B	197	SER	LEU	engineered mutation	UNP Q8C0E2
B	199	GLU	ARG	engineered mutation	UNP Q8C0E2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	162	Total 162	O 162	0	0
2	B	146	Total 146	O 146	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vacuolar protein sorting-associated protein 26B



- Molecule 1: Vacuolar protein sorting-associated protein 26B



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.35Å 118.35Å 214.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.54 – 2.40 42.54 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.54-2.40) 99.9 (42.54-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.39Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.199 , 0.228 0.198 , 0.224	Depositor DCC
R_{free} test set	3041 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	54.6	Xtrriage
Anisotropy	0.003	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5077	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2427	0.80	2/3265 (0.1%)
1	B	0.49	0/2445	0.86	6/3291 (0.2%)
All	All	0.50	0/4872	0.83	8/6556 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	56	ASN	CA-C-N	7.09	128.71	119.84
1	B	56	ASN	C-N-CA	7.09	128.71	119.84
1	A	179	ASN	N-CA-C	5.71	119.43	112.23
1	B	155	TYR	CA-C-N	5.59	126.83	119.84
1	B	155	TYR	C-N-CA	5.59	126.83	119.84
1	A	80	ARG	N-CA-C	5.43	117.65	111.02
1	B	124	VAL	CB-CA-C	-5.38	100.80	110.48
1	B	179	ASN	N-CA-C	5.14	118.70	112.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2376	0	2372	47	0
1	B	2393	0	2392	61	0
2	A	162	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	146	0	0	7	0
All	All	5077	0	4764	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:LEU:HD11	1:A:202:ILE:HD11	1.37	1.03
1:B:156:PRO:HB2	1:B:158:LEU:H	1.40	0.85
1:B:179:ASN:HD22	1:B:179:ASN:H	1.24	0.82
1:A:237:ALA:N	1:A:238:PRO:HD3	1.98	0.79
1:A:281:GLU:HG3	1:B:31:ASP:HB3	1.69	0.73
1:B:240:ARG:HD2	2:B:440:HOH:O	1.88	0.73
1:B:273:ASN:HD22	1:B:287:GLN:HB3	1.54	0.73
1:B:111:HIS:N	2:B:446:HOH:O	2.22	0.72
1:A:26:GLU:OE1	1:A:34:LYS:HE3	1.89	0.72
1:B:259:MET:HE2	1:B:266:PHE:HE1	1.54	0.71
1:B:192:LYS:HD2	1:B:194:TYR:CZ	2.26	0.71
1:B:259:MET:CE	1:B:266:PHE:HE1	2.06	0.69
1:A:80:ARG:O	2:A:391:HOH:O	2.12	0.68
1:A:99:ILE:HD12	1:A:103:GLN:OE1	1.95	0.67
1:A:172:LEU:HD11	1:A:202:ILE:CD1	2.20	0.67
1:B:259:MET:HE2	1:B:266:PHE:CE1	2.30	0.66
1:A:236:GLY:C	1:A:238:PRO:HD3	2.20	0.66
1:B:76:LEU:HD23	1:B:124:VAL:CG1	2.28	0.64
1:B:259:MET:CE	1:B:266:PHE:CE1	2.80	0.64
1:A:273:ASN:HD22	1:A:287:GLN:HB3	1.63	0.63
1:A:202:ILE:HD12	1:A:276:LEU:HD21	1.81	0.63
1:B:110:THR:HA	2:B:446:HOH:O	1.99	0.62
1:A:26:GLU:HG2	1:A:34:LYS:HG2	1.81	0.62
1:B:56:ASN:HB2	1:B:59:LYS:HB3	1.82	0.61
1:B:179:ASN:ND2	1:B:190:VAL:H	1.99	0.60
1:B:55:LYS:O	1:B:57:PRO:HD3	2.02	0.60
1:A:239:VAL:HG22	1:A:242:GLU:CD	2.27	0.59
1:B:259:MET:HE3	2:B:396:HOH:O	2.03	0.58
1:B:156:PRO:HB2	1:B:158:LEU:N	2.14	0.58
1:B:60:ARG:HG2	1:B:98:GLU:OE2	2.03	0.58
1:B:76:LEU:HD23	1:B:124:VAL:HG12	1.86	0.57
1:A:35:GLU:HG3	1:A:36:LYS:N	2.22	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:ASN:HD21	1:B:190:VAL:H	1.53	0.54
1:A:100:THR:O	1:A:101:GLN:HB2	2.07	0.54
1:A:23:LYS:HG3	1:A:39:LEU:HB3	1.90	0.54
1:A:237:ALA:N	1:A:238:PRO:CD	2.69	0.54
1:A:20:GLU:CD	1:A:20:GLU:H	2.16	0.53
1:B:60:ARG:HG3	1:B:100:THR:HG22	1.89	0.53
1:B:158:LEU:C	1:B:159:ASN:HD22	2.16	0.53
1:B:163:LYS:HE2	1:B:175:GLU:OE2	2.09	0.53
1:B:20:GLU:H	1:B:20:GLU:CD	2.16	0.52
1:B:156:PRO:HB2	1:B:157:GLU:HA	1.92	0.52
1:B:172:LEU:HG	1:B:276:LEU:HD21	1.90	0.52
1:A:10:VAL:HG21	1:A:61:LEU:HD22	1.90	0.52
1:B:192:LYS:HD3	1:B:243:SER:OG	2.10	0.52
1:B:69:GLU:HG2	1:B:90:VAL:HG22	1.92	0.52
1:A:172:LEU:HD21	1:A:202:ILE:HD13	1.94	0.49
1:B:156:PRO:CB	1:B:158:LEU:H	2.16	0.49
1:A:10:VAL:HG21	1:A:61:LEU:CD2	2.41	0.49
1:B:179:ASN:H	1:B:179:ASN:ND2	2.02	0.49
1:A:235:ASP:OD2	1:A:235:ASP:C	2.56	0.48
1:B:243:SER:HB2	2:B:423:HOH:O	2.14	0.48
1:B:56:ASN:HB2	1:B:59:LYS:CB	2.45	0.47
1:B:119:TYR:CG	1:B:259:MET:HE1	2.50	0.46
1:A:263:ASN:HD22	1:A:263:ASN:HA	1.58	0.46
1:B:174:ILE:HG12	1:B:205:MET:HE1	1.98	0.46
1:A:201:LYS:HB2	1:A:279:GLU:OE2	2.15	0.46
1:B:155:TYR:HB3	1:B:156:PRO:HD2	1.97	0.46
1:A:61:LEU:HB3	1:A:99:ILE:HB	1.98	0.45
1:B:12:VAL:O	1:B:143:LYS:HD2	2.16	0.45
1:A:174:ILE:HG22	1:A:175:GLU:N	2.31	0.45
1:A:208:ASP:HB2	1:A:273:ASN:HB3	1.98	0.45
1:A:29:THR:HB	2:A:344:HOH:O	2.15	0.45
1:B:13:GLU:HB3	1:B:15:LEU:HD21	1.98	0.45
1:B:29:THR:CG2	2:B:378:HOH:O	2.65	0.45
1:B:278:ASP:OD1	1:B:278:ASP:C	2.60	0.45
1:B:156:PRO:HB2	1:B:157:GLU:CA	2.47	0.45
1:B:59:LYS:HD3	1:B:60:ARG:O	2.17	0.45
1:B:217:THR:CG2	1:B:264:LYS:O	2.65	0.44
1:B:23:LYS:HB2	2:B:418:HOH:O	2.17	0.44
1:A:35:GLU:HG3	1:A:36:LYS:H	1.82	0.44
1:B:23:LYS:HE2	1:B:23:LYS:HB3	1.72	0.44
1:B:188:VAL:HG22	1:B:249:PHE:CD1	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:VAL:HB	1:B:274:LEU:HD23	2.00	0.44
1:B:33:LYS:H	1:B:33:LYS:HG2	1.52	0.43
1:B:91:LYS:HB2	1:B:91:LYS:HE2	1.75	0.42
1:A:122:GLN:HG2	1:A:224[A]:GLU:HG3	2.01	0.42
1:A:80:ARG:HB3	1:B:284:TYR:HD1	1.84	0.42
1:A:260:ARG:HD2	1:A:260:ARG:HA	1.86	0.42
1:A:56:ASN:C	1:A:58:ASN:N	2.75	0.42
1:A:28:LYS:HE3	1:A:28:LYS:HB3	1.78	0.42
1:B:189:ILE:HB	1:B:248:LEU:HB2	2.02	0.42
1:B:54:LEU:H	1:B:102:SER:HG	1.65	0.42
1:A:10:VAL:CG2	1:A:61:LEU:HD22	2.50	0.42
1:A:166:VAL:HG13	1:A:276:LEU:HD12	2.02	0.41
1:A:192:LYS:HE3	1:A:243:SER:OG	2.20	0.41
1:A:238:PRO:HA	1:A:242:GLU:OE1	2.20	0.41
1:A:126:LEU:C	1:A:126:LEU:HD23	2.45	0.41
1:A:163:LYS:HD3	1:A:175:GLU:OE2	2.20	0.41
1:B:184:HIS:HA	1:B:293:TRP:O	2.21	0.41
1:A:239:VAL:O	1:A:240:ARG:C	2.64	0.41
1:A:205:MET:HB3	1:A:205:MET:HE2	1.80	0.41
1:B:56:ASN:HB3	1:B:59:LYS:HB2	2.02	0.41
1:A:198:VAL:HG12	1:A:198:VAL:O	2.21	0.41
1:B:99:ILE:HD12	1:B:103:GLN:OE1	2.21	0.41
1:A:244:ILE:HA	1:A:245:PRO:HD3	1.87	0.41
1:B:77:TYR:N	1:B:77:TYR:CD1	2.89	0.40
1:B:119:TYR:CB	1:B:259:MET:HE1	2.51	0.40
1:B:143:LYS:HE3	1:B:143:LYS:HB2	1.93	0.40
1:B:214:THR:O	1:B:266:PHE:HA	2.21	0.40
1:A:75:GLU:OE2	1:A:127:ARG:HD3	2.22	0.40
1:A:91:LYS:HE2	1:A:91:LYS:HB2	1.92	0.40
1:A:195:PHE:HD2	1:A:242:GLU:O	2.04	0.40
1:A:137:ARG:HG3	2:A:476:HOH:O	2.21	0.40
1:B:20:GLU:CD	1:B:20:GLU:N	2.80	0.40
1:B:155:TYR:O	1:B:157:GLU:CG	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/340 (83%)	267 (94%)	13 (5%)	3 (1%)	11	18
1	B	287/340 (84%)	265 (92%)	15 (5%)	7 (2%)	4	5
All	All	570/680 (84%)	532 (93%)	28 (5%)	10 (2%)	6	9

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	156	PRO
1	B	31	ASP
1	A	101	GLN
1	B	57	PRO
1	B	160	SER
1	A	59	LYS
1	B	32	GLY
1	B	218	GLY
1	B	219	PRO
1	A	237	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/308 (85%)	240 (91%)	23 (9%)	9	16
1	B	265/308 (86%)	244 (92%)	21 (8%)	11	19
All	All	528/616 (86%)	484 (92%)	44 (8%)	10	17

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	A	10	VAL
1	A	11	GLU
1	A	20	GLU
1	A	49	LYS
1	A	51	SER
1	A	54	LEU
1	A	59	LYS
1	A	60	ARG
1	A	98	GLU
1	A	113	GLU
1	A	137	ARG
1	A	139	ASN
1	A	166	VAL
1	A	234	MET
1	A	235	ASP
1	A	254	GLU
1	A	261	ASP
1	A	263	ASN
1	A	276	LEU
1	A	286	LYS
1	A	288	GLN
1	A	297	ASP
1	B	10	VAL
1	B	21	SER
1	B	31	ASP
1	B	33	LYS
1	B	49	LYS
1	B	54	LEU
1	B	95	ARG
1	B	110	THR
1	B	113	GLU
1	B	124	VAL
1	B	143	LYS
1	B	147	ILE
1	B	154	THR
1	B	157	GLU
1	B	158	LEU
1	B	163	LYS
1	B	166	VAL
1	B	174	ILE
1	B	179	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	243	SER
1	B	261	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	HIS
1	A	223	HIS
1	A	263	ASN
1	A	273	ASN
1	B	111	HIS
1	B	123	ASN
1	B	159	ASN
1	B	179	ASN
1	B	223	HIS
1	B	273	ASN
1	B	288	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/340 (84%)	0.25	26 (9%) 15 12	26, 56, 96, 131	1 (0%)
1	B	288/340 (84%)	0.22	22 (7%) 20 16	27, 56, 95, 126	1 (0%)
All	All	574/680 (84%)	0.23	48 (8%) 17 14	26, 56, 96, 131	2 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	218	GLY	6.6
1	B	158	LEU	6.5
1	A	218	GLY	5.2
1	B	296	GLY	5.2
1	A	59	LYS	5.2
1	B	156	PRO	5.0
1	A	217	THR	4.8
1	A	156	PRO	4.5
1	A	32	GLY	4.2
1	A	297	ASP	4.1
1	A	234	MET	4.0
1	A	9	SER	3.9
1	B	55	LYS	3.9
1	B	219	PRO	3.9
1	A	219	PRO	3.7
1	A	231	TYR	3.5
1	A	57	PRO	3.5
1	B	9	SER	3.2
1	B	57	PRO	3.2
1	A	100	THR	3.2
1	B	217	THR	3.0
1	A	58	ASN	2.8
1	A	160	SER	2.8
1	B	216	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	99	ILE	2.7
1	B	59	LYS	2.7
1	B	60	ARG	2.7
1	A	28	LYS	2.6
1	B	234	MET	2.6
1	B	20	GLU	2.6
1	A	216	GLY	2.6
1	A	235	ASP	2.6
1	A	236	GLY	2.5
1	A	55	LYS	2.4
1	B	34	LYS	2.4
1	B	186	LYS	2.3
1	A	137	ARG	2.3
1	A	240	ARG	2.3
1	B	58	ASN	2.2
1	A	33	LYS	2.2
1	B	56	ASN	2.2
1	A	61	LEU	2.1
1	B	231	TYR	2.1
1	A	237	ALA	2.1
1	B	235	ASP	2.1
1	B	19	ALA	2.0
1	B	201	LYS	2.0
1	A	239	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.