



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:23 AM UTC

PDB ID : 3LHX / pdb_00003lhx
Title : Crystal structure of a Ketodeoxygluconokinase (kdgk) from *Shigella flexneri*
Authors : Ramagopal, U.A.; Toro, R.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-01-23
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	 6% 84% 10% 6%
1	B	319	 5% 85% 7% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ketodeoxygluconokinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	Se	0	1	0
			2335	1471	403	451	5	5			
1	B	295	Total	C	N	O	S	Se	0	1	0
			2288	1445	392	441	5	5			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	72	MSE	-	expression tag	UNP Q83J55
A	73	SER	-	expression tag	UNP Q83J55
A	74	LEU	-	expression tag	UNP Q83J55
A	383	GLU	-	expression tag	UNP Q83J55
A	384	GLY	-	expression tag	UNP Q83J55
A	385	HIS	-	expression tag	UNP Q83J55
A	386	HIS	-	expression tag	UNP Q83J55
A	387	HIS	-	expression tag	UNP Q83J55
A	388	HIS	-	expression tag	UNP Q83J55
A	389	HIS	-	expression tag	UNP Q83J55
A	390	HIS	-	expression tag	UNP Q83J55
B	76	MSE	-	expression tag	UNP Q83J55
B	77	SER	-	expression tag	UNP Q83J55
B	78	LEU	-	expression tag	UNP Q83J55
B	387	GLU	-	expression tag	UNP Q83J55
B	388	GLY	-	expression tag	UNP Q83J55
B	389	HIS	-	expression tag	UNP Q83J55
B	390	HIS	-	expression tag	UNP Q83J55
B	391	HIS	-	expression tag	UNP Q83J55
B	392	HIS	-	expression tag	UNP Q83J55
B	393	HIS	-	expression tag	UNP Q83J55
B	394	HIS	-	expression tag	UNP Q83J55


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	87	Total 87	O 87	0	0
2	B	95	Total 95	O 95	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ketodeoxygluconokinase



• Molecule 1: Ketodeoxygluconokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.87Å 169.01Å 44.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.51 – 1.87 84.51 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.0 (84.51-1.87) 99.0 (84.51-1.87)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.87Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.215 , 0.259 0.221 , 0.260	Depositor DCC
R_{free} test set	2785 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 27.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4805	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	1/2370 (0.0%)	0.93	2/3204 (0.1%)
1	B	1.03	3/2323 (0.1%)	1.01	3/3142 (0.1%)
All	All	1.01	4/4693 (0.1%)	0.97	5/6346 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	153	MSE	SE-CE	-6.11	1.77	1.95
1	B	157	MSE	SE-CE	-5.95	1.77	1.95
1	B	323	VAL	CA-CB	5.26	1.58	1.54
1	B	82	ILE	CA-CB	5.15	1.60	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	328	LEU	CA-C-N	-6.60	113.18	119.85
1	B	328	LEU	C-N-CA	-6.60	113.18	119.85
1	A	346	VAL	N-CA-C	5.96	116.61	110.82
1	B	339	ALA	N-CA-C	-5.43	105.26	111.07
1	A	129	THR	N-CA-C	-5.24	106.28	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2335	0	2312	13	0
1	B	2288	0	2264	16	0
2	A	87	0	0	0	0
2	B	95	0	0	0	0
All	All	4805	0	4576	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:326:VAL:HG12	1:B:326:VAL:O	1.70	0.92
1:A:291:HIS:HE1	1:A:310:SER:OG	1.67	0.76
1:B:138:GLN:OE1	1:B:156:ARG:NH1	2.21	0.73
1:B:295:HIS:HE1	1:B:314:SER:OG	1.72	0.72
1:A:128:GLY:O	1:A:152:ARG:HD2	1.93	0.67
1:A:130:ASP:OD2	1:A:132:PHE:HB2	1.97	0.65
1:B:336:THR:HG22	1:B:336:THR:O	2.00	0.61
1:A:325:PRO:HD2	1:A:328:LYS:HD2	1.84	0.59
1:B:326:VAL:O	1:B:326:VAL:CG1	2.39	0.59
1:A:134:GLN:OE1	1:A:152:ARG:NH1	2.35	0.59
1:B:375:ARG:CG	1:B:375:ARG:HH21	2.16	0.59
1:A:330:ILE:HD12	1:A:369:GLN:O	2.03	0.58
1:B:375:ARG:HH21	1:B:375:ARG:HB2	1.68	0.58
1:A:291:HIS:HD2	1:A:295:VAL:O	1.89	0.56
1:B:375:ARG:HH21	1:B:375:ARG:CB	2.21	0.54
1:A:380:MSE:HE3	1:A:381:PRO:HD2	1.91	0.53
1:A:325:PRO:HG2	1:A:328:LYS:HE2	1.91	0.52
1:B:375:ARG:HH21	1:B:375:ARG:HG2	1.76	0.50
1:A:128:GLY:O	1:A:152:ARG:CD	2.62	0.47
1:B:326:VAL:HG23	1:B:365:HIS:HB3	1.97	0.47
1:B:295:HIS:HD2	1:B:299:VAL:O	1.99	0.46
1:A:247:TRP:CG	1:A:253:THR:HG1	2.33	0.46
1:A:370:TYR:HB2	1:A:375:ILE:HG22	1.99	0.45
1:B:244:ASN:O	1:B:245:ASN:C	2.60	0.43
1:A:339:PHE:C	1:A:339:PHE:CD1	2.96	0.43
1:B:167:ILE:HG12	1:B:177:PHE:CE2	2.54	0.42
1:B:305:LYS:HD3	1:B:343:PHE:CD1	2.53	0.42
1:B:91:GLU:HB2	1:B:165:TYR:CE2	2.55	0.42
1:B:132:GLY:O	1:B:156:ARG:HD2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/319 (92%)	289 (98%)	5 (2%)	1 (0%)	36	26
1	B	290/319 (91%)	286 (99%)	3 (1%)	1 (0%)	36	26
All	All	585/638 (92%)	575 (98%)	8 (1%)	2 (0%)	36	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	ASN
1	B	245	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/254 (97%)	238 (97%)	8 (3%)	33	16
1	B	240/254 (94%)	236 (98%)	4 (2%)	53	40
All	All	486/508 (96%)	474 (98%)	12 (2%)	42	26

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	120	THR
1	A	156	ARG
1	A	165	THR
1	A	206	LEU
1	A	292	ASN
1	A	305	ASP
1	A	374	ILE
1	B	156	ARG
1	B	210	LEU
1	B	334	ILE
1	B	375	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	122	HIS
1	A	254	GLN
1	A	291	HIS
1	A	292	ASN
1	A	369	GLN
1	B	109	ASN
1	B	126	HIS
1	B	295	HIS
1	B	296	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/319 (92%)	0.52	18 (6%) 27 29	17, 34, 53, 63	1 (0%)
1	B	290/319 (90%)	0.35	15 (5%) 33 34	18, 31, 47, 56	1 (0%)
All	All	585/638 (91%)	0.43	33 (5%) 30 32	17, 32, 50, 63	2 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	385	PRO	4.6
1	B	99	VAL	4.2
1	A	129	THR	3.5
1	B	337	THR	3.4
1	B	383	ALA	3.4
1	B	93	SER	3.1
1	A	94	ASP	3.1
1	A	381	PRO	3.0
1	A	322	VAL	2.9
1	B	166	TYR	2.8
1	A	165	THR	2.7
1	B	176	THR	2.7
1	A	187	ALA	2.6
1	A	231	ALA	2.6
1	A	333	THR	2.6
1	B	252	ALA	2.5
1	A	95	VAL	2.5
1	B	376	GLY	2.5
1	B	374	TYR	2.4
1	A	173	PHE	2.4
1	B	333	VAL	2.4
1	A	370	TYR	2.3
1	A	170	GLU	2.3
1	A	178	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	161	TYR	2.3
1	B	326	VAL	2.2
1	B	167	ILE	2.2
1	A	321	ALA	2.2
1	A	171	ARG	2.2
1	A	154	GLU	2.1
1	B	338	ALA	2.1
1	B	177	PHE	2.1
1	A	141	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.