



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:39 PM UTC

PDB ID : 6LII / pdb_00006lii
Title : A quinone oxidoreductase
Authors : Hakoshima, T.; Kim, S.-Y.; Mori, T.
Deposited on : 2019-12-11
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	 81% 16% ..
1	B	353	 76% 20% .
1	C	353	 79% 16% ..
1	D	353	 79% 17% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Synaptic vesicle membrane protein VAT-1 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	0	0
			2609	1671	449	471	18			
1	B	340	Total	C	N	O	S	0	0	0
			2552	1629	441	465	17			
1	C	345	Total	C	N	O	S	0	0	0
			2582	1658	443	463	18			
1	D	342	Total	C	N	O	S	0	0	0
			2577	1650	442	468	17			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	GLY	-	expression tag	UNP Q99536
A	42	PRO	-	expression tag	UNP Q99536
B	41	GLY	-	expression tag	UNP Q99536
B	42	PRO	-	expression tag	UNP Q99536
C	41	GLY	-	expression tag	UNP Q99536
C	42	PRO	-	expression tag	UNP Q99536
D	41	GLY	-	expression tag	UNP Q99536
D	42	PRO	-	expression tag	UNP Q99536

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	56	Total	O	0	0
			56	56		
2	B	53	Total	O	0	0
			53	53		
2	C	58	Total	O	0	0
			58	58		
2	D	58	Total	O	0	0
			58	58		

GLY	PRO	ALA	SER	P45	L51	Y59	D60	P71	P72	G77	Q78	L79	R82	R96	Y100	D101	R102	L103	P110	I119	A120	V121	G122	E123	R128	K129	M135	R139	W143	Q144	E145	I157	P158	M161	A166	L169	L170	Y173	L174	M178
V192	L193	M196	A197	A198	M203	L208	T211	L229	N232	G233	V234	Y245	I249	L259	K271	L276	V282	P284	K285	R296	N297	L298	MET	ALA	LEU	A302	N307	L337	V341	W362	A370	M371	K372	Q373	M374	Q375	E376	K377	V380	
G381	K382	L385	P389	GLU	LYS	GLU	ASN																																	

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	69.35Å 69.35Å 581.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.68 – 2.30 48.68 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.68-2.30) 99.3 (48.68-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.60 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.16_3549, PHENIX 1.16_3549	Depositor
R, R_{free}	0.180 , 0.203 0.180 , 0.199	Depositor DCC
R_{free} test set	3734 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	43.3	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.469 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10545	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	5/2667 (0.2%)	0.39	0/3625
1	B	0.27	0/2605	0.36	0/3538
1	C	0.66	10/2640 (0.4%)	0.55	0/3592
1	D	0.31	2/2634 (0.1%)	0.35	0/3582
All	All	0.45	17/10546 (0.2%)	0.42	0/14337

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	294	PRO	C-O	-8.32	1.13	1.24
1	C	356	PRO	C-O	-6.89	1.14	1.23
1	A	49	ARG	C-O	-6.63	1.16	1.23
1	C	354	ILE	C-O	-6.57	1.17	1.24
1	C	96	ARG	C-O	-6.49	1.16	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2609	0	2659	41	0
1	B	2552	0	2591	55	0
1	C	2582	0	2624	47	0
1	D	2577	0	2610	44	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	56	0	0	1	0
2	B	53	0	0	0	0
2	C	58	0	0	1	0
2	D	58	0	0	6	0
All	All	10545	0	10484	182	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:121:VAL:CG1	1:B:125:VAL:HB	1.93	0.97
1:B:121:VAL:HG11	1:B:125:VAL:CG1	1.94	0.97
1:A:240:TYR:HE2	1:A:241:HIS:CE1	1.95	0.84
1:D:196:MET:CE	1:D:377:LYS:HE3	2.07	0.84
1:B:121:VAL:CG1	1:B:125:VAL:CB	2.57	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/353 (97%)	338 (98%)	5 (2%)	0	100	100
1	B	336/353 (95%)	330 (98%)	6 (2%)	0	100	100
1	C	343/353 (97%)	334 (97%)	9 (3%)	0	100	100
1	D	338/353 (96%)	331 (98%)	7 (2%)	0	100	100
All	All	1360/1412 (96%)	1333 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/284 (98%)	277 (100%)	1 (0%)	84	92
1	B	271/284 (95%)	270 (100%)	1 (0%)	84	92
1	C	271/284 (95%)	269 (99%)	2 (1%)	76	87
1	D	273/284 (96%)	272 (100%)	1 (0%)	84	92
All	All	1093/1136 (96%)	1088 (100%)	5 (0%)	81	90

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	240	TYR
1	B	135	MET
1	C	355	LYS
1	C	377	LYS
1	D	79	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	327	HIS
1	D	308	GLN
1	C	153	GLN
1	D	350	ASN
1	D	214	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/353 (97%)	-1.47	0 100 100	24, 39, 55, 90	0
1	B	340/353 (96%)	-1.36	0 100 100	26, 46, 75, 117	0
1	C	345/353 (97%)	-1.43	0 100 100	26, 41, 65, 146	0
1	D	342/353 (96%)	-1.37	0 100 100	30, 44, 66, 95	0
All	All	1372/1412 (97%)	-1.41	0 100 100	24, 43, 67, 146	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.