



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 06:01 AM UTC

PDB ID : 9LIK / pdb_00009lik
Title : Crystal structure of the C-tagged I7L protease from monkeypox virus
Authors : You, M.Y.; Wang, Y.H.; Su, H.X.; Li, M.J.; Xu, Y.C.
Deposited on : 2025-01-14
Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2053 (2.66-2.62)
Clashscore	190562	2097 (2.66-2.62)
Ramachandran outliers	187476	2066 (2.66-2.62)
Sidechain outliers	187428	2066 (2.66-2.62)
RSRZ outliers	180081	2052 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	22% 91% 9%
1	B	429	7% 85% 6% 9%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called I7L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	392	Total	C	N	O	S	0	0	0
			3087	1994	495	575	23			
1	A	427	Total	C	N	O	S	0	0	0
			3264	2093	532	613	26			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	424	GLU	-	expression tag	UNP Q8V512
B	425	ASN	-	expression tag	UNP Q8V512
B	426	LEU	-	expression tag	UNP Q8V512
B	427	TYR	-	expression tag	UNP Q8V512
B	428	PHE	-	expression tag	UNP Q8V512
B	429	GLN	-	expression tag	UNP Q8V512
A	424	GLU	-	expression tag	UNP Q8V512
A	425	ASN	-	expression tag	UNP Q8V512
A	426	LEU	-	expression tag	UNP Q8V512
A	427	TYR	-	expression tag	UNP Q8V512
A	428	PHE	-	expression tag	UNP Q8V512
A	429	GLN	-	expression tag	UNP Q8V512


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	4	Total	O	0	0
			4	4		
2	A	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: I7L



• Molecule 1: I7L



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	187.16Å 187.16Å 69.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.95 – 2.64 44.95 – 2.64	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.95-2.64) 99.5 (44.95-2.64)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.65Å)	Xtriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, R_{free}	0.242 , 0.270 (Not available) , 0.261	Depositor DCC
R_{free} test set	2060 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	67.3	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6361	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3338	0.62	0/4523
1	B	0.31	0/3160	0.59	0/4277
All	All	0.31	0/6498	0.61	0/8800

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3264	0	3036	21	0
1	B	3087	0	2937	18	0
2	A	6	0	0	0	0
2	B	4	0	0	0	0
All	All	6361	0	5973	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:ASP:HA	1:A:145:ARG:HE	1.56	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:TYR:HA	1:B:180:GLN:HG2	1.77	0.67
1:B:119:PHE:O	1:B:180:GLN:NE2	2.33	0.62
1:B:206:LYS:NZ	1:B:293:ASN:OD1	2.33	0.62
1:A:198:LEU:O	1:A:206:LYS:NZ	2.35	0.60
1:A:119:PHE:O	1:A:180:GLN:NE2	2.36	0.58
1:B:411:LEU:HD23	1:A:404:ILE:HG21	1.89	0.54
1:B:252:CYS:HB3	1:B:311:ALA:HA	1.92	0.51
1:A:15:LEU:HD11	1:A:363:LYS:HD3	1.91	0.51
1:A:274:TYR:O	1:A:315:CYS:HA	2.11	0.51
1:A:32:SER:HB3	1:A:342:THR:HG21	1.93	0.51
1:A:230:ARG:HG3	1:A:250:LYS:HG2	1.93	0.51
1:A:41:THR:HG23	1:A:43:CYS:H	1.77	0.49
1:A:150:THR:OG1	1:A:152:GLU:O	2.26	0.49
1:B:196:ARG:HG2	1:B:199:ASN:HD21	1.78	0.48
1:A:343:PRO:HA	1:A:344:PRO:HD3	1.84	0.45
1:B:404:ILE:HG21	1:A:411:LEU:HD23	1.99	0.44
1:A:132:ASP:HB2	1:A:324:LEU:HD22	2.00	0.43
1:A:8:VAL:HG11	1:A:157:VAL:HG13	2.01	0.43
1:A:52:ASP:OD2	1:A:57:ALA:HA	2.19	0.43
1:B:269:HIS:CG	1:B:287:ARG:HA	2.54	0.43
1:B:301:LEU:HD23	1:B:301:LEU:HA	1.87	0.42
1:A:132:ASP:OD2	1:A:324:LEU:HB3	2.19	0.42
1:B:132:ASP:OD1	1:B:132:ASP:N	2.44	0.42
1:B:68:MET:HB3	1:B:92:LEU:HD13	2.00	0.42
1:B:241:HIS:ND1	1:B:325:GLU:OE2	2.50	0.42
1:B:111:THR:HG21	1:B:115:LEU:HD21	2.01	0.42
1:B:196:ARG:HE	1:B:196:ARG:HB2	1.66	0.42
1:B:246:ILE:HD13	1:B:344:PRO:HD2	2.02	0.41
1:A:64:PRO:HG2	1:A:67:MET:HG2	2.02	0.41
1:A:115:LEU:HA	1:A:118:PHE:HD2	1.86	0.41
1:B:277:SER:HB2	1:B:319:GLU:OE2	2.20	0.41
1:A:36:VAL:HG13	1:A:371:ILE:HG21	2.02	0.41
1:B:391:LYS:HZ2	1:B:391:LYS:HG2	1.76	0.41
1:A:125:GLU:HB3	1:A:167:ALA:HB1	2.03	0.41
1:B:419:VAL:HG13	1:B:420:ASP:H	1.85	0.40
1:A:301:LEU:O	1:A:304:PHE:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/429 (90%)	380 (99%)	5 (1%)	0	100	100
1	B	276/429 (64%)	270 (98%)	4 (1%)	2 (1%)	18	27
All	All	661/858 (77%)	650 (98%)	9 (1%)	2 (0%)	36	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	419	VAL
1	B	420	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/399 (86%)	342 (100%)	0	100	100
1	B	336/399 (84%)	336 (100%)	0	100	100
All	All	678/798 (85%)	678 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	192	GLN
1	B	212	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	262	ASN
1	B	268	HIS
1	A	149	HIS
1	A	153	ASN
1	A	268	HIS
1	A	283	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/429 (99%)	1.28	94 (22%) 2 1	57, 79, 140, 156	0
1	B	392/429 (91%)	0.69	28 (7%) 22 18	50, 68, 95, 137	0
All	All	819/858 (95%)	1.00	122 (14%) 5 4	50, 73, 129, 156	0

All (122) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	421	ASP	5.8
1	A	65	ILE	5.6
1	A	100	TYR	5.6
1	B	129	LYS	5.3
1	A	146	LYS	4.9
1	B	419	VAL	4.7
1	B	423	GLU	4.6
1	B	422	ASP	4.4
1	B	420	ASP	4.3
1	A	130	ALA	4.3
1	A	166	GLY	4.3
1	A	60	VAL	4.3
1	A	118	PHE	4.2
1	A	86	LEU	4.2
1	A	131	ILE	4.2
1	A	98	THR	4.1
1	A	105	ASP	4.1
1	A	96	LEU	4.1
1	A	133	PHE	4.0
1	A	163	PRO	3.9
1	A	99	ARG	3.9
1	B	250	LYS	3.8
1	A	103	ILE	3.7
1	A	97	THR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	168	TRP	3.7
1	B	131	ILE	3.5
1	A	165	ARG	3.5
1	A	140	ILE	3.5
1	A	161	ILE	3.5
1	A	132	ASP	3.5
1	A	425	ASN	3.3
1	A	381	TYR	3.3
1	A	43	CYS	3.2
1	A	41	THR	3.2
1	A	77	LEU	3.2
1	A	162	GLU	3.2
1	A	250	LYS	3.2
1	A	382	ILE	3.2
1	A	115	LEU	3.2
1	A	92	LEU	3.1
1	A	104	TYR	3.1
1	A	135	GLN	3.1
1	A	79	ARG	3.1
1	A	129	LYS	3.1
1	B	135	GLN	3.1
1	A	380	LEU	3.0
1	A	421	ASP	3.0
1	A	63	ILE	3.0
1	A	127	VAL	3.0
1	A	80	PRO	3.0
1	B	134	SER	3.0
1	B	159	MET	3.0
1	A	378	LEU	3.0
1	A	95	GLU	3.0
1	A	149	HIS	2.9
1	B	128	SER	2.9
1	B	208	ASP	2.9
1	B	130	ALA	2.9
1	A	68	MET	2.9
1	A	69	LEU	2.9
1	A	107	PHE	2.9
1	A	427	TYR	2.9
1	A	90	LYS	2.8
1	A	221	LYS	2.8
1	A	144	SER	2.8
1	A	106	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	168	TRP	2.8
1	A	109	LEU	2.7
1	A	179	SER	2.7
1	A	94	ASP	2.7
1	A	420	ASP	2.7
1	A	288	HIS	2.6
1	A	62	CYS	2.6
1	B	418	ILE	2.6
1	A	152	GLU	2.6
1	B	132	ASP	2.6
1	A	76	HIS	2.6
1	A	232	VAL	2.6
1	B	209	ALA	2.5
1	A	88	GLN	2.5
1	B	395	ARG	2.5
1	A	101	HIS	2.5
1	A	58	GLY	2.5
1	A	134	SER	2.5
1	B	12	ILE	2.5
1	A	78	SER	2.5
1	A	93	THR	2.4
1	A	145	ARG	2.4
1	B	426	LEU	2.4
1	A	125	GLU	2.4
1	A	83	SER	2.4
1	A	164	GLU	2.4
1	A	287	ARG	2.4
1	B	425	ASN	2.3
1	A	102	SER	2.3
1	A	307	CYS	2.3
1	A	155	LYS	2.3
1	A	82	SER	2.3
1	A	128	SER	2.3
1	A	326	SER	2.3
1	A	239	LEU	2.3
1	A	85	GLU	2.3
1	A	72	VAL	2.3
1	A	286	HIS	2.3
1	A	47	VAL	2.3
1	A	87	ASP	2.2
1	B	239	LEU	2.2
1	B	378	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	71	LEU	2.2
1	B	15	LEU	2.2
1	B	384	GLU	2.1
1	A	171	ASN	2.1
1	A	74	SER	2.1
1	A	407	ILE	2.1
1	B	401	LYS	2.1
1	A	366	THR	2.1
1	A	110	PRO	2.1
1	A	45	GLY	2.0
1	B	325	GLU	2.0
1	A	325	GLU	2.0
1	A	342	THR	2.0
1	A	143	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.