



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:50 AM UTC

PDB ID : 1LKS / pdb_00001lks
Title : HEN EGG WHITE LYSOZYME NITRATE
Authors : Steinrauf, L.K.
Deposited on : 1997-10-09
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1656 (1.12-1.08)
Ramachandran outliers	187476	1616 (1.12-1.08)
Sidechain outliers	187428	1612 (1.12-1.08)
RSRZ outliers	180081	1625 (1.12-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NO3	A	133	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	1014	618	197	189	10	0	6	0

- Molecule 2 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	A	1	Total	N	O	0	0
			4	1	3		
2	A	1	Total	N	O	0	0
			4	1	3		
2	A	1	Total	N	O	0	0
			4	1	3		
2	A	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	197	Total	O	0	0
			197	197		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	31.97Å 65.08Å 27.25Å 94.57° 111.56° 82.98°	Depositor
Resolution (Å)	20.00 – 1.10 20.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	78.8 (20.00-1.10) 76.9 (20.00-1.75)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.75 (at 1.74Å)	Xtriage
Refinement program	SHELXL-93	Depositor
R, R_{free}	(Not available) , (Not available) 0.168 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 75.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.54	EDS
Total number of atoms	1235	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 99.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.0206e-14. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.41	8/1064 (0.8%)	1.67	20/1435 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	127	CYS	C-N	-9.96	1.19	1.33
1	A	103	ASN	CA-C	-7.68	1.43	1.52
1	A	62	TRP	CD1-NE1	7.64	1.53	1.37
1	A	105	MET	C-N	-7.08	1.24	1.33
1	A	106	ASN	C-N	-6.24	1.25	1.33
1	A	104	GLY	N-CA	6.18	1.54	1.45
1	A	112	ARG	CD-NE	-5.50	1.38	1.46
1	A	5	ARG	CB-CG	-5.31	1.36	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	112	ARG	CD-NE-CZ	19.26	151.37	124.40
1	A	128	ARG	O-C-N	13.15	140.08	122.59
1	A	103	ASN	CA-C-N	-10.31	109.55	122.75
1	A	103	ASN	C-N-CA	-10.31	109.55	122.75
1	A	103	ASN	CA-CB-CG	-9.39	103.21	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	112	ARG	NE-CZ-NH2	7.20	125.68	119.20
1	A	103	ASN	N-CA-C	7.19	121.20	109.84
1	A	103	ASN	CB-CG-ND2	-6.47	106.69	116.40
1	A	62	TRP	CB-CG-CD2	-6.37	117.89	126.80
1	A	128	ARG	CA-C-N	6.31	133.06	121.70
1	A	128	ARG	C-N-CA	6.31	133.06	121.70
1	A	99	VAL	O-C-N	-6.27	114.72	121.80
1	A	103	ASN	CB-CA-C	6.12	122.17	109.38
1	A	5	ARG	CA-CB-CG	6.09	126.28	114.10
1	A	128	ARG	CB-CA-C	-5.91	98.67	110.42
1	A	103	ASN	O-C-N	-5.58	116.38	123.24
1	A	103	ASN	OD1-CG-ND2	5.48	128.08	122.60
1	A	68	ARG	NE-CZ-NH2	5.25	123.93	119.20
1	A	84	LEU	CA-C-N	5.03	129.74	121.39
1	A	84	LEU	C-N-CA	5.03	129.74	121.39

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	103	ASN	CA

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	103	ASN	Sidechain
1	A	127	CYS	Peptide
1	A	61[A]	ARG	Sidechain
1	A	61[B]	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1014	0	964	22	0
2	A	24	0	0	3	0
3	A	197	0	0	4	0
All	All	1235	0	964	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ARG:HB3	1:A:129:LEU:HG	1.16	1.13
1:A:128:ARG:HG3	1:A:129:LEU:H	1.10	1.09
1:A:128:ARG:HG3	1:A:129:LEU:N	1.73	1.03
1:A:106:ASN:ND2	2:A:133:NO3:O2	1.95	1.00
1:A:128:ARG:CG	1:A:129:LEU:H	1.40	0.97
1:A:128:ARG:HB3	1:A:129:LEU:CG	1.99	0.93
1:A:106:ASN:ND2	2:A:133:NO3:O1	2.03	0.89
1:A:106:ASN:OD1	3:A:228:HOH:O	1.92	0.88
1:A:128:ARG:CG	1:A:129:LEU:N	2.22	0.83
1:A:106:ASN:ND2	2:A:133:NO3:N	2.29	0.80
1:A:102[B]:GLY:O	1:A:103:ASN:HB3	1.98	0.63
1:A:128:ARG:HG3	1:A:129:LEU:CA	2.31	0.59
1:A:128:ARG:O	1:A:129:LEU:O	2.26	0.54
1:A:128:ARG:HG2	1:A:129:LEU:HD23	1.91	0.52
1:A:104:GLY:N	3:A:213:HOH:O	2.44	0.47
1:A:99:VAL:HA	1:A:103:ASN:O	2.15	0.46
1:A:128:ARG:C	1:A:129:LEU:O	2.60	0.45
1:A:68:ARG:NH2	3:A:173:HOH:O	2.45	0.45
1:A:68:ARG:HG3	1:A:68:ARG:HH11	1.82	0.44
1:A:128:ARG:CG	1:A:129:LEU:HD23	2.48	0.43
1:A:128:ARG:HG3	1:A:129:LEU:HA	1.99	0.43
1:A:106:ASN:HB2	3:A:213:HOH:O	2.19	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/129 (102%)	130 (98%)	2 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/105 (105%)	108 (98%)	2 (2%)	51	15

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	ASN
1	A	129	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NO3	A	132	-	1,3,3	0.06	0	0,3,3	-	-
2	NO3	A	134	-	1,3,3	0.72	0	0,3,3	-	-
2	NO3	A	130	-	1,3,3	1.12	0	0,3,3	-	-
2	NO3	A	131	-	1,3,3	0.69	0	0,3,3	-	-
2	NO3	A	133	-	1,3,3	0.26	0	0,3,3	-	-
2	NO3	A	135	-	1,3,3	0.19	0	0,3,3	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	133	NO3	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	127:CYS	C	128:ARG	N	1.19

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/129 (100%)	3.15	95 (73%) 0 0	8, 13, 28, 56	6 (4%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	129	LEU	10.9
1	A	123	TRP	8.2
1	A	102[A]	GLY	8.1
1	A	99	VAL	7.2
1	A	103	ASN	7.1
1	A	43[A]	THR	7.0
1	A	128	ARG	6.3
1	A	106	ASN	6.2
1	A	109	VAL	5.9
1	A	62	TRP	5.8
1	A	87	ASP	5.7
1	A	100	SER	5.3
1	A	78	ILE	5.2
1	A	63	TRP	5.0
1	A	108	TRP	5.0
1	A	92	VAL	5.0
1	A	34	PHE	4.9
1	A	120	VAL	4.7
1	A	126	GLY	4.7
1	A	118	THR	4.5
1	A	124	ILE	4.5
1	A	84	LEU	4.5
1	A	56	LEU	4.5
1	A	122	ALA	4.5
1	A	47	THR	4.3
1	A	55	ILE	4.3
1	A	112	ARG	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	28	TRP	4.1
1	A	25	LEU	4.1
1	A	113	ASN	4.1
1	A	127	CYS	4.0
1	A	75	LEU	3.9
1	A	6	CYS	3.8
1	A	101	ASP	3.8
1	A	53	TYR	3.8
1	A	64	CYS	3.7
1	A	111	TRP	3.7
1	A	88	ILE	3.6
1	A	121	GLN	3.6
1	A	79	PRO	3.6
1	A	110	ALA	3.5
1	A	58	ILE	3.5
1	A	83	LEU	3.5
1	A	17	LEU	3.4
1	A	68	ARG	3.4
1	A	107	ALA	3.4
1	A	13	LYS	3.4
1	A	104	GLY	3.4
1	A	3	PHE	3.3
1	A	98	ILE	3.3
1	A	5	ARG	3.3
1	A	21	ARG	3.3
1	A	41	GLN	3.3
1	A	20	TYR	3.3
1	A	38	PHE	3.2
1	A	97	LYS	3.2
1	A	23	TYR	3.2
1	A	11	ALA	3.1
1	A	61[A]	ARG	3.1
1	A	86[A]	SER	3.1
1	A	29	VAL	3.0
1	A	2	VAL	3.0
1	A	125	ARG	2.9
1	A	70	PRO	2.9
1	A	18	ASP	2.9
1	A	19	ASN	2.8
1	A	14	ARG	2.7
1	A	82	ALA	2.7
1	A	46	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	114	ARG	2.6
1	A	89	THR	2.6
1	A	44	ASN	2.6
1	A	51	THR	2.6
1	A	15	HIS	2.5
1	A	32	ALA	2.5
1	A	4	GLY	2.4
1	A	67	GLY	2.4
1	A	37	ASN	2.4
1	A	76	CYS	2.4
1	A	45	ARG	2.3
1	A	81	SER	2.3
1	A	94	CYS	2.2
1	A	50	SER	2.2
1	A	93	ASN	2.2
1	A	9	ALA	2.2
1	A	95	ALA	2.2
1	A	72	SER	2.1
1	A	119	ASP	2.1
1	A	26	GLY	2.1
1	A	59	ASN	2.1
1	A	115	CYS	2.1
1	A	24	SER	2.0
1	A	30	CYS	2.0
1	A	69	THR	2.0
1	A	105	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NO3	A	133	4/4	0.42	0.22	38,42,44,44	4
2	NO3	A	135	4/4	0.50	0.24	42,42,42,47	4
2	NO3	A	130	4/4	0.60	0.25	17,20,20,21	4
2	NO3	A	134	4/4	0.71	0.20	25,25,28,31	4
2	NO3	A	131	4/4	0.76	0.15	20,22,26,27	0
2	NO3	A	132	4/4	0.81	0.12	33,33,34,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.