



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:44 PM UTC

PDB ID : 7LKT / pdb_00007lkt
Title : 1.50 Å resolution structure of SARS-CoV-2 3CL protease in complex with inhibitor 2k
Authors : Kashipathy, M.M.; Lovell, S.; Battaile, K.P.; Chamandi, S.D.; Rathnayake, A.D.; Nguyen, H.N.; Baird, M.A.; Kim, Y.; Shadipeni, N.; Chang, K.O.; Groutas, W.C.
Deposited on : 2021-02-02
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	12% 89% 7% .
1	B	309	10% 92% 6% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	6	0
			2272	1442	381	425	24			
1	B	300	Total	C	N	O	S	0	2	0
			2286	1455	379	429	23			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	SER	-	expression tag	UNP P0DTD1
A	-2	ASN	-	expression tag	UNP P0DTD1
A	-1	ILE	-	expression tag	UNP P0DTD1
A	0	GLY	-	expression tag	UNP P0DTD1
B	-3	SER	-	expression tag	UNP P0DTD1
B	-2	ASN	-	expression tag	UNP P0DTD1
B	-1	ILE	-	expression tag	UNP P0DTD1
B	0	GLY	-	expression tag	UNP P0DTD1

- Molecule 2 is (1S,2S)-2-((S)-2-(((adamantan-1-ylmethoxy)carbonyl)amino)-4-methylpentanamido)-1-hydroxy-3-((S)-2-oxopyrrolidin-3-yl)propane-1-sulfonic acid (CCD ID: Y7M) (formula: C₂₅H₄₁N₃O₈S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 33	C 25	N 3	O 5	0	1
2	B	1	Total 33	C 25	N 3	O 5	0	1

- Molecule 3 is (1R,2S)-2-((S)-2-(((adamantan-1-ylmethoxy)carbonyl)amino)-4-methylpentanamido)-1-hydroxy-3-((S)-2-oxopyrrolidin-3-yl)propane-1-sulfonic acid (CCD ID: Y4V) (formula: C₂₅H₄₁N₃O₈S).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	1
			33	25	3	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	1
			33	25	3	5		

- Molecule 4 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			13	8	5		
5	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	202	Total	O	0	0
			202	202		
6	B	218	Total	O	0	0
			218	218		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3C-like proteinase



• Molecule 1: 3C-like proteinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.10Å 98.86Å 59.42Å 90.00° 107.91° 90.00°	Depositor
Resolution (Å)	37.21 – 1.50 37.21 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.21-1.50) 99.9 (37.21-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.50Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.178 , 0.205 0.186 , 0.210	Depositor DCC
R_{free} test set	4828 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5149	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Y7M, PG4, Y4V, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2338	0.60	0/3183
1	B	0.43	0/2342	0.61	0/3188
All	All	0.42	0/4680	0.61	0/6371

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2272	0	2211	12	0
1	B	2286	0	2209	14	0
2	A	33	0	0	0	0
2	B	33	0	0	0	0
3	A	33	0	0	0	0
3	B	33	0	0	0	0
4	B	13	0	5	0	0
5	B	26	0	36	4	0
6	A	202	0	0	0	0
6	B	218	0	0	2	0
All	All	5149	0	4461	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:279:ARG:NH2	6:B:501:HOH:O	2.20	0.74
1:A:92:ASP:OD1	1:A:93:THR:HG23	2.02	0.60
1:A:188:ARG:HG2	1:A:190[A]:THR:HG23	1.89	0.55
1:B:3:PHE:CZ	1:B:210:ALA:HB1	2.43	0.54
1:A:109:GLY:HA2	1:A:200:ILE:HD13	1.90	0.54
1:A:86[A]:VAL:HG13	1:A:179:GLY:HA2	1.89	0.53
1:B:86[A]:VAL:HG13	1:B:179:GLY:HA2	1.90	0.53
1:A:126:TYR:CD1	1:B:4:ARG:HD2	2.44	0.52
1:B:269:LYS:NZ	6:B:507:HOH:O	2.40	0.52
1:A:165[B]:MET:HE1	1:A:192:GLN:NE2	2.27	0.49
1:A:22[B]:CYS:SG	1:A:66:PHE:CD1	3.08	0.47
1:B:90:LYS:CD	5:B:405:PG4:H11	2.46	0.46
1:A:233:VAL:HG11	1:A:269:LYS:HG3	1.98	0.45
1:A:297:VAL:O	1:A:300:CYS:N	2.39	0.45
1:A:31:TRP:CE2	1:A:95:ASN:HB2	2.51	0.44
1:B:113:SER:O	1:B:149:GLY:HA2	2.17	0.44
1:B:90:LYS:CD	5:B:405:PG4:O1	2.65	0.44
1:A:113:SER:O	1:A:149:GLY:HA2	2.18	0.44
1:A:257:THR:HB	1:A:259:ILE:HD12	2.00	0.44
1:B:75:LEU:HD21	1:B:93:THR:HB	2.00	0.43
1:B:90:LYS:HD3	5:B:405:PG4:C1	2.48	0.43
1:B:31:TRP:CE2	1:B:75:LEU:HD11	2.53	0.43
1:B:45:THR:O	1:B:49:MET:HG3	2.19	0.42
1:B:90:LYS:HD2	5:B:405:PG4:H11	2.02	0.42
1:B:126:TYR:HE1	1:B:128:CYS:SG	2.44	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/309 (97%)	294 (98%)	7 (2%)	0	100	100
1	B	300/309 (97%)	294 (98%)	6 (2%)	0	100	100
All	All	601/618 (97%)	588 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/265 (94%)	248 (100%)	0	100	100
1	B	247/265 (93%)	247 (100%)	0	100	100
All	All	495/530 (93%)	495 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	274	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FLC	B	401	-	12,12,12	1.55	4 (33%)	17,17,17	1.98	4 (23%)
3	Y4V	A	402[B]	1	36,36,40	1.98	8 (22%)	48,52,60	2.02	10 (20%)
5	PG4	B	405	-	12,12,12	0.17	0	11,11,11	0.62	0
2	Y7M	A	401[A]	1	36,36,40	1.65	6 (16%)	48,52,60	1.93	9 (18%)
5	PG4	B	404	-	12,12,12	0.14	0	11,11,11	0.46	0
2	Y7M	B	402[A]	1	36,36,40	1.94	9 (25%)	48,52,60	1.91	7 (14%)
3	Y4V	B	403[B]	1	36,36,40	2.12	12 (33%)	48,52,60	2.05	12 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FLC	B	401	-	-	5/16/16/16	-
3	Y4V	A	402[B]	1	-	5/28/65/73	0/5/4/4
5	PG4	B	405	-	-	3/10/10/10	-
2	Y7M	A	401[A]	1	-	0/28/65/73	0/5/4/4
5	PG4	B	404	-	-	0/10/10/10	-
2	Y7M	B	402[A]	1	-	5/28/65/73	0/5/4/4
3	Y4V	B	403[B]	1	-	2/28/65/73	0/5/4/4

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403[B]	Y4V	C14-N15	6.78	1.40	1.33
2	B	402[A]	Y7M	C14-N15	6.72	1.40	1.33
3	A	402[B]	Y4V	C14-N15	6.46	1.40	1.33
2	A	401[A]	Y7M	C14-N15	6.03	1.40	1.33
3	B	403[B]	Y4V	C19-C11	3.95	1.59	1.52
2	B	402[A]	Y7M	C32-C24	3.82	1.58	1.53
3	B	403[B]	Y4V	C32-C24	3.71	1.58	1.53
3	A	402[B]	Y4V	C19-C11	3.61	1.59	1.52
2	B	402[A]	Y7M	C19-C11	3.46	1.58	1.52
2	B	402[A]	Y7M	C17-C16	-3.45	1.48	1.53
2	A	401[A]	Y7M	C17-C16	-3.43	1.48	1.53
3	B	403[B]	Y4V	C23-C24	3.37	1.61	1.53
3	A	402[B]	Y4V	C23-C24	3.37	1.61	1.53
3	A	402[B]	Y4V	C32-C24	3.33	1.58	1.53
3	B	403[B]	Y4V	C17-C16	-3.09	1.48	1.53
3	A	402[B]	Y4V	C17-C16	-3.08	1.48	1.53
2	A	401[A]	Y7M	C32-C24	3.02	1.57	1.53
3	B	403[B]	Y4V	C02-N03	2.95	1.42	1.34
3	B	403[B]	Y4V	C09-N10	2.92	1.40	1.34
3	A	402[B]	Y4V	C09-N10	2.89	1.40	1.34
4	B	401	FLC	OB1-CBC	2.85	1.31	1.22
4	B	401	FLC	OA1-CAC	2.83	1.31	1.22
4	B	401	FLC	OA2-CAC	-2.66	1.22	1.30
2	B	402[A]	Y7M	C23-C24	2.57	1.59	1.53
3	B	403[B]	Y4V	O22-C02	2.52	1.40	1.35
2	B	402[A]	Y7M	C09-N10	2.28	1.38	1.34
2	B	402[A]	Y7M	C02-N03	2.28	1.40	1.34
2	A	401[A]	Y7M	C23-C24	2.28	1.58	1.53
4	B	401	FLC	OB2-CBC	-2.20	1.22	1.30
2	A	401[A]	Y7M	C12-C11	2.16	1.58	1.53
2	A	401[A]	Y7M	C19-C11	2.13	1.56	1.52
3	B	403[B]	Y4V	O21-C09	2.11	1.27	1.23
2	B	402[A]	Y7M	C12-C11	2.10	1.58	1.53
3	A	402[B]	Y4V	C02-N03	2.09	1.39	1.34
3	B	403[B]	Y4V	C04-N03	2.07	1.50	1.45
3	B	403[B]	Y4V	C12-C11	2.06	1.58	1.53
3	A	402[B]	Y4V	O21-C09	2.04	1.27	1.23
2	B	402[A]	Y7M	O21-C09	2.03	1.27	1.23
3	B	403[B]	Y4V	C05-C04	2.01	1.59	1.53

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[A]	Y7M	C29-C24-C25	7.16	114.00	108.47
2	B	402[A]	Y7M	C29-C24-C25	6.94	113.83	108.47
3	B	403[B]	Y4V	C29-C24-C25	5.77	112.93	108.47
3	B	403[B]	Y4V	O22-C02-N03	5.66	122.57	110.45
3	A	402[B]	Y4V	C29-C24-C25	5.57	112.77	108.47
4	B	401	FLC	OB2-CBC-CB	5.14	123.00	113.14
2	A	401[A]	Y7M	C32-C24-C25	-5.11	104.53	108.47
3	A	402[B]	Y4V	O22-C02-N03	5.10	121.36	110.45
2	B	402[A]	Y7M	O22-C02-N03	5.08	121.31	110.45
3	A	402[B]	Y4V	C32-C24-C25	-4.93	104.68	108.47
4	B	401	FLC	OB1-CBC-CB	-4.77	112.84	122.09
2	A	401[A]	Y7M	O18-C14-N15	-4.71	119.17	125.76
3	B	403[B]	Y4V	O18-C14-N15	-4.55	119.41	125.76
3	A	402[B]	Y4V	O18-C14-N15	-4.34	119.70	125.76
3	B	403[B]	Y4V	C32-C24-C25	-4.29	105.17	108.47
2	B	402[A]	Y7M	C32-C24-C25	-4.18	105.25	108.47
2	B	402[A]	Y7M	O18-C14-N15	-4.15	119.96	125.76
3	B	403[B]	Y4V	C16-N15-C14	-4.03	105.23	113.79
3	B	403[B]	Y4V	O22-C02-O01	-3.98	116.62	124.26
3	A	402[B]	Y4V	C16-N15-C14	-3.88	105.55	113.79
2	B	402[A]	Y7M	C16-N15-C14	-3.88	105.56	113.79
3	A	402[B]	Y4V	O22-C02-O01	-3.83	116.92	124.26
2	A	401[A]	Y7M	C16-N15-C14	-3.71	105.91	113.79
2	A	401[A]	Y7M	O22-C02-N03	3.54	118.01	110.45
2	B	402[A]	Y7M	O22-C02-O01	-3.47	117.60	124.26
3	A	402[B]	Y4V	C24-C25-C26	3.33	112.75	110.43
3	B	403[B]	Y4V	C04-C09-N10	3.19	123.43	116.63
4	B	401	FLC	OA1-CAC-CA	-3.10	114.16	122.95
3	B	403[B]	Y4V	O20-C19-C11	-3.03	104.11	111.73
2	A	401[A]	Y7M	O22-C02-O01	-3.00	118.50	124.26
3	A	402[B]	Y4V	C04-C09-N10	2.94	122.91	116.63
3	A	402[B]	Y4V	O20-C19-C11	-2.74	104.84	111.73
3	B	403[B]	Y4V	C24-C25-C26	2.67	112.29	110.43
3	B	403[B]	Y4V	O21-C09-N10	-2.66	118.20	122.96
4	B	401	FLC	OA2-CAC-CA	2.60	122.58	114.35
3	B	403[B]	Y4V	O01-C02-N03	-2.47	120.81	124.86
3	A	402[B]	Y4V	O21-C09-N10	-2.43	118.60	122.96
2	A	401[A]	Y7M	O21-C09-N10	-2.32	118.80	122.96
2	B	402[A]	Y7M	O01-C02-N03	-2.30	121.09	124.86
3	B	403[B]	Y4V	C32-C24-C29	-2.28	106.71	108.47
2	A	401[A]	Y7M	C24-C25-C26	2.15	111.93	110.43
2	A	401[A]	Y7M	C06-C05-C04	-2.02	109.97	115.40

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	402[A]	Y7M	O22-C23-C24-C25
2	B	402[A]	Y7M	O22-C23-C24-C29
2	B	402[A]	Y7M	O22-C23-C24-C32
4	B	401	FLC	CA-CB-CG-CGC
4	B	401	FLC	CBC-CB-CG-CGC
2	B	402[A]	Y7M	N03-C02-O22-C23
4	B	401	FLC	OHB-CB-CG-CGC
2	B	402[A]	Y7M	O01-C02-O22-C23
5	B	405	PG4	O1-C1-C2-O2
5	B	405	PG4	C3-C4-O3-C5
4	B	401	FLC	CAC-CA-CB-OHB
4	B	401	FLC	CAC-CA-CB-CG
3	A	402[B]	Y4V	O22-C23-C24-C25
3	A	402[B]	Y4V	N03-C02-O22-C23
3	B	403[B]	Y4V	N03-C02-O22-C23
3	A	402[B]	Y4V	O22-C02-N03-C04
3	B	403[B]	Y4V	O01-C02-O22-C23
3	A	402[B]	Y4V	O01-C02-O22-C23
3	A	402[B]	Y4V	O01-C02-N03-C04
5	B	405	PG4	O2-C3-C4-O3


There are no ring outliers.

1 monomer is involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	405	PG4	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand Y4V A 402 (B)



Ligand Y7M A 401 (A)





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/309 (96%)	0.73	37 (12%) 8 8	11, 24, 46, 62	6 (2%)
1	B	300/309 (97%)	0.52	30 (10%) 12 13	11, 21, 44, 54	2 (0%)
All	All	597/618 (96%)	0.62	67 (11%) 10 10	11, 22, 45, 62	8 (1%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	223	PHE	4.7
1	B	3	PHE	4.5
1	B	2	GLY	4.4
1	B	300	CYS	4.1
1	A	300	CYS	3.8
1	B	298	ARG	3.8
1	B	301	SER	3.6
1	A	224	THR	3.5
1	B	277	ASN	3.4
1	B	297	VAL	3.4
1	A	218	TRP	3.3
1	A	213	ILE	3.3
1	B	191	ALA	3.2
1	A	216	ASP	3.1
1	A	212	VAL	3.1
1	A	255	ALA	3.1
1	A	93	THR	3.1
1	B	154	TYR	3.0
1	B	155	ASP	3.0
1	A	222	ARG	3.0
1	A	294	PHE	2.9
1	B	46	SER	2.9
1	B	223	PHE	2.9
1	A	4	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	280	THR	2.9
1	A	282	LEU	2.8
1	A	72	ASN	2.8
1	B	196	THR	2.8
1	A	298	ARG	2.8
1	A	283	GLY	2.7
1	A	207	TRP	2.6
1	B	222	ARG	2.6
1	A	219	PHE	2.6
1	A	154	TYR	2.6
1	A	46	SER	2.6
1	A	259	ILE	2.5
1	A	278	GLY	2.5
1	A	210	ALA	2.5
1	B	137	LYS	2.4
1	B	190	THR	2.4
1	B	193	ALA	2.4
1	A	249	ILE	2.4
1	B	23	GLY	2.4
1	A	297	VAL	2.4
1	A	155	ASP	2.3
1	B	299	GLN	2.3
1	B	49	MET	2.3
1	B	4	ARG	2.3
1	A	253	LEU	2.3
1	A	281	ILE	2.3
1	B	48	ASP	2.2
1	B	168	PRO	2.2
1	A	275	GLY	2.2
1	A	291	PHE	2.2
1	A	287	LEU	2.2
1	B	50	LEU	2.2
1	A	220	LEU	2.1
1	A	260	ALA	2.1
1	A	209	TYR	2.1
1	B	283	GLY	2.1
1	B	47	GLU	2.1
1	B	195	GLY	2.1
1	B	278	GLY	2.1
1	A	45	THR	2.1
1	A	299	GLN	2.1
1	B	280	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	128	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FLC	B	401	13/13	0.69	0.17	33,53,61,69	0
5	PG4	B	405	13/13	0.83	0.15	34,37,46,54	0
3	Y4V	B	403[B]	33/37	0.90	0.12	19,25,37,39	33
2	Y7M	B	402[A]	33/37	0.92	0.11	19,26,37,39	33
3	Y4V	A	402[B]	33/37	0.95	0.07	13,17,24,26	33
5	PG4	B	404	13/13	0.97	0.06	17,19,29,32	0
2	Y7M	A	401[A]	33/37	0.97	0.07	13,17,24,26	33

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Y4V B 403 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Y7M B 402 (A):


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Y4V A 402 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.