



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 01:52 AM UTC

PDB ID : 1LL0 / pdb_00001ll0
Title : Crystal Structure of Rabbit Muscle Glycogenin
Authors : Gibbons, B.J.; Roach, P.J.; Hurley, T.D.
Deposited on : 2002-04-26
Resolution : 3.43 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1736 (3.50-3.38)
Clashscore	190562	1808 (3.50-3.38)
Ramachandran outliers	187476	1776 (3.50-3.38)
Sidechain outliers	187428	1777 (3.50-3.38)
RSRZ outliers	180081	1736 (3.50-3.38)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	6% 39% 33% 6% 22%
1	B	339	7% 39% 32% 7% 21%
1	C	339	5% 40% 32% 6% 23%
1	D	339	4% 41% 29% 7% 23%
1	E	339	4% 40% 31% 6% 23%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	339	7%39%32%6%23%
1	G	339	6%38%33%6%23%
1	H	339	8%39%30%7%23%
1	I	339	6%38%33%6%22%
1	J	339	13%40%29%5%26%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 20736 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCOGENIN-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2092	1346	342	396	8			
1	B	267	Total	C	N	O	S	0	0	0
			2115	1360	350	397	8			
1	C	262	Total	C	N	O	S	0	0	0
			2077	1337	340	392	8			
1	D	261	Total	C	N	O	S	0	0	0
			2071	1333	339	392	7			
1	E	262	Total	C	N	O	S	0	0	0
			2081	1339	342	393	7			
1	F	261	Total	C	N	O	S	0	0	0
			2071	1333	339	392	7			
1	G	262	Total	C	N	O	S	0	0	0
			2079	1338	340	393	8			
1	H	260	Total	C	N	O	S	0	0	0
			2064	1327	338	391	8			
1	I	264	Total	C	N	O	S	0	0	0
			2095	1346	347	395	7			
1	J	251	Total	C	N	O	S	0	0	0
			1991	1281	327	377	6			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	VAL	-	expression tag	UNP P13280
A	-6	PRO	-	expression tag	UNP P13280
A	-5	ARG	-	expression tag	UNP P13280
A	-4	GLY	-	expression tag	UNP P13280
A	-3	SER	-	expression tag	UNP P13280
A	-2	HIS	-	expression tag	UNP P13280
B	-7	VAL	-	expression tag	UNP P13280
B	-6	PRO	-	expression tag	UNP P13280
B	-5	ARG	-	expression tag	UNP P13280

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP P13280
B	-3	SER	-	expression tag	UNP P13280
B	-2	HIS	-	expression tag	UNP P13280
C	-7	VAL	-	expression tag	UNP P13280
C	-6	PRO	-	expression tag	UNP P13280
C	-5	ARG	-	expression tag	UNP P13280
C	-4	GLY	-	expression tag	UNP P13280
C	-3	SER	-	expression tag	UNP P13280
C	-2	HIS	-	expression tag	UNP P13280
D	-7	VAL	-	expression tag	UNP P13280
D	-6	PRO	-	expression tag	UNP P13280
D	-5	ARG	-	expression tag	UNP P13280
D	-4	GLY	-	expression tag	UNP P13280
D	-3	SER	-	expression tag	UNP P13280
D	-2	HIS	-	expression tag	UNP P13280
E	-7	VAL	-	expression tag	UNP P13280
E	-6	PRO	-	expression tag	UNP P13280
E	-5	ARG	-	expression tag	UNP P13280
E	-4	GLY	-	expression tag	UNP P13280
E	-3	SER	-	expression tag	UNP P13280
E	-2	HIS	-	expression tag	UNP P13280
F	-7	VAL	-	expression tag	UNP P13280
F	-6	PRO	-	expression tag	UNP P13280
F	-5	ARG	-	expression tag	UNP P13280
F	-4	GLY	-	expression tag	UNP P13280
F	-3	SER	-	expression tag	UNP P13280
F	-2	HIS	-	expression tag	UNP P13280
G	-7	VAL	-	expression tag	UNP P13280
G	-6	PRO	-	expression tag	UNP P13280
G	-5	ARG	-	expression tag	UNP P13280
G	-4	GLY	-	expression tag	UNP P13280
G	-3	SER	-	expression tag	UNP P13280
G	-2	HIS	-	expression tag	UNP P13280
H	-7	VAL	-	expression tag	UNP P13280
H	-6	PRO	-	expression tag	UNP P13280
H	-5	ARG	-	expression tag	UNP P13280
H	-4	GLY	-	expression tag	UNP P13280
H	-3	SER	-	expression tag	UNP P13280
H	-2	HIS	-	expression tag	UNP P13280
I	-7	VAL	-	expression tag	UNP P13280
I	-6	PRO	-	expression tag	UNP P13280
I	-5	ARG	-	expression tag	UNP P13280

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	-4	GLY	-	expression tag	UNP P13280
I	-3	SER	-	expression tag	UNP P13280
I	-2	HIS	-	expression tag	UNP P13280
J	-7	VAL	-	expression tag	UNP P13280
J	-6	PRO	-	expression tag	UNP P13280
J	-5	ARG	-	expression tag	UNP P13280
J	-4	GLY	-	expression tag	UNP P13280
J	-3	SER	-	expression tag	UNP P13280
J	-2	HIS	-	expression tag	UNP P13280


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GLYCOGENIN-1




• Molecule 1: GLYCOGENIN-1




VAL SER SER SER GLU GLU ARG LYS
GLU ARG TRP GLU GLY GLN ASP
ALA ASP MET GLY ALA


• Molecule 1: GLYCOGENIN-1




• Molecule 1: GLYCOGENIN-1



• Molecule 1: GLYCOGENIN-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	139.46Å 139.46Å 416.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 3.43 25.00 – 3.43	Depositor EDS
% Data completeness (in resolution range)	95.8 (25.00-3.43) 95.4 (25.00-3.43)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.44 (at 3.47Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.252 , 0.287 0.245 , 0.246	Depositor DCC
R_{free} test set	2837 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	90.2	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 90.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	20736	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.70	1/2146 (0.0%)	1.11	20/2927 (0.7%)
1	B	0.88	6/2170 (0.3%)	1.18	28/2958 (0.9%)
1	C	0.65	1/2130 (0.0%)	1.04	8/2904 (0.3%)
1	D	0.65	1/2123 (0.0%)	1.14	18/2893 (0.6%)
1	E	0.63	0/2135	1.06	11/2911 (0.4%)
1	F	0.59	1/2124 (0.0%)	1.08	14/2896 (0.5%)
1	G	0.61	0/2132	1.08	18/2906 (0.6%)
1	H	0.86	6/2117 (0.3%)	1.15	20/2885 (0.7%)
1	I	0.68	2/2149 (0.1%)	1.15	26/2928 (0.9%)
1	J	0.52	0/2043	1.04	12/2787 (0.4%)
All	All	0.69	18/21269 (0.1%)	1.10	175/28995 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
1	G	0	1
1	H	0	1
1	I	0	1
All	All	0	7

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	241	HIS	C-N	19.02	1.50	1.33
1	B	188	LEU	N-CA	17.02	1.68	1.46
1	H	241	HIS	CA-C	14.27	1.70	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	188	LEU	N-CA	12.65	1.62	1.46
1	B	-4	GLY	C-N	9.97	1.47	1.33

The worst 5 of 175 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	187	ASN	O-C-N	-18.36	99.45	122.21
1	D	241	HIS	CA-C-N	-15.70	101.73	118.85
1	D	241	HIS	C-N-CA	-15.70	101.73	118.85
1	A	208	LYS	N-CA-CB	14.25	131.07	110.40
1	D	187	ASN	O-C-N	-13.74	104.95	121.99

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	187	ASN	Mainchain
1	D	187	ASN	Mainchain
1	E	187	ASN	Mainchain
1	F	187	ASN	Mainchain
1	G	187	ASN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2092	0	2048	159	0
1	B	2115	0	2074	166	0
1	C	2077	0	2037	153	0
1	D	2071	0	2026	115	0
1	E	2081	0	2034	137	0
1	F	2071	0	2027	128	0
1	G	2079	0	2036	145	0
1	H	2064	0	2016	126	0
1	I	2095	0	2048	145	0
1	J	1991	0	1946	103	0
All	All	20736	0	20292	1337	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 1337 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:188:LEU:N	1:B:188:LEU:CA	1.68	1.52
1:A:193:ILE:CB	1:A:240:THR:HG21	1.55	1.37
1:B:200:PHE:CE2	1:B:240:THR:HB	1.58	1.36
1:A:193:ILE:HB	1:A:240:THR:CG2	1.57	1.32
1:C:200:PHE:CE2	1:C:240:THR:HB	1.66	1.30

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/339 (77%)	223 (86%)	31 (12%)	6 (2%)	5	27
1	B	263/339 (78%)	223 (85%)	31 (12%)	9 (3%)	3	21
1	C	258/339 (76%)	220 (85%)	33 (13%)	5 (2%)	6	30
1	D	256/339 (76%)	219 (86%)	31 (12%)	6 (2%)	5	27
1	E	258/339 (76%)	224 (87%)	28 (11%)	6 (2%)	5	27
1	F	257/339 (76%)	221 (86%)	29 (11%)	7 (3%)	4	24
1	G	258/339 (76%)	225 (87%)	28 (11%)	5 (2%)	6	30
1	H	256/339 (76%)	221 (86%)	26 (10%)	9 (4%)	3	20
1	I	260/339 (77%)	226 (87%)	30 (12%)	4 (2%)	8	34
1	J	247/339 (73%)	213 (86%)	30 (12%)	4 (2%)	7	33
All	All	2573/3390 (76%)	2215 (86%)	297 (12%)	61 (2%)	4	26

5 of 61 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	240	THR
1	H	241	HIS
1	A	128	PRO
1	A	190	SER
1	A	215	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/299 (78%)	212 (91%)	22 (9%)	8	30
1	B	236/299 (79%)	215 (91%)	21 (9%)	9	32
1	C	232/299 (78%)	212 (91%)	20 (9%)	10	33
1	D	231/299 (77%)	209 (90%)	22 (10%)	8	30
1	E	232/299 (78%)	210 (90%)	22 (10%)	8	30
1	F	231/299 (77%)	210 (91%)	21 (9%)	9	31
1	G	232/299 (78%)	210 (90%)	22 (10%)	8	30
1	H	230/299 (77%)	206 (90%)	24 (10%)	7	26
1	I	233/299 (78%)	212 (91%)	21 (9%)	9	32
1	J	222/299 (74%)	202 (91%)	20 (9%)	9	32
All	All	2313/2990 (77%)	2098 (91%)	215 (9%)	8	31

5 of 215 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	138	TYR
1	G	194	TYR
1	J	89	TRP
1	F	155	GLN
1	G	21	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 107 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	93	GLN
1	G	147	GLN
1	J	93	GLN
1	F	139	GLN
1	F	181	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	238:THR	C	241:HIS	N	5.99

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	264/339 (77%)	0.44	20 (7%)	20 15	12, 58, 138, 194	0
1	B	267/339 (78%)	0.51	23 (8%)	16 13	16, 49, 129, 200	0
1	C	262/339 (77%)	0.49	18 (6%)	23 17	18, 49, 119, 189	0
1	D	261/339 (76%)	0.58	14 (5%)	31 21	25, 61, 143, 200	0
1	E	262/339 (77%)	0.35	13 (4%)	34 22	23, 56, 122, 200	0
1	F	261/339 (76%)	0.70	23 (8%)	15 12	27, 73, 151, 189	0
1	G	262/339 (77%)	0.50	19 (7%)	21 16	21, 56, 130, 199	0
1	H	260/339 (76%)	0.62	27 (10%)	11 10	30, 75, 154, 200	0
1	I	264/339 (77%)	0.61	21 (7%)	18 14	19, 59, 134, 187	0
1	J	251/339 (74%)	1.14	45 (17%)	3 4	48, 99, 160, 200	0
All	All	2614/3390 (77%)	0.59	223 (8%)	16 13	12, 63, 145, 200	0

The worst 5 of 223 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	207	ALA	6.7
1	G	240	THR	6.0
1	F	241	HIS	5.6
1	I	-1	MET	5.6
1	B	-7	VAL	5.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.