



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:52 PM UTC

PDB ID : 6LL4 / pdb_00006ll4
Title : Oxygen-exposed carbazole-soaked reduced terminal oxygenase of carbazole
1,9a-dioxygenase
Authors : Wang, Y.X.; Suzuki-Minakuchi, C.; Nojiri, H.
Deposited on : 2019-12-21
Resolution : 2.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	
1	B	392	
1	C	392	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	B	410	-	-	X	-
5	EDO	B	414	-	-	X	-

2 Entry composition i

There are 11 unique types of molecules in this entry. The entry contains 10038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Terminal oxygenase component of carbazole.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	383	Total 3207	C 2049	N 545	O 600	S 13	0	18	0
1	B	383	Total 3081	C 1970	N 523	O 575	S 13	0	0	0
1	C	383	Total 3092	C 1978	N 524	O 577	S 13	0	2	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	385	LEU	-	expression tag	UNP Q84II6
A	386	GLU	-	expression tag	UNP Q84II6
A	387	HIS	-	expression tag	UNP Q84II6
A	388	HIS	-	expression tag	UNP Q84II6
A	389	HIS	-	expression tag	UNP Q84II6
A	390	HIS	-	expression tag	UNP Q84II6
A	391	HIS	-	expression tag	UNP Q84II6
A	392	HIS	-	expression tag	UNP Q84II6
B	385	LEU	-	expression tag	UNP Q84II6
B	386	GLU	-	expression tag	UNP Q84II6
B	387	HIS	-	expression tag	UNP Q84II6
B	388	HIS	-	expression tag	UNP Q84II6
B	389	HIS	-	expression tag	UNP Q84II6
B	390	HIS	-	expression tag	UNP Q84II6
B	391	HIS	-	expression tag	UNP Q84II6
B	392	HIS	-	expression tag	UNP Q84II6
C	385	LEU	-	expression tag	UNP Q84II6
C	386	GLU	-	expression tag	UNP Q84II6
C	387	HIS	-	expression tag	UNP Q84II6
C	388	HIS	-	expression tag	UNP Q84II6
C	389	HIS	-	expression tag	UNP Q84II6
C	390	HIS	-	expression tag	UNP Q84II6
C	391	HIS	-	expression tag	UNP Q84II6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	392	HIS	-	expression tag	UNP Q84II6

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
2	A	1	4	2	2	0	0
2	B	1	4	2	2	0	0
2	C	1	4	2	2	0	0

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0

- Molecule 4 is (9aR)-9a-(dioxidanyl)-1,9-dihydrocarbazole (CCD ID: EHX) (formula: C₁₂H₁₁NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	15	12	1	2	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	A	1	4	2	2	0	0
5	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	0
6	A	1	Total C O 7 4 3	0	0
6	A	1	Total C O 7 4 3	0	0
6	A	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	C	1	Total C O 7 4 3	0	0

- Molecule 7 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 10 6 4	0	0
7	A	1	Total C O 10 6 4	0	0
7	A	1	Total C O 10 6 4	0	0
7	B	1	Total C O 10 6 4	0	0
7	C	1	Total C O 10 6 4	0	0

- Molecule 8 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	O	S	0	0
			4	2	1	1		
8	B	1	Total	C	O	S	0	0
			4	2	1	1		
8	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 9 is 2'-amino[1,1'-biphenyl]-2,3-diol (CCD ID: 9G0) (formula: C₁₂H₁₁NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	N	O	0	0
			15	12	1	2		
9	C	1	Total	C	N	O	0	0
			15	12	1	2		

- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	1	Total	Mg	0	0
			1	1		

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	126	Total	O	0	0
			126	126		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	140	Total 140	O 140	0	0
11	C	112	Total 112	O 112	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	91.89Å 91.89Å 242.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.23 – 2.20 48.23 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.23-2.20) 97.9 (48.23-2.20)	Depositor EDS
R_{merge}	0.37	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.169 , 0.240 0.177 , 0.244	Depositor DCC
R_{free} test set	2969 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.055 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10038	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PEG, FE, 9G0, EDO, FES, MG, EHX, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/3297	0.94	1/4473 (0.0%)
1	B	0.79	0/3163	0.98	2/4294 (0.0%)
1	C	0.75	0/3180	0.97	0/4317
All	All	0.76	0/9640	0.96	3/13084 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	3
All	All	0	4

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	289	ILE	N-CA-C	-5.78	106.18	111.67
1	B	21	ALA	CA-C-N	5.17	127.20	120.28
1	B	21	ALA	C-N-CA	5.17	127.20	120.28

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	98	ARG	Sidechain
1	C	310	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	C	337	ARG	Sidechain
1	C	72	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3207	0	3114	72	0
1	B	3081	0	2992	39	0
1	C	3092	0	3009	32	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	15	0	0	0	0
5	A	20	0	30	1	0
5	B	40	0	60	8	0
5	C	20	0	30	4	0
6	A	28	0	40	2	0
6	B	42	0	60	4	0
6	C	7	0	10	0	0
7	A	30	0	42	2	0
7	B	10	0	14	0	0
7	C	10	0	14	1	0
8	A	4	0	6	0	0
8	B	8	0	12	2	0
9	B	15	0	0	1	0
9	C	15	0	0	2	0
10	B	1	0	0	0	0
11	A	126	0	0	1	0
11	B	140	0	0	2	0
11	C	112	0	0	1	0
All	All	10038	0	9433	141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204[B]:PHE:CD2	1:A:231[B]:ILE:HD12	1.42	1.53
1:A:204[B]:PHE:CE2	1:A:231[B]:ILE:HD12	1.62	1.33
1:A:204[B]:PHE:CD2	1:A:231[B]:ILE:CD1	2.24	1.20
1:A:204[B]:PHE:CE2	1:A:231[B]:ILE:CD1	2.28	1.15
1:A:204[B]:PHE:CG	1:A:231[B]:ILE:HD12	1.99	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	399/392 (102%)	360 (90%)	29 (7%)	10 (2%)	4 2
1	B	381/392 (97%)	363 (95%)	17 (4%)	1 (0%)	36 42
1	C	383/392 (98%)	364 (95%)	17 (4%)	2 (0%)	24 27
All	All	1163/1176 (99%)	1087 (94%)	63 (5%)	13 (1%)	16 10

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208[A]	GLY
1	A	208[B]	GLY
1	A	209[A]	ASP
1	A	209[B]	ASP
1	A	231[A]	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/339 (101%)	327 (96%)	15 (4%)	25	34
1	B	330/339 (97%)	323 (98%)	7 (2%)	47	63
1	C	332/339 (98%)	324 (98%)	8 (2%)	43	58
All	All	1004/1017 (99%)	974 (97%)	30 (3%)	39	49

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	383	SER
1	C	281	MET
1	B	220	ASP
1	C	356	PHE
1	C	191	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	315	GLN
1	C	330	ASN
1	C	213	GLN
1	C	234	HIS
1	C	282	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 49 ligands modelled in this entry, 4 are monoatomic - leaving 45 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FES	C	401	1	0,4,4	-	-	-		
5	EDO	C	405	-	3,3,3	0.33	0	2,2,2	0.49	0
2	FES	B	401	1	0,4,4	-	-	-		
5	EDO	A	404	-	3,3,3	0.44	0	2,2,2	0.49	0
5	EDO	B	413	-	3,3,3	0.42	0	2,2,2	0.26	0
6	PEG	A	409	-	6,6,6	0.61	0	5,5,5	0.57	0
5	EDO	A	408	-	3,3,3	0.43	0	2,2,2	0.18	0
5	EDO	C	404	-	3,3,3	0.29	0	2,2,2	0.51	0
6	PEG	B	417	-	6,6,6	0.65	0	5,5,5	0.89	0
5	EDO	B	407	-	3,3,3	0.44	0	2,2,2	0.46	0
5	EDO	A	407	-	3,3,3	0.41	0	2,2,2	0.62	0
7	PGE	A	415	-	9,9,9	0.59	0	8,8,8	0.35	0
6	PEG	A	412	-	6,6,6	0.62	0	5,5,5	0.35	0
8	DMS	B	423	-	3,3,3	0.54	0	3,3,3	0.80	0
6	PEG	A	410	-	6,6,6	0.49	0	5,5,5	0.38	0
8	DMS	B	422	-	3,3,3	0.67	0	3,3,3	0.84	0
5	EDO	C	407	-	3,3,3	0.37	0	2,2,2	0.36	0
5	EDO	B	408	-	3,3,3	0.40	0	2,2,2	0.39	0
4	EHX	A	403	3	14,17,17	4.09	5 (35%)	14,25,25	3.01	5 (35%)
5	EDO	A	406	-	3,3,3	0.30	0	2,2,2	0.61	0
5	EDO	B	411	-	3,3,3	0.50	0	2,2,2	0.40	0
6	PEG	A	411	-	6,6,6	0.51	0	5,5,5	0.24	0
7	PGE	A	413	-	9,9,9	0.48	0	8,8,8	0.37	0
6	PEG	B	416	-	6,6,6	0.43	0	5,5,5	0.28	0
7	PGE	B	421	-	9,9,9	0.51	0	8,8,8	0.56	0
7	PGE	C	410	-	9,9,9	0.46	0	8,8,8	0.47	0
5	EDO	B	406	-	3,3,3	0.34	0	2,2,2	0.31	0
8	DMS	A	416	-	3,3,3	0.43	0	3,3,3	0.89	0
5	EDO	B	405	-	3,3,3	0.52	0	2,2,2	0.42	0
9	9G0	B	403	3	16,16,16	2.18	4 (25%)	22,22,22	1.76	5 (22%)
5	EDO	B	410	-	3,3,3	0.18	0	2,2,2	0.52	0
5	EDO	C	408	-	3,3,3	0.71	0	2,2,2	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	409	-	3,3,3	0.50	0	2,2,2	0.33	0
5	EDO	B	414	-	3,3,3	0.57	0	2,2,2	0.22	0
5	EDO	C	406	-	3,3,3	0.46	0	2,2,2	0.50	0
6	PEG	B	418	-	6,6,6	0.38	0	5,5,5	0.80	0
5	EDO	B	412	-	3,3,3	0.65	0	2,2,2	0.26	0
6	PEG	B	420	-	6,6,6	0.55	0	5,5,5	0.51	0
6	PEG	C	409	-	6,6,6	0.40	0	5,5,5	0.60	0
6	PEG	B	419	-	6,6,6	0.55	0	5,5,5	0.75	0
9	9G0	C	403	3	16,16,16	2.31	4 (25%)	22,22,22	1.31	3 (13%)
2	FES	A	401	1	0,4,4	-	-	-	-	-
5	EDO	A	405	-	3,3,3	0.51	0	2,2,2	0.32	0
6	PEG	B	415	-	6,6,6	0.53	0	5,5,5	0.97	0
7	PGE	A	414	-	9,9,9	0.80	0	8,8,8	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FES	C	401	1	-	-	0/1/1/1
5	EDO	C	405	-	-	1/1/1/1	-
2	FES	B	401	1	-	-	0/1/1/1
5	EDO	A	404	-	-	0/1/1/1	-
5	EDO	B	413	-	-	1/1/1/1	-
6	PEG	A	409	-	-	4/4/4/4	-
5	EDO	A	408	-	-	1/1/1/1	-
5	EDO	C	404	-	-	0/1/1/1	-
6	PEG	B	417	-	-	1/4/4/4	-
5	EDO	B	407	-	-	0/1/1/1	-
5	EDO	A	407	-	-	1/1/1/1	-
7	PGE	A	415	-	-	4/7/7/7	-
6	PEG	A	412	-	-	3/4/4/4	-
6	PEG	A	410	-	-	2/4/4/4	-
5	EDO	C	407	-	-	1/1/1/1	-
5	EDO	B	408	-	-	1/1/1/1	-
4	EHX	A	403	3	-	0/0/26/26	0/3/3/3
5	EDO	A	406	-	-	1/1/1/1	-
5	EDO	B	411	-	-	0/1/1/1	-
6	PEG	A	411	-	-	1/4/4/4	-
7	PGE	A	413	-	-	5/7/7/7	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	B	416	-	-	2/4/4/4	-
7	PGE	B	421	-	-	3/7/7/7	-
7	PGE	C	410	-	-	4/7/7/7	-
5	EDO	B	406	-	-	1/1/1/1	-
5	EDO	B	405	-	-	0/1/1/1	-
9	9G0	B	403	3	-	0/4/4/4	0/2/2/2
5	EDO	B	410	-	-	1/1/1/1	-
5	EDO	C	408	-	-	1/1/1/1	-
5	EDO	B	409	-	-	1/1/1/1	-
5	EDO	B	414	-	-	1/1/1/1	-
5	EDO	C	406	-	-	0/1/1/1	-
6	PEG	B	418	-	-	2/4/4/4	-
5	EDO	B	412	-	-	1/1/1/1	-
6	PEG	B	420	-	-	1/4/4/4	-
6	PEG	C	409	-	-	1/4/4/4	-
6	PEG	B	419	-	-	2/4/4/4	-
9	9G0	C	403	3	-	0/4/4/4	0/2/2/2
2	FES	A	401	1	-	-	0/1/1/1
5	EDO	A	405	-	-	0/1/1/1	-
6	PEG	B	415	-	-	4/4/4/4	-
7	PGE	A	414	-	-	5/7/7/7	-

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	EHX	C4-C4A	12.21	1.47	1.35
4	A	403	EHX	C4B-C8A	7.53	1.50	1.41
9	C	403	9G0	CA1-CA2	5.65	1.50	1.40
9	C	403	9G0	CA3-CA2	5.06	1.45	1.40
9	B	403	9G0	CA1-CA2	5.03	1.49	1.40

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	EHX	C4B-C8A-N9	-7.48	108.45	111.82
4	A	403	EHX	C8-C8A-C4B	-4.79	117.55	122.19
4	A	403	EHX	C8A-C4B-C4A	-4.21	104.23	106.58
9	B	403	9G0	CA6-CA1-CB1	4.06	126.94	118.74
9	B	403	9G0	CB1-CA1-CA2	-3.99	115.25	121.80

There are no chirality outliers.

5 of 57 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	415	PEG	C4-C3-O2-C2
7	A	415	PGE	O2-C3-C4-O3
7	B	421	PGE	O2-C3-C4-O3
7	A	414	PGE	O2-C3-C4-O3
6	A	410	PEG	O2-C3-C4-O4

There are no ring outliers.

14 monomers are involved in 27 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	408	EDO	1	0
5	C	404	EDO	3	0
6	B	417	PEG	1	0
7	A	415	PGE	2	0
6	A	412	PEG	1	0
8	B	423	DMS	2	0
6	A	410	PEG	1	0
5	C	407	EDO	1	0
7	C	410	PGE	1	0
9	B	403	9G0	1	0
5	B	410	EDO	4	0
5	B	414	EDO	4	0
6	B	419	PEG	3	0
9	C	403	9G0	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/392 (97%)	-0.07	9 (2%) 61 58	16, 31, 60, 97	18 (4%)
1	B	383/392 (97%)	-0.33	1 (0%) 90 88	20, 29, 53, 83	0
1	C	383/392 (97%)	-0.07	6 (1%) 70 67	22, 35, 59, 79	2 (0%)
All	All	1149/1176 (97%)	-0.16	16 (1%) 73 71	16, 32, 59, 97	20 (1%)

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	210[A]	ARG	3.2
1	A	335	TRP	2.9
1	C	335	TRP	2.8
1	A	230[A]	LEU	2.7
1	A	235	GLY	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	C	408	4/4	0.74	0.16	48,51,51,52	0
5	EDO	B	405	4/4	0.78	0.18	57,60,62,63	0
7	PGE	A	414	10/10	0.78	0.17	50,59,63,65	0
8	DMS	A	416	4/4	0.79	0.23	65,66,68,70	0
5	EDO	B	411	4/4	0.81	0.14	53,53,53,56	0
6	PEG	B	415	7/7	0.81	0.16	34,42,50,53	0
6	PEG	A	410	7/7	0.82	0.15	57,58,66,66	0
7	PGE	A	413	10/10	0.82	0.16	62,65,68,69	0
5	EDO	B	409	4/4	0.83	0.14	60,61,62,64	0
6	PEG	A	412	7/7	0.83	0.14	54,59,61,62	0
5	EDO	A	407	4/4	0.84	0.16	54,57,58,62	0
5	EDO	B	414	4/4	0.84	0.15	46,49,49,50	0
6	PEG	A	411	7/7	0.84	0.15	59,63,66,66	0
5	EDO	C	406	4/4	0.84	0.16	50,53,57,60	0
6	PEG	B	420	7/7	0.85	0.13	46,55,61,63	0
6	PEG	B	418	7/7	0.85	0.15	37,42,45,47	0
5	EDO	B	410	4/4	0.87	0.22	38,42,42,52	0
6	PEG	B	419	7/7	0.87	0.13	39,44,50,55	0
5	EDO	C	404	4/4	0.87	0.19	39,41,41,42	0
5	EDO	A	408	4/4	0.87	0.12	54,55,55,57	0
5	EDO	B	412	4/4	0.87	0.15	40,47,48,53	0
7	PGE	A	415	10/10	0.87	0.11	55,68,71,72	0
6	PEG	B	417	7/7	0.87	0.15	25,36,43,46	0
8	DMS	B	423	4/4	0.87	0.18	76,76,79,80	0
5	EDO	B	407	4/4	0.88	0.13	39,49,51,54	0
7	PGE	C	410	10/10	0.88	0.13	44,54,62,66	0
4	EHX	A	403	15/15	0.89	0.12	43,51,63,64	0
6	PEG	A	409	7/7	0.89	0.12	43,47,54,55	0
9	9G0	B	403	15/15	0.89	0.15	42,56,61,63	0
9	9G0	C	403	15/15	0.89	0.14	48,54,62,65	0
6	PEG	C	409	7/7	0.90	0.13	48,51,56,56	0
7	PGE	B	421	10/10	0.91	0.10	40,52,59,60	0
5	EDO	B	408	4/4	0.91	0.10	43,44,46,48	0
5	EDO	A	405	4/4	0.91	0.14	34,40,41,42	0
5	EDO	A	404	4/4	0.92	0.09	40,40,40,40	0
5	EDO	B	413	4/4	0.93	0.10	38,46,48,53	0
6	PEG	B	416	7/7	0.93	0.12	45,46,49,53	0
5	EDO	A	406	4/4	0.93	0.11	46,50,53,59	0
5	EDO	B	406	4/4	0.93	0.09	43,43,43,44	0
5	EDO	C	407	4/4	0.94	0.10	41,45,47,53	0
5	EDO	C	405	4/4	0.94	0.12	55,57,57,60	0
8	DMS	B	422	4/4	0.98	0.07	37,37,37,39	0
3	FE	A	402	1/1	0.98	0.03	46,46,46,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FES	A	401	4/4	0.99	0.02	20,20,21,21	0
3	FE	B	402	1/1	0.99	0.02	23,23,23,23	0
3	FE	C	402	1/1	0.99	0.01	32,32,32,32	0
2	FES	B	401	4/4	0.99	0.02	25,26,28,29	0
2	FES	C	401	4/4	0.99	0.04	32,33,34,34	0
10	MG	B	404	1/1	0.99	0.03	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.