



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 07:41 PM EDT

PDB ID : 3LMH / pdb_00003lmh
Title : Crystal Structure of the Alpha-kinase Domain of Myosin Heavy Chain Kinase A Complex with ADP
Authors : Ye, Q.; Jia, Z.
Deposited on : 2010-01-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	
1	B	307	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin heavy chain kinase A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	P	S	0	0	0
			1978	1270	329	366	1	12			
1	B	240	Total	C	N	O	P	S	0	0	0
			1915	1231	317	354	1	12			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	535	MET	-	expression tag	UNP P42527
A	536	GLY	-	expression tag	UNP P42527
A	537	GLY	-	expression tag	UNP P42527
A	538	HIS	-	expression tag	UNP P42527
A	539	HIS	-	expression tag	UNP P42527
A	540	HIS	-	expression tag	UNP P42527
A	541	HIS	-	expression tag	UNP P42527
A	542	HIS	-	expression tag	UNP P42527
A	543	HIS	-	expression tag	UNP P42527
A	544	GLY	-	expression tag	UNP P42527
A	545	GLU	-	expression tag	UNP P42527
A	546	ASN	-	expression tag	UNP P42527
A	547	LEU	-	expression tag	UNP P42527
A	548	TYR	-	expression tag	UNP P42527
A	549	PHE	-	expression tag	UNP P42527
A	550	GLN	-	expression tag	UNP P42527
A	551	GLY	-	expression tag	UNP P42527
B	535	MET	-	expression tag	UNP P42527
B	536	GLY	-	expression tag	UNP P42527
B	537	GLY	-	expression tag	UNP P42527
B	538	HIS	-	expression tag	UNP P42527
B	539	HIS	-	expression tag	UNP P42527
B	540	HIS	-	expression tag	UNP P42527
B	541	HIS	-	expression tag	UNP P42527
B	542	HIS	-	expression tag	UNP P42527

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	543	HIS	-	expression tag	UNP P42527
B	544	GLY	-	expression tag	UNP P42527
B	545	GLU	-	expression tag	UNP P42527
B	546	ASN	-	expression tag	UNP P42527
B	547	LEU	-	expression tag	UNP P42527
B	548	TYR	-	expression tag	UNP P42527
B	549	PHE	-	expression tag	UNP P42527
B	550	GLN	-	expression tag	UNP P42527
B	551	GLY	-	expression tag	UNP P42527

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	204	Total	O	0	0
			204	204		
5	B	164	Total	O	0	0
			164	164		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.82Å 110.05Å 78.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.70 – 2.00 19.70 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.70-2.00) 99.6 (19.70-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.63 (at 2.01Å)	Xtriage
Refinement program	REFMAC 6.1.0	Depositor
R, R_{free}	0.216 , 0.259 0.215 , 0.258	Depositor DCC
R_{free} test set	2484 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	29.5	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4327	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4, ADP, PHD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2006	0.81	0/2701
1	B	0.57	0/1943	0.79	0/2618
All	All	0.58	0/3949	0.80	0/5319

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1978	0	1993	41	0
1	B	1915	0	1929	39	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	204	0	0	10	0
5	B	164	0	0	9	0
All	All	4327	0	3946	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:743:PHE:CZ	1:B:747:LEU:HD12	2.00	0.95
1:B:659:LEU:HD11	5:B:94:HOH:O	1.73	0.87
1:B:743:PHE:CZ	1:B:747:LEU:CD1	2.61	0.83
1:A:770:HIS:HD2	1:A:782:LEU:H	1.28	0.81
1:A:770:HIS:CD2	1:A:782:LEU:H	1.99	0.80
1:A:671:ARG:NE	5:A:281:HOH:O	2.13	0.75
1:A:730:VAL:HG23	1:A:793:THR:CG2	2.19	0.73
1:A:706:GLN:HB3	1:A:707:PRO:HD2	1.74	0.70
1:A:728:GLY:O	1:A:793:THR:HG21	1.95	0.66
1:A:671:ARG:NH1	5:A:25:HOH:O	2.28	0.66
1:A:743:PHE:CZ	1:A:747:LEU:HD22	2.33	0.63
1:B:773:ASP:HB2	5:B:81:HOH:O	1.98	0.63
1:A:771:THR:O	1:A:783:GLY:HA2	1.98	0.63
1:A:706:GLN:CB	1:A:707:PRO:HD2	2.29	0.62
1:A:648:LYS:O	1:A:649:LYS:HB2	1.99	0.62
1:A:766:PHD:P	5:A:283:HOH:O	2.57	0.61
1:B:743:PHE:CE2	1:B:747:LEU:HD12	2.36	0.60
1:B:743:PHE:CE1	1:B:747:LEU:CD1	2.84	0.60
1:A:671:ARG:NH1	1:A:690:MET:HG2	2.16	0.60
1:A:581:VAL:HG11	1:A:699:ARG:NH2	2.17	0.59
1:A:552:ILE:HG22	1:A:601:GLY:HA3	1.85	0.58
1:B:741:SER:OG	1:B:757:ILE:CD1	2.51	0.58
1:B:592:ARG:NH1	5:B:282:HOH:O	2.35	0.58
1:B:743:PHE:CE1	1:B:747:LEU:HD12	2.40	0.55
1:A:728:GLY:HA3	1:A:789:LYS:HG3	1.89	0.54
1:B:616:LEU:N	1:B:618:PRO:O	2.41	0.54
1:A:730:VAL:HG23	1:A:793:THR:HG21	1.90	0.54
1:B:727:TYR:CE1	1:B:789:LYS:HG3	2.42	0.53
1:A:656:SER:OG	1:A:659:LEU:HD13	2.08	0.53
1:B:743:PHE:CE1	1:B:747:LEU:HD11	2.44	0.53
1:B:655:ALA:HA	1:B:659:LEU:HD23	1.90	0.52
1:A:571:LYS:HE3	5:A:143:HOH:O	2.09	0.52
1:B:755:VAL:HA	1:B:757:ILE:HD12	1.91	0.52
1:A:552:ILE:N	5:A:255:HOH:O	2.42	0.51
1:B:773:ASP:OD2	1:B:775:LYS:HB2	2.10	0.51
1:B:727:TYR:CZ	1:B:789:LYS:HG3	2.45	0.51
1:A:745:TYR:CZ	1:A:750:LYS:HG2	2.45	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:671:ARG:NH1	5:B:42:HOH:O	2.43	0.51
1:B:741:SER:OG	1:B:757:ILE:HD13	2.12	0.49
1:B:771:THR:O	1:B:783:GLY:HA2	2.12	0.49
1:A:617:PHE:HA	1:A:618:PRO:C	2.37	0.49
1:B:659:LEU:HB2	5:B:296:HOH:O	2.12	0.49
1:A:689:LEU:HB3	1:A:714:PRO:HD2	1.95	0.48
1:A:581:VAL:HG11	1:A:699:ARG:HH22	1.78	0.48
1:B:676:LYS:HE3	1:B:747:LEU:HD13	1.95	0.47
1:A:706:GLN:OE1	1:A:706:GLN:N	2.47	0.47
1:A:745:TYR:CE1	1:A:750:LYS:HG2	2.50	0.47
1:A:706:GLN:CB	1:A:707:PRO:CD	2.92	0.47
1:A:726:ASN:O	1:A:770:HIS:CE1	2.68	0.47
1:A:610:LEU:HD22	1:A:616:LEU:HD11	1.96	0.47
1:B:755:VAL:HA	1:B:757:ILE:CD1	2.44	0.47
1:A:766:PHD:P	5:A:12:HOH:O	2.72	0.46
1:B:773:ASP:HB3	1:B:775:LYS:H	1.80	0.46
1:A:766:PHD:OP1	5:A:12:HOH:O	2.20	0.46
1:B:741:SER:OG	1:B:757:ILE:HD11	2.15	0.46
1:B:794:HIS:HD2	5:B:30:HOH:O	1.98	0.45
1:A:671:ARG:HH12	1:A:690:MET:HG2	1.80	0.45
1:A:581:VAL:HG12	5:A:15:HOH:O	2.16	0.45
1:A:677:PHE:CE2	1:A:682:PRO:HG3	2.52	0.45
1:B:583:ARG:NH1	1:B:584:LYS:HE3	2.32	0.44
1:B:671:ARG:NH2	5:B:42:HOH:O	2.50	0.44
1:A:682:PRO:HB2	5:A:20:HOH:O	2.16	0.44
1:B:583:ARG:HH12	1:B:584:LYS:HE3	1.83	0.43
1:B:743:PHE:CZ	1:B:747:LEU:HD11	2.48	0.43
1:A:730:VAL:HG23	1:A:793:THR:HG22	2.01	0.43
1:A:730:VAL:CG2	1:A:793:THR:CG2	2.93	0.43
1:B:617:PHE:CG	1:B:618:PRO:HA	2.54	0.42
1:A:762:ASP:OD1	1:A:764:TYR:OH	2.20	0.42
1:B:563:TRP:CH2	1:B:574:ARG:HD2	2.55	0.42
1:A:626:ILE:HD13	1:B:750:LYS:HD2	2.01	0.42
1:B:616:LEU:N	1:B:617:PHE:C	2.78	0.41
1:B:806:ASP:O	1:B:807:VAL:HB	2.21	0.41
1:B:742:HIS:HE1	5:B:66:HOH:O	2.03	0.41
1:A:706:GLN:HB3	1:A:707:PRO:CD	2.47	0.41
1:A:794:HIS:HD2	5:A:51:HOH:O	2.04	0.41
1:B:555:GLU:HB2	1:B:600:LEU:CD2	2.51	0.41
1:B:571:LYS:HE3	5:B:106:HOH:O	2.21	0.40
1:B:686:ILE:HG22	1:B:762:ASP:HB3	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:724:ASN:HD21	1:B:757:ILE:HD13	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/307 (78%)	234 (98%)	3 (1%)	2 (1%)	16	11
1	B	231/307 (75%)	226 (98%)	5 (2%)	0	100	100
All	All	470/614 (76%)	460 (98%)	8 (2%)	2 (0%)	30	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	649	LYS
1	A	706	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/269 (81%)	217 (99%)	2 (1%)	70	78
1	B	213/269 (79%)	208 (98%)	5 (2%)	44	49
All	All	432/538 (80%)	425 (98%)	7 (2%)	55	62

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	552	ILE
1	A	793	THR
1	B	552	ILE
1	B	556	THR
1	B	616	LEU
1	B	773	ASP
1	B	793	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	607	ASN
1	A	654	GLN
1	A	675	ASN
1	A	770	HIS
1	A	794	HIS
1	B	607	ASN
1	B	742	HIS
1	B	794	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PHD	B	766	1	9,11,12	0.98	1 (11%)	9,15,17	1.86	2 (22%)
1	PHD	A	766	1	9,11,12	1.27	1 (11%)	9,15,17	2.11	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PHD	B	766	1	-	2/8/11/13	-
1	PHD	A	766	1	-	1/8/11/13	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	766	PHD	P-OD1	3.38	1.65	1.59
1	B	766	PHD	P-OD1	2.21	1.63	1.59

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	766	PHD	OD1-CG-CB	4.75	122.10	110.95
1	B	766	PHD	OD1-CG-CB	4.16	120.71	110.95
1	A	766	PHD	OD2-CG-CB	-3.27	117.06	124.65
1	B	766	PHD	OD2-CG-CB	-2.63	118.53	124.65

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	766	PHD	CA-CB-CG-OD1
1	B	766	PHD	CA-CB-CG-OD2
1	A	766	PHD	CA-CB-CG-OD2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	766	PHD	3	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	A	1511	-	28,29,29	1.69	5 (17%)	43,45,45	1.78	7 (16%)
2	ADP	B	1512	-	28,29,29	1.49	6 (21%)	43,45,45	1.73	9 (20%)
4	PO4	A	842	-	4,4,4	1.17	0	6,6,6	1.34	1 (16%)
4	PO4	B	842	-	4,4,4	0.98	0	6,6,6	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1511	-	-	3/16/32/32	0/3/3/3
2	ADP	B	1512	-	-	0/16/32/32	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1511	ADP	C5-C4	5.04	1.48	1.39
2	B	1512	ADP	C5-C4	4.89	1.47	1.39
2	A	1511	ADP	PA-O3A	4.64	1.64	1.59
2	B	1512	ADP	PA-O3A	3.05	1.62	1.59
2	A	1511	ADP	C5-C6	2.65	1.48	1.41
2	B	1512	ADP	C5-C6	2.57	1.48	1.41
2	A	1511	ADP	C8-N7	2.47	1.36	1.31
2	A	1511	ADP	C5-N7	-2.43	1.34	1.39
2	B	1512	ADP	C5-N7	-2.39	1.34	1.39
2	B	1512	ADP	C8-N7	2.20	1.35	1.31
2	B	1512	ADP	C4-N9	-2.09	1.33	1.37

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1511	ADP	C5-C4-N3	-6.05	118.39	126.72
2	B	1512	ADP	C5-C4-N3	-5.46	119.20	126.72
2	A	1511	ADP	N3-C4-N9	4.64	135.06	127.17
2	B	1512	ADP	N3-C4-N9	4.19	134.29	127.17
2	A	1511	ADP	C2-N3-C4	3.77	121.05	111.83
2	B	1512	ADP	C4-C5-N7	-3.70	106.35	110.58
2	B	1512	ADP	C2-N3-C4	3.53	120.45	111.83
2	A	1511	ADP	N3-C2-N1	-3.32	123.56	128.58
2	A	1511	ADP	C4-C5-N7	-3.20	106.92	110.58
2	B	1512	ADP	N3-C2-N1	-3.14	123.83	128.58
2	B	1512	ADP	C5-N7-C8	2.73	107.75	103.45
2	B	1512	ADP	C4-N9-C8	2.57	108.43	105.74
2	B	1512	ADP	C6-C5-N7	2.46	136.83	132.09
4	A	842	PO4	O4-P-O2	2.33	115.15	107.91
2	A	1511	ADP	C5-N7-C8	2.25	106.99	103.45
2	B	1512	ADP	N9-C8-N7	-2.10	110.96	113.94
2	A	1511	ADP	C2-N1-C6	2.08	122.15	118.73

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1511	ADP	PA-O3A-PB-O3B
2	A	1511	ADP	PA-O3A-PB-O1B
2	A	1511	ADP	PA-O3A-PB-O2B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/307 (80%)	0.26	12 (4%) 35 34	15, 28, 45, 57	0
1	B	239/307 (77%)	0.56	19 (7%) 18 17	19, 33, 46, 63	0
All	All	486/614 (79%)	0.41	31 (6%) 25 24	15, 30, 46, 63	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	796	CYS	9.4
1	A	705	GLY	6.4
1	B	552	ILE	6.4
1	B	554	SER	6.3
1	B	616	LEU	5.2
1	A	552	ILE	4.7
1	B	655	ALA	4.2
1	B	807	VAL	4.0
1	B	804	ASP	3.5
1	A	809	LEU	3.3
1	B	806	ASP	3.3
1	B	556	THR	3.0
1	B	707	PRO	2.9
1	A	706	GLN	2.8
1	B	553	SER	2.6
1	B	555	GLU	2.6
1	B	700	SER	2.6
1	B	634	THR	2.5
1	A	704	ASN	2.5
1	B	773	ASP	2.5
1	B	792	THR	2.5
1	B	793	THR	2.4
1	A	649	LYS	2.4
1	A	650	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	793	THR	2.3
1	A	658	GLU	2.3
1	A	804	ASP	2.3
1	B	801	ALA	2.1
1	A	791	ILE	2.1
1	B	583	ARG	2.1
1	A	555	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PHD	A	766	12/13	0.80	0.14	19,27,38,40	0
1	PHD	B	766	12/13	0.80	0.17	26,34,46,46	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ADP	A	1511	27/27	0.87	0.13	27,38,55,55	0
4	PO4	B	842	5/5	0.90	0.13	42,42,46,47	0
2	ADP	B	1512	27/27	0.91	0.10	30,42,60,61	0
4	PO4	A	842	5/5	0.92	0.10	35,35,37,38	0
3	ZN	A	1	1/1	0.97	0.03	27,27,27,27	0
3	ZN	B	2	1/1	0.98	0.03	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.




Electron density around ADP A 1511:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ADP B 1512:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.