



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:26 PM UTC

PDB ID : 4LMQ / pdb_00004lmq
Title : Development and Preclinical Characterization of a Humanized Antibody Targeting CXCL12
Authors : Zhong, Z.; Wang, J.; Li, B.; Xiang, H.; Ultsch, M.; Coons, M.; Wong, T.; Chiang, N.Y.; Clark, S.; Clark, R.; Quintana, L.; Gribling, P.; Suto, E.; Barck, K.; Corpuz, R.; Yao, J.; Takkar, R.; Lee, W.P.; Damico-Beyer, L.A.; Carano, R.D.; Adams, C.; Kelley, R.F.; Wang, W.; Ferrara, N.
Deposited on : 2013-07-10
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	61	15% 59% 28% • 10%
1	F	61	8% 54% 34% 5% • 5%
2	E	221	22% 66% 19% • 12%
2	H	221	6% 86% 13% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	214	 7% 77% 21% •
3	L	214	 2% 80% 19%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7398 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Stromal cell-derived factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	55	Total	C	N	O	S	0	0	0
			449	286	85	76	2			
1	F	58	Total	C	N	O	S	0	0	0
			472	300	90	78	4			

- Molecule 2 is a protein called hu30D8 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	195	Total	C	N	O	S	0	0	0
			1474	936	248	285	5			
2	H	220	Total	C	N	O	S	0	0	0
			1636	1031	276	324	5			

- Molecule 3 is a protein called hu30D8 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	214	Total	C	N	O	S	0	0	0
			1654	1039	274	335	6			
3	L	214	Total	C	N	O	S	0	0	0
			1654	1039	274	335	6			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	8	Total	O	0	0
			8	8		
4	E	10	Total	O	0	0
			10	10		
4	F	7	Total	O	0	0
			7	7		
4	H	11	Total	O	0	0
			11	11		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	I	11	Total	O	0	0
			11	11		
4	L	12	Total	O	0	0
			12	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Stromal cell-derived factor 1




- Molecule 1: Stromal cell-derived factor 1




- Molecule 2: hu30D8 Fab heavy chain





- Molecule 2: hu30D8 Fab heavy chain







- Molecule 3: hu30D8 Fab light chain



- Molecule 3: hu30D8 Fab light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.95Å 73.10Å 106.63Å 90.00° 90.40° 90.00°	Depositor
Resolution (Å)	55.30 – 2.77 55.30 – 2.77	Depositor EDS
% Data completeness (in resolution range)	94.2 (55.30-2.77) 98.0 (55.30-2.77)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.253 , 0.301 0.225 , 0.241	Depositor DCC
R_{free} test set	1251 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtriage
Anisotropy	0.226	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7398	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.29	0/455	0.67	0/610
1	F	0.35	0/481	0.83	0/649
2	E	0.32	0/1508	0.76	1/2051 (0.0%)
2	H	0.30	0/1675	0.70	1/2284 (0.0%)
3	I	0.31	0/1691	0.72	2/2295 (0.1%)
3	L	0.30	0/1691	0.72	1/2295 (0.0%)
All	All	0.31	0/7501	0.73	5/10184 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	E	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	54	SER	N-CA-C	7.45	119.47	111.36
2	H	128	SER	N-CA-C	5.36	116.80	111.07
3	L	30	SER	CB-CA-C	-5.11	110.28	117.23
3	I	112	ALA	CA-C-N	5.07	124.83	119.76
3	I	112	ALA	C-N-CA	5.07	124.83	119.76

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	E	146	PHE	Peptide
2	E	148	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	449	0	467	12	0
1	F	472	0	484	16	0
2	E	1474	0	1434	21	0
2	H	1636	0	1599	12	0
3	I	1654	0	1609	22	0
3	L	1654	0	1609	15	0
4	D	8	0	0	1	0
4	E	10	0	0	0	0
4	F	7	0	0	1	0
4	H	11	0	0	0	0
4	I	11	0	0	0	0
4	L	12	0	0	0	0
All	All	7398	0	7202	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:16:SER:H	1:F:17:HIS:HA	1.45	0.81
3:I:197:THR:H	3:I:204:PRO:HB3	1.50	0.76
1:F:15:GLU:HB3	1:F:16:SER:HB2	1.67	0.75
2:E:34:VAL:HG11	2:E:78:VAL:HG21	1.68	0.73
2:E:51:LEU:HD11	2:E:71:ARG:HE	1.53	0.72

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	51/61 (84%)	47 (92%)	2 (4%)	2 (4%)	2	7
1	F	56/61 (92%)	51 (91%)	3 (5%)	2 (4%)	2	8
2	E	187/221 (85%)	172 (92%)	9 (5%)	6 (3%)	3	10
2	H	218/221 (99%)	205 (94%)	11 (5%)	2 (1%)	14	37
3	I	212/214 (99%)	198 (93%)	10 (5%)	4 (2%)	6	19
3	L	212/214 (99%)	202 (95%)	9 (4%)	1 (0%)	24	52
All	All	936/992 (94%)	875 (94%)	44 (5%)	17 (2%)	6	21

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	147	PRO
1	F	12	ARG
3	I	196	VAL
3	I	197	THR
1	D	12	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	49/56 (88%)	45 (92%)	4 (8%)	10	30
1	F	52/56 (93%)	44 (85%)	8 (15%)	2	8
2	E	163/184 (89%)	150 (92%)	13 (8%)	11	31
2	H	183/184 (100%)	175 (96%)	8 (4%)	25	56
3	I	190/190 (100%)	178 (94%)	12 (6%)	16	41
3	L	190/190 (100%)	177 (93%)	13 (7%)	14	38
All	All	827/860 (96%)	769 (93%)	58 (7%)	14	37

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	131	THR
3	L	201	LEU
3	I	18	ARG
3	L	181	LEU
3	L	132	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	39	GLN
2	H	164	HIS
3	L	38	GLN
3	I	50	ASN
1	F	30	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	55/61 (90%)	0.64	9 (16%) 4 4	15, 41, 88, 116	0
1	F	58/61 (95%)	0.51	5 (8%) 16 13	9, 39, 75, 104	0
2	E	195/221 (88%)	1.06	49 (25%) 1 1	10, 60, 109, 132	0
2	H	220/221 (99%)	0.27	14 (6%) 25 20	8, 32, 81, 142	0
3	I	214/214 (100%)	0.56	14 (6%) 25 20	13, 45, 97, 124	0
3	L	214/214 (100%)	-0.03	4 (1%) 66 59	7, 30, 67, 113	0
All	All	956/992 (96%)	0.47	95 (9%) 13 10	7, 36, 97, 142	0

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	16	SER	4.9
2	H	132	SER	4.5
1	F	11	CYS	3.7
2	E	205	THR	3.7
2	E	114	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.